


3.4. CANADA

Figure 3.4.1. **National agri-environmental and economic profile, 2002-04: Canada**



StatLink  <http://dx.doi.org/10.1787/288861614413>

1. Data refer to the year 1996.

2. Data refer to the year 1995.

3. Data refer to the year 2004.

Source: OECD Secretariat. For full details of these indicators, see Chapter 1 of the *Main Report*.

3.4.1. Agricultural sector trends and policy context

Growth in agricultural production was more than double the OECD average between 1990-92 and 2002-04, owing in part to recent strong growth in production and sales in the pig and horticultural sectors (Figure 3.4.2). Farming's contribution to the economy accounts for around 2% of employment and 1% of GDP, while the whole agriculture and agri-food system accounts for approximately 13% of employment and 8% of GDP [1] (Figure 3.4.1). Canada is a major world exporter of cereals, oilseeds, animals and red meats (around 3% of world farm export value), with nearly 25% of production exported in 2004 [1, 2].

Agricultural production is intensifying and concentrated in fewer farms [1, 3]. Farm size and intensity varies across Canada depending on commodity specialisation, geography and land availability. The range of climates, soil types, resource availability, population distribution and competing land uses across the country allows some regions to implement more intensive management practices than others, including higher uses of inputs such as fertiliser, pesticides, energy and water (Figure 3.4.2), and higher densities of livestock. The result has been a greater rise in annual multifactor productivity growth for the agriculture sector (3%) than for industrial sectors (1.5%) over the period 1997 to 2003 [1]. The increase in intensity began in the 1940s, in part, due to economies of scale associated with a change to more capital-intensive technologies, with both farm and herd size increasing ever since [1]. This is reflected in that only one-third of farms report sales over CAD 100 000 (USD 76 000) but account for nearly 90% of farm production [1].

Agricultural support has declined. Support to farmers (as measured by the OECD Producer Support Estimate – PSE) fell from 36% to 22% of farm receipts between 1986-88 and 2002-04, compared to the performance of the OECD area where the average decreased from 37% to 30%. The share of output and input linked support also fell from 82% in 1986-88 to 57% of the PSE in 2002-04 [4]. The 2003-08 *Agricultural Policy Framework* (APF) provides Federal, Provincial and Territorial support to the farm sector through various programmes that fall under the headings of: business risk management; food safety and quality; environment; science and innovation; and renewal. Total agricultural expenditure was CAD 10 (USD 7) billion annually over 2002-04, or just under 1% of GDP [4].

There is growing emphasis on the environment in agricultural policy. Over the 1990s much of the focus of agricultural policy was on economic and production objectives, but environmental considerations became a key part of the APF [5]. Agriculture and environment are shared responsibilities between Federal, Provincial and Territorial governments. Most APF agri-environmental programmes are cost-shared between the Federal, Provincial and Territorial Governments, with CAD 700 (USD 490) million of funding over 5 years (2004-08) provided by the Federal Government [4, 6]. Programmes under the environment chapter of the APF provide producers with assistance to improve their environmental management of soil, water, land and biodiversity by cost-sharing activities such as technical assistance, extension, research and demonstration activities, although there are some exceptions [7, 8].

Several national environmental and taxation policies impact agriculture. The *Canadian Environmental Protection Act* addresses air pollution and toxic substances and involves the agriculture sector when developing risk management plans for listed substances. The Pest Management Regulatory Authority monitors and regulates pesticide products and their use under the *Pest Control Products Act*. Farmers are supported with an on-farm fuel tax exemption, equal to CAD 285 (USD 200) million annually during 2002-04 [9]. Some farm inputs (e.g. fertilisers, pesticides) are exempt from the Federal Goods and Service Tax [9]. Irrigation water charges have risen from CAD 11 to 31 (USD 7 to 22) per m³ between the 1980s and 2000 [3]. *Integrated Water Resources Management* is being used to bring together Federal, Provincial and Municipal authorities in the planning and management of water policies [3, 10]. Biofuels are exempt from the Federal excise taxes on transport fuels [4].

Producers are also affected by commitments under several international environmental agreements. Under the *North America Free Trade Agreement* Canada, together with Mexico and the United States, is seeking greater harmonisation of pesticide regulations [2]. In eastern Canada producers are impacted by commitments made under the *Great Lakes Water Quality Agreement* with the US, co-ordinated through the International Joint Commission, which addresses concerns related to agricultural water pollution and water withdrawals for irrigation and other uses [3, 11]. A plan was initiated in 1997 to improve biodiversity conservation in agriculture as part of Canada's commitments under the *Convention on Biological Diversity*. [3]. Canada is a signatory to the *Kyoto Protocol* to address greenhouse gas emissions, the *Gothenburg Protocol* to reduce ammonia emissions (although emission targets have not yet been determined), and the *Montreal Protocol* to phase out ozone depleting substances, including methyl bromide.

3.4.2. The environmental performance of agriculture

The key environmental challenges concerning agriculture include soil, water and air quality.

The growing agricultural demand for water and the impact of farming on biodiversity are also important issues. There are a number of environmental concerns between farming and urban communities [12, 13], notably odours from livestock operations, and the conversion of farmland to urban use [14]. Canada is the second largest country by area in the world, but climate, topography and the range of soil types limit the land suitable for agriculture to approximately 7% (2002-04) of the total land area [15]. Between 1990-92 and 2002-04 the total area of farmland decreased by over 2%, largely because the land suitable for agriculture is already being used for that purpose (Figure 3.4.2). Approximately 60% of farmland is cultivated, 30% pasture and 10% used for other purposes (e.g. woodlots). The increase in cropped land is primarily due to the reduction in the use of summerfallow in rotations. Summerfallow area decreased by more than half between 1981 and 2001. The more intensive use of cropland is a result of the adoption of management practices that allow for continuous cropping or extended crop rotations [2].

Overall soil quality – erosion, soil organic carbon, salinity – has improved, during the period 1991 to 2001. Improvements include: an increase in the share of cropland under vegetative cover for more than 300 days annually; a higher share of cropland in low erosion (water, wind and tillage soil erosion) and salinisation risk classes; and a net accumulation of soil organic carbon in cropland since 1996 (Figure 3.4.2) [2, 16, 17, 18]. These developments are a result of: increased adoption of reduced tillage or no-till practices, rising from around 30% in 1991 to 60% of cropland in 2001; reduced use of summerfallow; and expansion in the area of perennial vegetation which primarily involves the conversion of marginal cropland to forage production. There is still room for improvement, however. Approximately 4% of cropland considered to be at high risk for soil degradation (erosion and salinity) was still under cultivation in 2001. In 2006 about 28% of agricultural land in Canada remains under conventional tillage practices, with a higher share in the Atlantic Provinces and Québec, largely due to crop type and climate, and 30% of cropland is still considered to be in the low soil cover class (especially in Ontario and Saskatchewan) [2].

Water contamination from agricultural sources is a concern and risk of water contamination from agriculture has increased since 1981 [3, 19, 20, 21, 22, 23, 24, 25]. Agriculture is a key source of nitrogen and phosphorus in the environment, although risk of contamination tends to be localised [26]. The increase in nutrient surpluses is reflected in the rising trend in the Indicator of Risk of Water Contamination by Nitrogen (IROWC-N) [2]. For instance, the share of farmland in the high to very high risk category for IROWC-N rose from 11% in 1991 to 16% by 2001, and was about 50% in certain regions [2]. Some regions in Canada are at higher risk of poor water quality than others, owing to: surrounding land uses; population density; increased use of inputs, such as fertilisers; and climatic conditions of heavy annual or seasonal precipitation. Overall water quality in Canada is high but it is difficult to provide a national overview as there is no comprehensive water quality monitoring system [3, 21, 27]. About 10% of the total population draws water from private rural household wells, which routinely do not meet drinking water quality standards for bacteria and nitrates. In some Provinces environmental water standards are exceeded for pesticides and phosphorus [3, 21, 28, 29, 30] which also impacts livestock water supplies [28]. About 15% of rural wells exceed guidelines for nitrates in drinking water (45 mg/litre) [3]. Depending on the region, 20-40% of surveyed rural wells have occurrences of coliform bacteria in excess of drinking water guidelines [3, 28, 30].

The Great Lakes ecosystem is stressed by farm nutrients, pathogens, pesticides and soil sediments, both from Canadian and US sources. These pollutants threaten recreational opportunities and raises costs of treating drinking water and dredging harbours [31, 32]. There has been some improvement in certain areas of the Great Lakes, such as the attainment of guideline levels of phosphorus for all lakes (except Lake Erie), due to a reduction of P inputs from agricultural, municipal and industrial sources. There is evidence that Canadian agricultural nutrient inputs (especially phosphate) to the Great Lakes could be declining as a result of improved farm management practices [33, 34]. Nutrient surpluses are an issue in some key watersheds, such as Lake Winnipeg which is showing signs of eutrophication [41], although farming is not the only source of nutrient pollution [35].

Agricultural nutrient surpluses per hectare are among the lowest in the OECD, however, they show the highest per cent increase across the OECD (Figure 3.4.2). In absolute values, the N surplus was 35 kg N/ha, about half the OECD average of 74 kg N/ha (2002-04). Both nitrogen (N) and phosphorus (P) surpluses grew respectively by 85% and 123% between 1990-92 and 2002-04. Nutrient surpluses (in tonnes) have grown in response to: greater inorganic fertiliser use – N fertiliser use rose by 35% between 1990-92 and 2002-04 and P use rose by 11% over the same period; the rise in pulse crop area (i.e. greater biological nitrogen fixation) without a concurrent reduction in fertiliser use; and higher livestock numbers generating growing quantities of manure [2]. In 1990-92, an estimated 40% of farmland suffered from a nitrogen deficit, however, this problem was addressed and by 2001 no land showed a nitrogen deficit. There are large regional variations in nutrient balances, owing to differing climates and types of soil, farming types and crops types, and also varying topography across the agricultural regions of Canada [2, 37].

Nutrient efficiency has declined, but the ratio is close to the OECD average for nitrogen and above it for phosphorus (nutrient efficiency is defined as the ratio of nutrient inputs and outputs). While the share of farms with formal nutrient management plans is low at 15% in 2001, several management practices are being adopted to protect water quality such as: establishing riparian areas adjacent to surface water on 75% of farms; conducting regular (1-5 years) soil nutrient tests on approximately two-thirds of farms; avoiding livestock feeding less than 100m from surface water during winter (on over 90% of farms); and preventing direct access of grazing livestock to surface water (nearly 60% of farms). Manure storage and application are key elements of most nutrient management plans, but between 1995 and 2001 manure application methods changed little, manure storage capacity was relatively low compared to manure production and timing of applications was not always optimal [2]. Between 1995 and 2001, 15% of producers adopted the optimal beneficial management practices for application of manure, representing 18% of total manure produced [2]. In 2001 10-11% of pig, poultry and dairy farms and 6% of beef farms, reported making environmental investments to reduce the risk of contamination to the environment from their operations [37].

Pesticide sales in Canada doubled between 1990 and 2003 [1, 2]. The risks associated with higher pesticide use, however, may to some extent be offset by: the use of new lower dose products that allow for targeted application; the expansion of genetically modified crops that are more pest-resistant; and the growth in organic farming, which accounted for under 1% of farmland and farms by 2003 and 1-2% of food sales despite its rapid growth in the past decade [1, 38, 39]. The growth in pesticide use is linked to the expansion in crop production, reduction in the use of summerfallow and greater intensity of farming [1]. Pesticides are used on over 80% of cropland [2, 40]. Over 60% of farmers are certified as

pesticide applicators, however, more efforts are required to encourage the uptake of beneficial management practices, such as recalibrating the sprayer before changing products, and spraying products at optimal times [1].

Under 10% of arable and horticultural farms in 2001 reported making environmental investments for pesticide storage and to combat water pollution from pesticides [37]. Pesticide residues have been detected in water bodies, but there is no systematic monitoring of pesticides in the environment [2, 19, 30]. Only 0.1% of rural wells were found to exceed drinking water standards for pesticides, which suggests management practices are helping to reduce risks [3]. The share of fresh fruit and vegetables with detectable pesticide residues decreased over the period 1995 to 2002 [30]. Since 1994 more than 20 instances of fish kills (with up to 35 000 dead fish collected in each incident) were attributed to pesticides in Prince Edward Island, and in British Columbia birds of prey were lost following the use of granular pesticides [40].

Agricultural water use is increasing. Water resources are abundant nationally; however, water availability varies across different regions of the country [2, 3]. In 1996 agriculture's share of total water use was over 10%, having increased by 3% from 1991 (Figure 3.4.2). Most of the growth in water use is being driven by the expansion in the area irrigated, which rose by 20% from 1990-92 to 2001-03, with most irrigation occurring in Alberta (55%) and British Columbia (21%) [19]. About 30% of irrigators in 2001 were fully or partially using best management practices. Water for irrigation is largely drawn from surface water [41, 42]. A study of Alberta shows improvements in irrigation efficiency over the past 30 years, but there is room for further progress with over 20% of the irrigated area using the less efficient gravity irrigation practices [43]. Increased risk of drought is a growing problem for farming in some regions, and one of Canada's most costly types of natural disaster [10], even in some of the usually more humid areas, such as the Atlantic Provinces [3].

Trends in harmful air emissions from agriculture have shown mixed results. The 3% growth in **ammonia emissions** between 1990 and 1995 was largely due to an intensification of livestock operations (Figure 3.4.2). Farming accounted for 80% of anthropogenic ammonia emissions, of which over 80% were from livestock. As industrial sources of acidifying substances (e.g. sulphur dioxide) have declined, the rise in agricultural ammonia emissions has eroded the benefits from this reduction [26]. In 2003 gaseous ammonia was listed on Schedule 1 of the *Canadian Environmental Protection Act* for its potential risk to human health as a precursor to fine particulate matter. Research is ongoing to learn more about ammonia emissions levels, transport, deposition and interaction with other substances in the air, and the contribution of the agriculture sector to the emissions. Over 45% of the total land area is highly sensitive to acid rain, with ammonia emissions contributing to the acidification of terrestrial and aquatic ecosystems [27, 44].

Canada has agreed to phase out its use of methyl bromide by 2005 under the Montreal Protocol. By 2004 use was reduced by over 70% from 1991 levels. In 2005 a *Critical Use Exemption* (CUE) was agreed, that allows methyl bromide use of up to 37 tonnes ozone depleting potential, which under the terms of the Protocol allows farmers more time to find substitutes for this pesticide.

Net greenhouse gas (GHGs) emissions from agriculture increased by around 1% between 1991 and 2001. This reflects an increase in both nitrous oxide, due to increased crop production and fertiliser use, and methane emissions, from the higher intensity in livestock operations, offset by a large net increase in carbon sequestration by soils as a

result of land use changes and improved management practices (Figure 3.4.3) [2, 45, 46]. Changes in agricultural management practices which are being implemented across Canada to reduce emissions, are largely market driven through innovations in equipment, as well as changes in relative prices of crops and inputs [47]. The increase of gross agricultural GHG emissions over the period 1990-92 to 2002-04 (18%) was substantially above the OECD average (-3%) but lower than the rise of 23% for total Canadian GHG emissions (Figure 3.4.2). Agriculture's share in total GHGs was 7% in 2002-04. Canada's commitment under the Kyoto Protocol is to reduce total GHG emissions by 6% by 2008-12, but recent announcements by the Government of Canada indicate that it may not be possible to meet this target.

Direct on-farm energy consumption rose by 5% between 1990-92 and 2002-04, which contributed to GHG emissions (Figure 3.4.2). Farm energy efficiency (the ratio of energy inputs to outputs) declined by 3% over the period 1989-93 to 1997-01, mainly due to the rise in diesel fuel and fertiliser use, the largest input components [2]. The production and consumption of **renewable energy** from agricultural biomass is minor compared to national total energy consumption, although under the new federal policy on biofuels the target is to achieve a 5% average renewable fuel content in transport fuel by 2010. This should create opportunities for biofuel producers to increase their renewable energy capacity [48, 49, 50].

Overall pressure on agricultural biodiversity continues. For agricultural genetic resources, Canada has in situ programmes and extensive *ex situ* collections of plant and animal genetic material, and efforts are underway to further expand this capacity [41, 51]. The number of major crop varieties and livestock breeds used in production has increased in diversity over the period 1990 to 2002. During this period the number of endangered livestock breeds rose from 47 to 51 (mainly cattle and sheep breeds), with only one breed under a conservation programme. This is in contrast to most other OECD countries where numbers of endangered breeds have declined as more livestock have come under conservation programmes, although two Canadian non-governmental organisations are involved in conserving rare livestock breeds [41].

There has been a substantial increase in the area under transgenic crops since the mid-1990s, accounting for 9% of the total agricultural land area in 2005, mainly canola with 70% of the sown crop genetically modified (GM) [35]. Canada is now the second major OECD producer, in terms of area, of transgenic crops after the United States.

The capacity of farmland to support wildlife showed a decline over the period 1991 to 2001. Over this 10 year period, 87% of Canada's farmland showed moderate to large decreases in habitat capacity compared to the 1981-2001, period when 30% of Canada's farmland showed a moderate to large decrease in habitat capacity (Figure 3.4.4). The agricultural intensification that has occurred in some areas of the country since 1981 is considered one of the drivers of the decrease in habitat capacity, such as the increase in cropland that occurred at the expense of more valuable habitats, for example wetlands, woodlots and natural pasture in Eastern Canada. Agricultural habitats, however, make a significant contribution to supporting many wild species by providing the necessary resources for breeding, feeding and cover [2].

Overall 24% of farms in 2001 were fully or partially using best management practices for wildlife conservation [41]. A number of regional studies suggest that the changing structure and fragmentation of agricultural habitats, and some farming practices, have raised concerns for the conservation of terrestrial and aquatic ecosystems, for example: the reduction in size and

loss of forest patches on farmland [52]; the fragmentation of native ecosystems [53, 54, 55]; the drainage of agricultural land and straightening of watercourses [55, 56, 57]; and run-off of excess nutrients and pesticides into surface water bodies.

The conversion of native ecosystems to farmland is considered to have been the main cause for the decline of most wild species, including threatened species [58]. The Canadian Wildlife Service grassland species breeding bird population index, decreased by almost 30% between 1990-92 to 2002-04, part of a longer term downward trend since the late 1960s, although from 2001 to 2004 there has been a small upward trend in the index of almost 10% [59]. Possible causes of the decline in grassland bird species include agricultural activities, urban growth into rural areas, and a decline in quality of wintering sites, among others. There is also evidence of recent increases in the Prairies breeding duck populations, although the longer term trend has been variable, for example declining in Southern Alberta, but expanding in Southern Saskatchewan [60].

3.4.3. Overall agri-environmental performance

Changes in farming practices and land use over the past decade have been successful in addressing environmental issues in some areas, but still need improvement in others. The adoption of soil management practices have resulted in improved soil quality, however the expansion and intensification of production over the past decade has increased environmental pressures in other areas [2, 61]. These include mainly water quality, especially in relation to manure management; growing competition for water resources; increase in ammonia and greenhouse gas emissions; and pressure on biodiversity. Given the size of Canada and its diversity of climate and soil types, there are wide regional differences in the environmental impacts of agriculture.

A comprehensive set of indicators to monitor the environmental performance of agriculture has been developed, within the context of *Canada's Agricultural Policy Framework (APF)* [2, 61]. Two agri-environmental indicator reports have been published to date (2000 and 2005), and a third is planned for 2008/09. Further development work is underway to strengthen the agri-environmental indicators in a number of areas, for example, soil biodiversity, particulate matter, and integrated pest management [2, 5]. A crucial challenge for indicator development and policy integration capacity are data limitations in key areas, such as pesticide use, agricultural water use, and a national monitoring network on water quality.

Canada is one of only a few OECD countries that does not regularly report the annual volume of pesticide use, although the Federal government stated in 1994 that it would establish a pesticide use database [40]. The lack of a national monitoring network on the quality of water (surface and groundwater) in rural areas has also been recognised as an impediment to effective policy analysis [33], while data related to agricultural water use are poor [42]. Efforts are being made, however, by the Federal government to collaborate with Provincial governments to fill these gaps, by conducting national surveys and establishing collaborative relationships with industry and academia. Agriculture and Agri-Food Canada is investigating the relationship between trends in critical habitat for wild species at risk and trends in agricultural land use.

Growing efforts by Federal and Provincial governments are tackling agri-environmental concerns. Under the environment pillar of the APF several programmes have been launched with the goal of reducing the sector's risk to the environment while remaining economically competitive. Programmes such as the *National Farm Stewardship Program*

provide technical support for producers to conduct environmental scans of their operations and develop *Environmental Farm Plans*. The Plans that identify actions to improve on-farm environmental performance, as well as providing cost-share support to implement these actions (i.e. fencing livestock out of water). There is still room for improvement to limit the impact of pesticides in the environment, however, work is ongoing to encourage producers to develop and adopt integrated pest management (IPM) systems which allow for continuous monitoring, adoption of alternative strategies for controlling pests, and targeted and efficient use of pesticides when required. The uptake of IPM practices is beginning to increase.

Within the APF the four-year CAD 60 (USD 45) million National Water Supply Expansion Program (2005) will address the growing risks of water shortages. The Program is making support available for on-farm water infrastructure, among other measures, and by providing a third of project costs [4]. The *Environmental Technology Assistance for Agriculture* programme evaluates innovative new technologies and production systems that are expected to contribute to improved on-farm economics and environmental performance, through nutrient management and the production of biofuels and renewable energy. Some of the key Provincial Government agri-environmental initiatives include: the implementation of a tax of CAD 1.2 (USD 0.8) per litre of pesticides in British Columbia; and Quebec's CAD 28 (USD 20) million *Prime-Vert Program* to control manure related pollution including a subsidy of 70-90% for the construction of manure storage facilities and restraints on manure spreading over winter [3, 5, 62].

The greenhouse Gas (GHG) Mitigation Program is an information and awareness programme, that encourages voluntary adoption of farm practices to reduce GHG emissions and increase carbon sinks. A comprehensive strategy to implement a 5% renewable fuels mandate for transport by 2010 is being established. The strategy plans to provide significant government incentives to support the expansion of the ethanol and biodiesel industry, and investment in research and development to encourage the growth of second generation biofuels, such as cellulosic ethanol.

A number of Provincial governments have in recent years introduced a range of measures to control water pollution from intensive livestock operations. These include, for example, the *Nutrient Management Act* in Ontario and the *Water Protection Plan* in Manitoba, which set targets for N and P levels in water bodies, and regulate some activities such as the timing of manure spreading to reduce risk of water contamination by agricultural sources [3, 19, 63]. Continued promotion of management practices that help reduce run-off of fertilisers and pesticides into the Great Lakes are planned as there are still improvements to be made [11]. Canada and the United States have also been working closely to develop an action plan to mitigate agricultural and industrial risks to the Great Lakes Basin under the *Great Lakes Regional Collaboration*, which aims to set goals to 2010 and 2015 to reduce agricultural pollutants into the Great Lakes, such as reducing livestock non-point source loading [31].


The agriculture sector is continuing efforts to reduce emissions of ammonia through the development and implementation of beneficial management practices that address manure management, storage and spreading and fertiliser application and storage. Research is ongoing to learn more about ammonia emission levels, transport, deposition and interaction with other substances in the air, as well as develop new beneficial management practices to reduce risk.

The projected expansion of agriculture to 2015 presents a considerable challenge to avoid an increase in environmental pressure [2, 64]. Changes in farming practices, especially the shift to reduced or no tillage, and land use changes, notably the reduction in summer fallow have yielded considerable environmental benefits, including: improved soil and water quality; lower energy use; reduced greenhouse gas emissions; and improvements for biodiversity. But these gains have partly been offset by the decreasing efficiency of nutrient and energy use. Rapidly growing nutrient surpluses could be offset with improvements to increase the uptake of best managements practices (BMPs), as only 15% of farms use BMPs to apply manure. Raising the efficiency of nutrient use would bring economic and environmental benefits. Subsidising on-farm fuel costs is a disincentive to improving energy use efficiency, reducing GHGs, and adopting conservation tillage (which requires less energy than conventional tillage) [65]. Only 6% of farms reported investment in environmental protection (i.e. manure storage, pesticide and fuel storage and waterway protection), averaging over CAD 19 200 (USD 12 400) or almost 4% of total farm investment in 2001 [37, 66].

A further challenge will be meeting Canada's international environmental commitments related to agriculture. The *International Joint Commission* has been requested to examine water diversions and removals from the Great Lakes, including for irrigation purposes, especially as water use conflicts and litigation have increased rapidly over the past decade [3]. Subsidised irrigation water and infrastructure do not facilitate the conservation of water resources and promotion of the efficient allocation of water between farming and other uses [3, 19]. While there has been success in lowering the use of *methyl bromide* since 1990, a further reduction will be required if Canada is to phase out its use as agreed under the *Montreal Protocol*. Given the increase in agricultural *ammonia* and gross GHG emissions it will also be a major challenge for Canada to meet its commitments to reduce emissions under the respective *Gothenburg* and *Kyoto Protocols*, although success has been achieved in increasing carbon sequestration in agricultural soils, helping to reduce net GHG emissions.

Figure 3.4.2. **National agri-environmental performance compared to the OECD average**

Percentage change 1990-92 to 2002-04¹



Absolute and economy-wide change/level


| Variable | Unit | | Canada | OECD |
|---------------------------------------|---------------------------------------|--------------------|--------|---------|
| Agricultural production volume | Index (1999-01 = 100) | 1990-92 to 2002-04 | 115 | 105 |
| Agricultural land area | 000 hectares | 1990-92 to 2002-04 | -1 521 | -48 901 |
| Agricultural nitrogen (N) balance | Kg N/hectare | 2002-04 | 35 | 74 |
| Agricultural phosphorus (P) balance | Kg P/hectare | 2002-04 | 1 | 10 |
| Agricultural pesticide use | Tonnes | 1990-92 to 2001-03 | n.a. | -46 762 |
| Direct on-farm energy consumption | 000 tonnes of oil equivalent | 1990-92 to 2002-04 | +184 | +1 997 |
| Agricultural water use | Million m ³ | 1990-92 to 2001-03 | +113 | +8 102 |
| Irrigation water application rates | Megalitres/ha of irrigated land | 2001-03 | 3.6 | 8.4 |
| Agricultural ammonia emissions | 000 tonnes | 1990-92 to 2001-03 | +14 | +115 |
| Agricultural greenhouse gas emissions | 000 tonnes CO ₂ equivalent | 1990-92 to 2002-04 | +8 043 | -30 462 |

n.a.: Data not available. Zero equals value between -0.5% to < +0.5%.

1. For agricultural water use, pesticide use, irrigation water application rates, and agricultural ammonia emissions the % change is over the period 1990-92 to 2001-03.
2. Percentage change in nitrogen and phosphorus balances in tonnes.


Source: OECD Secretariat. For full details of these indicators, see Chapter 1 of the Main Report.

Figure 3.4.3. **Share of cropland in different soil organic carbon change classes**



Source: Lefebvre, A., W. Eilers and B. Chunn (eds.) (2005), *Environmental Sustainability of Canadian Agriculture*, AEI. Report Series, Report 2, Agriculture and Agri-Food Canada, Ottawa.

Figure 3.4.4. **Share of farmland in different wildlife habitat capacity¹ change classes**



1. "Habitat capacity" is the capacity of agricultural land to sustain populations of wild terrestrial vertebrates, i.e. birds, mammals, reptiles and amphibians.

Source: Lefebvre, A., W. Eilers and B. Chunn (eds.) (2005), *Environmental Sustainability of Canadian Agriculture*, AEI Report Series, Report 2, Agriculture and Agri-Food Canada, Ottawa.

StatLink <http://dx.doi.org/10.1787/288868232073>

Bibliography

- [1] Agriculture and Agri-Food Canada (2005), *An Overview of the Canadian Agriculture and Agri-Food System*, Ottawa, Canada.
- [2] Lefebvre, A., W. Eliers and B. Chunn (eds.) (2005), *Environmental Sustainability of Canadian Agriculture: Agri-Environmental Indicator Report Series, Report 2*, Agriculture and Agri-Food Canada Ottawa, Canada, www.agr.gc.ca/env/naharp-pnarsa/index_e.php.
- [3] OECD (2004), *Environmental Performance Reviews: Canada*, Paris, France, www.oecd.org/env.
- [4] OECD (2005), *Agricultural Policies in OECD Countries: Monitoring and Evaluation 2005*, Paris, France, www.oecd.org/agr/policy.
- [5] McRae, T., C.A.S. Smith and L.J. Gregorich (eds.) (2000), *Environmental Sustainability of Canadian Agriculture: Report of the Agri-Environmental Indicator Project*, Agriculture and Agri-Food Canada, Ottawa, Canada, www.agr.gc.ca/policy/environment/pdfs/aei/fullreport.pdf.
- [6] Agriculture and Agri-Food Canada (2005), *Agricultural Policy Framework: Federal-Provincial-Territorial Programs*, Ottawa, Canada, www.agr.gc.ca/progser/pdf/APFProg_e.pdf.
- [7] Robinson, G.M. (2006), "Ontario's Environmental Farm Plan: Evaluation and research agenda", *Geoforum*, Vol. 37, Issue 5, September, pp. 859-873.
- [8] Smithers, J. and M. Furman (2003), "Environmental farm planning in Ontario: exploring participation and the endurance of change", *Land Use Policy*, Vol. 20, pp. 343-356.
- [9] OECD (2005), *Taxation and Social Security in Agriculture*, Paris, France, www.oecd.org/tad.
- [10] Kassem, A., T. McRae and M. Sydor (2006), "Integrated Water Resources Management", in OECD, *Water and Agriculture: Sustainability, Markets and Policies*, Paris, France, www.oecd.org/tad/env.
- [11] International Joint Commission (2004), *Twelfth Biennial Report on Great Lakes Water Quality*, September, Secretariat to the Great Lakes Water Quality Agreement, Ottawa, Canada, www.ijc.org/php/publications/html/12br/english/report/index.html.
- [12] Jayasinghe-Mudalige, J., A. Weersink, B. Deaton, M. Beaulieu and M. Trant (2005), "The urban-rural clash: Environmental management systems on Canadian farms", *Vista on the agri-food industry and the farm community*, May, Statistics Canada, Ottawa, Canada, www.statcan.ca.
- [13] Jayasinghe-Mudalige, J., A. Weersink, B. Deaton, M. Beaulieu and M. Trant (2005), "Effect of urbanisation on the adoption of environmental management systems in Canadian agriculture", Research Paper, Agriculture Division, Statistics Canada, Ottawa, Canada, www.statcan.ca.
- [14] Caldwell, W. and S. Hilts (2005), "Farmland preservation: Innovative approaches in Ontario", *Journal of Soil and Water Conservation*, Vol. 60, No. 3, pp. 66A-69A.
- [15] Agriculture and Agri-Food Canada (2001), *Agriculture in Harmony with Nature – Agriculture and Agri-Food Canada's Sustainable Development Strategy 2001-2004*, Ottawa, Canada, www.agr.ca/policy/environment/.
- [16] Lobb, D.A., E. Huffman and D.C. Reicosky (2005), "Importance of information on tillage practices in the modelling of environmental processes and in the use of environmental indicators", in OECD, *Farm Management and the Environment: Developing Indicators for Policy Analysis*, Paris, France, www.oecd.org/tad/env/indicators.
- [17] Singh, B. and S.S. Malhi (2006), "Response of soil physical properties to tillage and residue management on two soils in a cool temperate environment", *Soil and Tillage Research*, Vol. 85, pp. 143-153.
- [18] See various papers on soil organic carbon in Canadian agricultural soils in the Canadian-hosted OECD Workshop, see OECD (2003), *OECD, Soil Organic Carbon and Agriculture: Developing Indicators for Policy Analysis*, Paris, France, www.oecd.org/tad/env/indicators; and also see Smith, C.A.S., D.A. Lobb and C.M. Monreal (eds.), "Estimating Regional Soil Organic Carbon Stocks", *Canadian Journal of Soil Science*, Vol. 89, No. 4.
- [19] Renzetti, D.S. (2005), *Canadian agricultural water use and management*, Working Paper Series, Department of Economics, Brock University, St. Catharines, Ontario, Canada, http://139.57.161.145/papers/Ag_Water_Chapter.pdf.
- [20] Cessna, A.J., E. van Bochove, J. Yang, R. de Jong, A. Farenhorst and E. Topp (2004), "Agri-environmental water quality indicators: The Canadian experience", in OECD, *Agricultural Impacts on Water Use and Water Quality: Developing Indicators for Policy Analysis*, Paris, France, www.oecd.org/tad/env/indicators.
- [21] Statistics Canada (2005), *Canadian Environmental Sustainability Indicators*, Ottawa, Canada.

- [22] Environment Canada (2001), *Threats to sources of drinking water and aquatic ecosystem health in Canada*, Ottawa, Canada, www.nwri.ca/threats/intro-e.html.
- [23] Office of the Auditor General of Canada (2005), *Report of the Commissioner of the Environment and Sustainable Development to the House of Commons, The Commissioner's Perspective*, Ottawa, Canada, www.oag-bvg.gc.ca.
- [24] Gannon, V.P.J., G.D. Duke, J.E. Thomas, J. Van Leeuwen, J. Byrne, D. Johnson, S.W. Kienzle, J. Little, T. Graham and B. Selinger, "Use of in-stream reservoirs to reduce bacterial contamination of rural watersheds", *Science of the Total Environment*, Vol. 348, pp. 19-31.
- [25] Loë, R.C. de and R.D. Kreutzwiser (2005), "Closing the groundwater protection implementation gap", *Geoforum*, Vol. 36, pp. 241-256.
- [26] Environment Canada (2002), *Nutrients in the Canadian Environment, Reporting on the State of Canada's Environment*, Ottawa, Canada.
- [27] Environment Canada (2001), *Tracking Key Environmental Issues*, Ottawa, Canada.
- [28] Corkal, D., W.C. Schutzman and C.R. Hilliard (2004), "Rural water safety from the source to the on-farm tap", *Journal of Toxicology and Environmental Health, Part A*, Vol. 67, pp. 1619-1642.
- [29] Coote, D.R. and L.J. Gregorich (eds.) (2000), *The health of our waters – towards sustainable agriculture in Canada*, Research Branch, Agriculture and Agri-Food Canada, Ottawa, Canada.
- [30] Government of Canada (2005), *Children's Health and the Environment in North America – A First Report on Available Indicators and Measures*, report provided to the Commission on Environmental Co-operation, Montreal, Canada, www.cec.org/pubs_docs/documents/index.cfm?varlan=english&iD=1917.
- [31] Environment Canada (2005), *A strategy to restore and protect the Great Lakes*, Great Lakes Regional Collaboration, Draft Action Plan in collaboration with the United States Environmental Protection Agency, Ottawa, Canada, www.glrc.us/.
- [32] Environment Canada (2003), *The State of the Great Lakes*, Ottawa, Canada, <http://binational.net/sogl2003/sogl03eng.pdf>.
- [33] OECD (2003), "Some Environmental Aspects of Sustainable Development – Water Quality", pp. 88-91, *OECD Economic Surveys – Canada*, Vol. 2003/14, September, Paris, France.
- [34] Shear, H. (2006), "The Great Lakes, an ecosystem rehabilitated, but still under threat", *Environmental Monitoring and Assessment*, Vol. 113, pp. 199-225.
- [35] Venema, H.D. (2006), "From Cumulative Threats to Integrated Responses: A Review of Ag-Water Policy Issues in Prairie Canada", in *OECD, Water and Agriculture: Sustainability, Markets and Policies*, Paris, France, www.oecd.org/tad/env.
- [36] Janzen, H.H., K.A. Beauchemin, Y. Bruinsma, C.A. Campbell, R.L. Desjardins, B.H. Ellert and E.G. Smith (2003), "The fate of nitrogen in agro-ecosystems: An illustration using Canadian estimates", *Nutrient Cycling in Agroecosystems*, Vol. 67, pp. 85-102.
- [37] Grimard, J. (2004), "Partial portrait of farm investments in environmental protection", *Vista on the agri-food industry and the farm community*, January, Statistics Canada, Ottawa, Canada, www.statcan.ca.
- [38] Natural Resources Canada (2005), *Energy Efficiency Trends in Canada 1990 to 2003*, Office of Energy Efficiency, Ottawa, Canada.
- [39] Magnusson, E. and J.A.L. Cranfield (2005), "Consumer demand for pesticide free food products in Canada: A probit analysis", *Canadian Journal of Agricultural Economics*, Vol. 53, pp. 67-81.
- [40] Office of the Auditor General of Canada (2003), *Report of the Commissioner of the Environment and Sustainable Development to the House of Commons: Chapter 1 Managing the Safety and Accessibility of Pesticides*, Ottawa, Canada, www.oag-bvg.gc.ca.
- [41] Canadian response to the OECD Agri-environmental Indicator Questionnaire, unpublished.
- [42] Environment Canada (2004), *Threats to water availability in Canada*, Ottawa, Canada, www.nwri.ca/threats2full/intro-e.html.
- [43] Bjornlund, H., L. Nicol and K.K. Klein (2006), *Alberta's Water for Life Strategy: Some early indications of its acceptance by the irrigation industry in Southern Alberta*, paper presented to the Annual Meeting of the Canadian Agricultural Economics Society, Montreal, Quebec.
- [44] Kurvits, T. and T. Marta (1998), "Agricultural NH₃ and NO_x emissions in Canada", *Environmental Pollution*, Vol. 102, S1, pp. 187-194.

- [45] Boame, A.K. (2005), "Zero tillage: a greener way for Canadian farms", *Vista on the agri-food industry and the farm community*, November, Statistics Canada, Ottawa, Canada, www.statcan.ca.
- [46] Desjardins, R.L., W. Smith, B. Grant, C. Campbell and R. Riznek (2005), "Management strategies to sequester carbon in agricultural soils and to mitigate greenhouse gas emissions", *Climate Change*, Vol. 70, pp. 283-297.
- [47] Smith, E.G. and B.M. Upadhyay (2005), *Greenhouse gas mitigation on diversified farms*, paper presented to the joint Annual Meeting of the Canadian Agricultural Economics Society and Western Agricultural Economics Association, San Francisco, July.
- [48] IEA (2004), *Energy Policies of IEA Countries – Canada 2004 Review*, Paris, France, www.iea.org.
- [49] Tupper, D. (2005), "The Canadian Situation – Biomass and Agriculture", in OECD, *Biomass and Agriculture: Sustainability, Markets and Policies*, Paris, France, www.oecd.org/tad/env.
- [50] OECD (2006), *Agricultural Market Impacts of Future Growth in the Production of Biofuels*, Paris, France, www.oecd.org/dataoecd/58/62/36074135.pdf.
- [51] Environment Canada (2005), *Canada – Third National Report to the Convention on Biological Diversity*, Secretariat to the Convention on Biological Diversity, Montreal, Canada, www.biodiv.org/reports/list.aspx?type=all.
- [52] Silva, M., L. Hartling and S.B. Opps (2005), "Small mammals in agricultural landscapes of Prince Edward Island (Canada): Effects of habitat characteristics at three different spatial scales", *Biological Conservation*, Vol. 126, pp. 556-568.
- [53] McLachlan, S.M. and A.L. Knispel (2005), "Assessment of long-term tallgrass prairie restoration in Manitoba, Canada", *Biological Conservation*, Vol. 124, pp. 75-88.
- [54] Dunford, W. and K. Freemark (2004), "Matrix matters: effects of surrounding land uses on forest birds near Ottawa, Canada", *Landscape Ecology*, Vol. 20, pp. 497-511.
- [55] Kerr, J.T. and J. Cihlar (2003), "Land use and cover with intensity of agriculture for Canada from satellite and census data", *Global Ecology and Biogeography*, Vol. 12, pp. 161-172.
- [56] Walters, D. and D. Shrubsole (2003), "Agricultural drainage and wetland management in Ontario", *Journal of Environmental Management*, Vol. 69, pp. 369-379.
- [57] Boutin, C., B. Jobin and L. Bélanger (2003), "Importance of riparian habitats to flora conservation in farming landscapes of southern Québec, Canada", *Agriculture, Ecosystems and Environment*, Vol. 94, pp. 73-87.
- [58] Kerr, J.T. and I. Deguise (2004), "Habitat loss and the limits to endangered species recovery", *Ecology Letters*, Vol. 7, pp. 1163-1169.
- [59] Canadian Wildlife Service (2006), *Canadian Bird Trends Database*, website database, Environment Canada, Hull, Canada, www.cws-scf.ec.gc.ca.
- [60] Wilkins, K.A. and M.C. Otto (2003), *Trends in duck breeding populations, 1955-2003*, US Fish and Wildlife Service, Laurel, Maryland, United States.
- [61] McRae, T., L. Heigh, B. Junkins and A. Lefebvre (2005), "Using Environmental Indicators to Support the Development of Agricultural Policy: The Canadian Experience", in OECD, *Farm Management and the Environment: Developing Indicators for Policy Analysis*, Paris, France, www.oecd.org/tad/env/indicators.
- [62] Boutin, D. (2006), "The Challenge of Reconciling Water and Agricultural Policies – The Role of Public Hearings", in OECD, *Water and Agriculture: Sustainability, Markets and Policies*, Paris, France, www.oecd.org/tad/env.
- [63] Cantin, B., S. Kalff and I. Campbell (2006), "Assessing the feasibility of water quality trading to address agricultural sources of pollution in Canada", in OECD, *Water and Agriculture: Sustainability, Markets and Policies*, Paris, France, www.oecd.org/tad/env.
- [64] OECD (2006), *OECD-FAO Agricultural Outlook 2006-2015*, Paris, France, www.oecd.org/tad.
- [65] Weersink, A, D. Pannell, M. Fulton and A. Meyer-Aurich (2005), "Agriculture's likely role in meeting Canada's Kyoto commitments", *Canadian Journal of Agricultural Economics*, Vol. 53, pp. 425-441.
- [66] Agriculture and Agri-Food Canada (2004), *Quantitative analysis of the Impact of Agricultural Management Strategies on Environmental Indicators*, Ottawa, Canada, www.agr.gc.ca/index_e.php?s1=info&s2=pub.

Table of Contents

| | |
|---|----|
| I. Highlights | 15 |
| Overall agri-environmental performance. | 15 |
| Agri-environmental performance in specific areas | 16 |
| Caveats and limitations | 19 |
| Matching indicator criteria. | 20 |
| II. Background and Scope of the Report. | 23 |
| 1. Objectives and scope. | 23 |
| 2. Data and information sources. | 24 |
| 3. Progress made since the OECD 2001 Agri-environmental Indicator Report | 25 |
| 4. Structure of the Report | 26 |
| Bibliography | 28 |
| Annex II.A1. List of indicators in Chapter 1 | 29 |
| Annex II.A2. Indicators in Chapter 1 assessed according to the OECD indicator criteria | 31 |
| Chapter 1. OECD Trends of Environmental Conditions related to Agriculture since 1990 | 37 |
| 1.1. Agricultural production and land | 38 |
| 1.1.1. Introduction | 39 |
| 1.1.2. Agricultural production | 39 |
| 1.1.3. Agricultural land use. | 40 |
| 1.1.4. Linkages between agricultural production and land use. | 46 |
| Bibliography | 47 |
| 1.2. Nutrients | 48 |
| 1.2.1. Nitrogen balance | 52 |
| 1.2.2. Phosphorus balance | 56 |
| 1.2.3. Regional (sub-national) nutrient balances. | 60 |
| Bibliography | 62 |
| 1.3. Pesticides | 63 |
| 1.3.1. Pesticide use | 63 |
| 1.3.2. Pesticide risk indicators | 67 |
| Bibliography | 74 |
| 1.4. Energy | 76 |
| Bibliography | 83 |
| 1.5. Soil | 84 |
| Bibliography | 90 |

| | |
|--|------------|
| 1.6. Water..... | 92 |
| 1.6.1. Water use | 93 |
| 1.6.2. Water quality | 100 |
| Bibliography | 108 |
| 1.7. Air | 109 |
| Background | 110 |
| 1.7.1. Ammonia emissions, acidification and eutrophication..... | 110 |
| 1.7.2. Methyl bromide use and ozone depletion | 117 |
| 1.7.3. Greenhouse gas emissions and climate change | 122 |
| Bibliography | 130 |
| 1.8. Biodiversity | 133 |
| Background | 134 |
| 1.8.1. Genetic diversity | 136 |
| 1.8.2. Wild species diversity | 146 |
| 1.8.3. Ecosystem diversity..... | 148 |
| Bibliography | 159 |
| 1.9. Farm management | 160 |
| 1.9.1. Overview of environmental farm management | 163 |
| 1.9.2. Nutrient management | 163 |
| 1.9.3. Pest management | 168 |
| 1.9.4. Soil management..... | 169 |
| 1.9.5. Water management..... | 172 |
| 1.9.6. Biodiversity management | 173 |
| 1.9.7. Organic management | 174 |
| Bibliography | 176 |
| Chapter 2. OECD Progress in Developing Agri-environmental Indicators | 179 |
| 2.1. Introduction..... | 180 |
| 2.2. Progress in developing OECD Agri-environmental Indicators | 180 |
| 2.2.1. Soil: Erosion, biodiversity and soil organic carbon | 180 |
| 2.2.2. Water: Use and water quality | 184 |
| 2.2.3. Biodiversity: Genetic, wild species and ecosystem diversity | 188 |
| 2.2.4. Land: Landscapes and ecosystem functions | 192 |
| 2.2.5. Farm management | 195 |
| 2.3. Overall assessment..... | 196 |
| Annex 2.A1. Agri-environmental Indicators of Regional Importance and/or under Development..... | 200 |
| Annex 2.A2. A Qualitative Assessment of the Agri-environmental Indicators included in Annex 2.A1 according to the OECD Indicator Criteria | 202 |
| Bibliography | 207 |
| Chapter 3. OECD Country Trends of Environmental Conditions related to Agriculture since 1990 | 209 |
| Background to the country sections | 210 |
| 3.1. Australia | 212 |
| 3.2. Austria | 224 |
| 3.3. Belgium..... | 234 |
| 3.4. Canada | 243 |

| | |
|--|------------|
| 3.5. Czech Republic | 256 |
| 3.6. Denmark..... | 269 |
| 3.7. Finland | 284 |
| 3.8. France | 296 |
| 3.9. Germany | 305 |
| 3.10. Greece..... | 313 |
| 3.11. Hungary | 324 |
| 3.12. Iceland | 336 |
| 3.13. Ireland..... | 344 |
| 3.14. Italy | 357 |
| 3.15. Japan..... | 366 |
| 3.16. Korea..... | 377 |
| 3.17. Luxembourg..... | 386 |
| 3.18. Mexico..... | 393 |
| 3.19. Netherlands | 402 |
| 3.20. New Zealand | 413 |
| 3.21. Norway | 423 |
| 3.22. Poland..... | 433 |
| 3.23. Portugal..... | 448 |
| 3.24. Slovak Republic | 459 |
| 3.25. Spain..... | 472 |
| 3.26. Sweden..... | 486 |
| 3.27. Switzerland | 498 |
| 3.28. Turkey..... | 507 |
| 3.29. United Kingdom | 522 |
| 3.30. United States | 532 |
| 3.31. European Union..... | 545 |
| Chapter 4. Using Agri-environmental Indicators for Policy Analysis | 551 |
| 4.1. Policy context to OECD agri-environmental performance | 552 |
| 4.2. Tracking agri-environmental performance..... | 554 |
| 4.2.1. Evolution of Agri-environmental Indicators to track sustainable development..... | 554 |
| 4.2.2. Tracking national agri-environmental performance | 556 |
| 4.2.3. International reporting on environmental conditions in agriculture | 559 |
| 4.2.4. Non-governmental organisations (NGOs) | 561 |
| 4.3. Using Agri-environmental Indicators for policy analysis | 562 |
| 4.3.1. OECD member countries | 563 |
| 4.3.2. International governmental organisations | 565 |
| 4.3.3. Research community | 567 |
| 4.4. Knowledge gaps in using Agri-environmental Indicators..... | 568 |
| Bibliography | 571 |
| | |
| List of boxes | |
| II.1. OECD Expert Meetings on Agri-environmental Indicators: 2001-04 | 25 |
| 1.7.1. Towards a net agricultural greenhouse gas balance indicator?..... | 123 |

| | |
|--|-----|
| 1.8.1. Defining agricultural biodiversity | 134 |
| 2.1. Soil biodiversity in agricultural land | 182 |
| 2.2. Agricultural livestock pathogens and water pollution | 187 |
| 2.3. The impact of agriculture on aquatic ecosystems | 188 |
| 4.1. Main agri-environmental measures in OECD countries | 553 |
| 4.2. Selected international and regional environmental agreements relevant to agriculture | 555 |

List of tables

| | |
|--|-----|
| 1.1.1. OECD and world agricultural production | 39 |
| 1.1.2. OECD and world agricultural exports | 40 |
| 1.3.1. Germany: Percentage risk indices | 70 |
| 1.7.1. Total OECD emissions of acidifying pollutants | 114 |
| 1.7.2. Ammonia emission targets to 2010 under the Convention on Long-range Transboundary Air Pollution | 116 |
| 1.7.3. Methyl bromide use and progress in meeting the phase-out schedule under the <i>Montreal Protocol</i> | 120 |
| 1.7.4. Critical Use Exemptions (CUEs) for methyl bromide agreed under the <i>Montreal Protocol</i> for 2005 | 121 |
| 1.7.5. Total OECD gross greenhouse gas emissions | 124 |
| 1.7.6. Main sources and types of gross greenhouse gas emissions | 127 |
| 1.8.1. Area of transgenic crops for major producing countries | 139 |
| 1.8.2. Plant genetic resource conservation activities for OECD countries | 139 |
| 1.8.3. Livestock genetic resource conservation activities for OECD countries | 144 |
| 1.8.4. Share of farm woodland in agricultural land area | 157 |
| 1.8.5. Share of farm fallow in agricultural land area | 157 |
| 1.9.1. Countries recording adoption of environmental farm management practices | 164 |
| 1.9.2. Overview of farmer incentives to adopt environmental farm management practices | 166 |
| 2.1. Net water balance in a Japanese rice field irrigation system: 2003 | 185 |

List of figures

| | |
|--|----|
| II.1. The Driving Force-State-Response framework: Coverage of indicators | 24 |
| 1.1.1. Production, yields and area harvested and future projections for selected commodities and OECD countries | 41 |
| 1.1.2. Volume of total agricultural production | 43 |
| 1.1.3. Share of agricultural land use in the national land area | 44 |
| 1.1.4. Agricultural land area | 45 |
| 1.1.5. Agricultural production volume index and agricultural land area | 46 |
| 1.2.1. Main elements in the OECD gross nutrient (nitrogen and phosphorus) balance calculation | 50 |
| 1.2.2. Gross nitrogen balance estimates | 51 |
| 1.2.3. Gross nitrogen balances for selected OECD countries | 53 |
| 1.2.4. Inorganic nitrogen fertilisers and livestock manure nitrogen input in nitrogen balances | 54 |

| | |
|---|-----|
| 1.2.5. Agricultural use of inorganic nitrogen and phosphate fertilisers | 54 |
| 1.2.6. Contribution of the main sources of nitrogen inputs and outputs in nitrogen balances | 56 |
| 1.2.7. Nitrogen efficiency based on gross nitrogen balances | 57 |
| 1.2.8. Gross phosphorus balance estimates | 58 |
| 1.2.9. Gross phosphorus balance for selected OECD countries | 59 |
| 1.2.10. Contribution of the main sources of phosphorus inputs and outputs in phosphorus balances | 60 |
| 1.2.11. Phosphorus efficiency based on phosphorus balances | 61 |
| 1.2.12. Spatial distribution of nitrogen balances in Canada and Poland | 62 |
| 1.3.1. Pesticide use in agriculture | 65 |
| 1.3.2. Pesticide use for selected OECD countries | 66 |
| 1.3.3. Belgium: Risk for aquatic species due to use of pesticides in arable land, horticulture and outside of agriculture | 69 |
| 1.3.4. Denmark: The annual trend in frequency of pesticide application | 70 |
| 1.3.5. The Netherlands: Potential chronic effects scores for aquatic and terrestrial organisms and leaching into groundwater | 71 |
| 1.3.6. Norway: Trends of health risk, environmental risk and sales of pesticides | 72 |
| 1.3.7. Sweden: National level pesticide risk indicators and the number of hectare doses | 73 |
| 1.3.8. United Kingdom (England and Wales): Total area of pesticide applications | 74 |
| 1.4.1. Simplified energy “model” of an agricultural system | 78 |
| 1.4.2. Direct on-farm energy consumption | 79 |
| 1.4.3. Direct on-farm energy consumption for selected OECD countries | 80 |
| 1.4.4. Agricultural employment and farm machinery use | 81 |
| 1.4.5. Composition of on-farm energy consumption in the EU15 and the United States | 82 |
| 1.5.1. Agricultural land area classified as having moderate to severe water erosion risk | 87 |
| 1.5.2. Trends in agricultural land area classified as having moderate to severe water erosion risk | 88 |
| 1.5.3. Agricultural land area classified as having moderate to severe wind erosion risk | 89 |
| 1.6.1. Agricultural water use | 95 |
| 1.6.2. Share of national water use in annual freshwater resources and share of agricultural water use in national use | 96 |
| 1.6.3. Irrigated area, irrigation water use and irrigation water application rates | 97 |
| 1.6.4. Share of agricultural groundwater use in total groundwater use, and total groundwater use in total water use | 99 |
| 1.6.5. Share of agriculture in total emissions of nitrates and phosphorus in surface water | 102 |
| 1.6.6. Share of agriculture in total emissions of nitrates and phosphorus in coastal water | 103 |
| 1.6.7. Share of monitoring sites in agricultural areas exceeding national drinking water limits for nitrates and phosphorus in surface water | 104 |
| 1.6.8. Share of monitoring sites in agricultural areas exceeding national drinking water limits for nitrates in groundwater | 105 |

| | |
|---|-----|
| 1.6.9. Share of monitoring sites in agricultural areas where one or more pesticides are present in surface and groundwater | 106 |
| 1.6.10. Share of monitoring sites in agricultural areas exceeding national drinking water limits for pesticides in surface water and groundwater | 107 |
| 1.7.1. Impacts of agriculture on air quality: Multi-pollutants, multi-effects | 110 |
| 1.7.2. Ammonia emissions from agriculture | 112 |
| 1.7.3. Emissions of acidifying airborne pollutants for the EU15, US and OECD. | 113 |
| 1.7.4. Agricultural ammonia emission trends for selected OECD countries | 114 |
| 1.7.5. Share of the main sources of agricultural ammonia emissions in OECD countries | 117 |
| 1.7.6. Methyl bromide use | 119 |
| 1.7.7. Global methyl bromide use by major sectors. | 121 |
| 1.7.8. Agricultural gross greenhouse gas emissions | 125 |
| 1.7.9. Gross agricultural greenhouse gas emissions in carbon dioxide equivalent for selected OECD countries | 126 |
| 1.7.10. Agricultural production and agricultural greenhouse gas emissions. | 128 |
| 1.7.11. Main sources of methane and nitrous oxide emissions in OECD agriculture | 129 |
| 1.7.12. Contribution of main sources in agricultural greenhouse gas emissions | 130 |
| 1.8.1. OECD agri-biodiversity indicators framework | 135 |
| 1.8.2. Change in the number of plant varieties registered and certified for marketing | 137 |
| 1.8.3. Change in the share of the one-to-five dominant crop varieties in total marketed crop production | 138 |
| 1.8.4. Change in the number of livestock breeds registered and certified for marketing | 141 |
| 1.8.5. Change in the share of the three major livestock breeds in total livestock numbers. | 142 |
| 1.8.6. Total number of cattle, pigs, poultry and sheep in endangered and critical risk status and under conservation programmes | 143 |
| 1.8.7. Share of selected wild species that use agricultural land as primary habitat. | 148 |
| 1.8.8. Population trends of farmland birds | 149 |
| 1.8.9. Change in agricultural land use and other uses of land. | 152 |
| 1.8.10. Permanent pasture and arable and permanent cropland | 155 |
| 1.8.11. Share of arable and permanent cropland, permanent pasture and other agricultural land in total agricultural land area. | 156 |
| 1.8.12. Share of national Important Bird Areas where intensive agricultural practices pose a serious threat or a high impact on the areas' ecological functions | 158 |
| 1.9.1. OECD farm management indicator framework | 162 |
| 1.9.2. Share of agricultural land area under nutrient management plans. | 168 |
| 1.9.3. Share of total number of farms under nutrient management plans | 169 |
| 1.9.4. Share of total number of farms using soil nutrient testing | 170 |
| 1.9.5. Share of total arable and permanent crop area under integrated pest management. | 171 |
| 1.9.6. Share of arable crop area under soil conservation practices | 172 |
| 1.9.7. Share of total arable and permanent crop area under all-year vegetative cover | 173 |
| 1.9.8. Share of irrigated land area using different irrigation technology systems | 174 |

| | |
|---|-----|
| 1.9.9. Share of agricultural land area under biodiversity management plans | 175 |
| 1.9.10. Share of agricultural land area under certified organic farm management | 176 |
| 2.1. Canadian soil organic carbon stocks in agricultural soils by different classes | 183 |
| 2.2. United States soil organic carbon stocks in agricultural soils by different classes | 184 |
| 2.3. Agricultural, industrial, and household water charges | 186 |
| 2.4. National crop varieties that are endangered | 189 |
| 2.5. National crop varieties that are not at risk. | 190 |
| 2.6. Edge density of agricultural fields in Finland. | 190 |
| 2.7. Share of Canadian farmland in various classes of the habitat capacity index. | 191 |
| 2.8. Cultural landscape features on agricultural land | 193 |
| 2.9. Water retaining capacity of agriculture | 194 |
| 2.10. Water retaining capacity for agricultural facilities | 195 |
| 2.11. Share of farmers participating in agri-environmental education programmes | 197 |
| 3.1.1. National agri-environmental and economic profile, 2002-04: Australia | 212 |
| 3.1.2. National agri-environmental performance compared to the OECD average. | 220 |
| 3.1.3. National Landcare membership. | 220 |
| 3.1.4. Annual quantities of insecticide and acaricide applied to the cotton crop | 220 |
| 3.2.1. National agri-environmental and economic profile, 2002-04: Austria | 224 |
| 3.2.2. National agri-environmental performance compared to the OECD average. | 231 |
| 3.2.3. Area under non-use of inputs, organic farming and erosion control measures of the ÖPUL agri-environmental programme. | 231 |
| 3.2.4. Greenhouse gas emissions from agriculture | 231 |
| 3.3.1. National agri-environmental and economic profile, 2002-04: Belgium | 234 |
| 3.3.2. National agri-environmental performance compared to the OECD average. | 240 |
| 3.3.3. Total pesticide use | 240 |
| 3.3.4. Greenhouse gas emissions and sinks | 240 |
| 3.4.1. National agri-environmental and economic profile, 2002-04: Canada | 243 |
| 3.4.2. National agri-environmental performance compared to the OECD average. | 252 |
| 3.4.3. Share of cropland in different soil organic carbon change classes. | 252 |
| 3.4.4. Share of farmland in different wildlife habitat capacity change classes. | 252 |
| 3.5.1. National agri-environmental and economic profile, 2002-04: Czech Republic | 256 |
| 3.5.2. National agri-environmental performance compared to the OECD average. | 265 |
| 3.5.3. Share of samples above Czech drinking water standards for nitrates in surface water | 265 |
| 3.5.4. Monitored numbers of partridge population | 265 |
| 3.6.1. National agri-environmental and economic profile, 2002-04: Denmark | 269 |
| 3.6.2. National agri-environmental performance compared to the OECD average. | 280 |
| 3.6.3. Share of monitoring sites with occurrences of pesticides in groundwater used for drinking | 280 |
| 3.6.4. Share of meadows and dry grasslands, heath, and bogs and marshes in the total land area | 280 |
| 3.7.1. National agri-environmental and economic profile, 2002-04: Finland | 284 |
| 3.7.2. National agri-environmental performance compared to the OECD average. | 292 |
| 3.7.3. Nitrogen fluxes in the Paimionjoki river and agricultural nitrogen balances | 292 |

| | |
|---|-----|
| 3.7.4. Population trends of Finnish farmland butterflies in three ecological species groups. | 292 |
| 3.8.1. National agri-environmental and economic profile, 2002-04: France. | 296 |
| 3.8.2. National agri-environmental performance compared to the OECD average. | 302 |
| 3.8.3. Trends in key agri-environmental indicators. | 302 |
| 3.8.4. Trends in key agri-environmental indicators. | 302 |
| 3.9.1. National agri-environmental and economic profile, 2002-04: Germany | 305 |
| 3.9.2. National agri-environmental performance compared to the OECD average. | 310 |
| 3.9.3. Share of the number of farms and Utilised Agricultural Area (UAA) under organic farming. | 310 |
| 3.9.4. Share of renewable biomass and energy crop area in the total agricultural land area | 310 |
| 3.10.1. National agri-environmental and economic profile, 2002-04: Greece | 313 |
| 3.10.2. National agri-environmental performance compared to the OECD average. | 321 |
| 3.10.3. Irrigated area and irrigation water application rates | 321 |
| 3.10.4. <i>Ex situ</i> accessions of plant landraces, wild and weedy relatives. | 321 |
| 3.11.1. National agri-environmental and economic profile, 2002-04: Hungary | 324 |
| 3.11.2. National agri-environmental performance compared to the OECD average. | 333 |
| 3.11.3. Agricultural land affected by various classes of water erosion | 333 |
| 3.11.4. Support payments for agri-environmental schemes and the number of paid applications. | 333 |
| 3.12.1. National agri-environmental and economic profile, 2002-04: Iceland | 336 |
| 3.12.2. National agri-environmental performance compared to the OECD average. | 342 |
| 3.12.3. Annual afforestation | 342 |
| 3.12.4. Annual area of wetland restoration. | 342 |
| 3.13.1. National agri-environmental and economic profile, 2002-04: Ireland | 344 |
| 3.13.2. National agri-environmental performance compared to the OECD average. | 353 |
| 3.13.3. River water quality | 353 |
| 3.13.4. Population changes for key farmland bird populations | 353 |
| 3.14.1. National agri-environmental and economic profile, 2002-04: Italy. | 357 |
| 3.14.2. National agri-environmental performance compared to the OECD average. | 363 |
| 3.14.3. Actual soil water erosion risk. | 363 |
| 3.14.4. Regional change in agricultural land area: 1990 to 2000. | 363 |
| 3.15.1. National agri-environmental and economic profile, 2002-04: Japan | 366 |
| 3.15.2. National agri-environmental performance compared to the OECD average. | 373 |
| 3.15.3. National water retaining capacity of agriculture. | 373 |
| 3.15.4. Share of eco-farmers in the total number of farmers. | 373 |
| 3.16.1. National agri-environmental and economic profile, 2002-04: Korea | 377 |
| 3.16.2. National agri-environmental performance compared to the OECD average. | 383 |
| 3.16.3. Composition of soils | 383 |
| 3.16.4. National water retaining capacity of agriculture. | 383 |
| 3.17.1. National agri-environmental and economic profile, 2002-04: Luxembourg | 386 |
| 3.17.2. National agri-environmental performance compared to the OECD average. | 391 |
| 3.17.3. Nitrate and phosphorus concentration in river sampling stations. | 391 |
| 3.17.4. Agricultural land under agri-environmental schemes | 391 |
| 3.18.1. National agri-environmental and economic profile, 2002-04: Mexico | 393 |
| 3.18.2. National agri-environmental performance compared to the OECD average. | 399 |

| | |
|--|-----|
| 3.18.3. Trends in key agri-environmental indicators | 399 |
| 3.18.4. Trends in key agri-environmental indicators | 399 |
| 3.19.1. National agri-environmental and economic profile, 2002-04: Netherlands | 402 |
| 3.19.2. National agri-environmental performance compared to the OECD average | 409 |
| 3.19.3. Annual mean concentrations of nitrogen and phosphorus in surface water of rural and agricultural water catchments | 409 |
| 3.19.4. Farmland bird populations | 409 |
| 3.20.1. National agri-environmental and economic profile, 2002-04: New Zealand | 413 |
| 3.20.2. National agri-environmental performance compared to the OECD average | 420 |
| 3.20.3. Sectoral use of pesticides: 2004 | 420 |
| 3.20.4. Dairy cattle enteric methane emissions per litre of milk | 420 |
| 3.21.1. National agri-environmental and economic profile, 2002-04: Norway | 423 |
| 3.21.2. National agri-environmental performance compared to the OECD average | 430 |
| 3.21.3. National sales of pesticides | 430 |
| 3.21.4. Net change in agricultural land for five counties | 430 |
| 3.22.1. National agri-environmental and economic profile, 2002-04: Poland | 433 |
| 3.22.2. National agri-environmental performance compared to the OECD average | 444 |
| 3.22.3. Agriculture and forest land at risk to erosion | 444 |
| 3.22.4. Index of population trends of farmland birds | 444 |
| 3.23.1. National agri-environmental and economic profile, 2002-04: Portugal | 448 |
| 3.23.2. National agri-environmental performance compared to the OECD average | 456 |
| 3.23.3. Numbers of local breeds under <i>in situ</i> conservation programmes: 2006 | 456 |
| 3.23.4. Relation between land use and Designated Nature Conservation Areas (DNCA): 2004 | 456 |
| 3.24.1. National agri-environmental and economic profile, 2002-04: Slovak Republic . . | 459 |
| 3.24.2. National agri-environmental performance compared to the OECD average | 468 |
| 3.24.3. Agricultural methane (CH ₄) and nitrous oxide (N ₂ O) emissions | 468 |
| 3.24.4. Share of agricultural land under different types of protected areas: 2003 | 468 |
| 3.25.1. National agri-environmental and economic profile, 2002-04: Spain | 472 |
| 3.25.2. National agri-environmental performance compared to the OECD average | 482 |
| 3.25.3. Area of organic farming | 482 |
| 3.25.4. Share of Dehesa area in total land area for five regions | 482 |
| 3.26.1. National agri-environmental and economic profile, 2002-04: Sweden | 486 |
| 3.26.2. National agri-environmental performance compared to the OECD average | 494 |
| 3.26.3. Losses of nutrients from arable areas and the root zone | 494 |
| 3.26.4. Cultural features on arable land | 494 |
| 3.27.1. National agri-environmental and economic profile, 2002-04: Switzerland | 498 |
| 3.27.2. National agri-environmental performance compared to the OECD average | 504 |
| 3.27.3. Support for agricultural semi-natural habitats | 504 |
| 3.27.4. Input/output efficiency of nitrogen, phosphorous and energy in agriculture . . . | 504 |
| 3.28.1. National agri-environmental and economic profile, 2002-04: Turkey | 507 |
| 3.28.2. National agri-environmental performance compared to the OECD average | 518 |
| 3.28.3. Trends in key agri-environmental indicators | 518 |
| 3.28.4. Trends in key agri-environmental indicators | 518 |
| 3.29.1. National agri-environmental and economic profile, 2002-04: United Kingdom | 522 |
| 3.29.2. National agri-environmental performance compared to the OECD average | 528 |

| | |
|---|-----|
| 3.29.3. Agri-environmental trends | 528 |
| 3.29.4. Greenhouse gas emission trends and projections. | 528 |
| 3.30.1. National agri-environmental and economic profile, 2002-04: United States. | 532 |
| 3.30.2. National agri-environmental performance compared to the OECD average. | 540 |
| 3.30.3. Soil erosion on cropland | 540 |
| 3.30.4. Change in palustrine and estuarine wetlands on non-federal land and water area | 540 |
| 3.31.1. National agri-environmental and economic profile, 2002-04: European Union (15) | 545 |
| 3.31.2. EU15 agri-environmental performance compared to the OECD average. | 548 |
| 3.31.3. Agri-environmental trends, EU15 | 548 |
| 3.31.4. Agri-environmental trends, EU15 | 548 |

This book has...



StatLinks 

**A service that delivers Excel® files
from the printed page!**

Look for the *StatLinks* at the bottom right-hand corner of the tables or graphs in this book. To download the matching Excel® spreadsheet, just type the link into your Internet browser, starting with the <http://dx.doi.org> prefix.

If you're reading the PDF e-book edition, and your PC is connected to the Internet, simply click on the link. You'll find *StatLinks* appearing in more OECD books.



From:
**Environmental Performance of Agriculture in
OECD Countries Since 1990**

Access the complete publication at:
<https://doi.org/10.1787/9789264040854-en>

Please cite this chapter as:

OECD (2008), "OECD Country Trends of Environmental Conditions related to Agriculture since 1990: Canada", in *Environmental Performance of Agriculture in OECD Countries Since 1990*, OECD Publishing, Paris.

DOI: <https://doi.org/10.1787/9789264040854-9-en>

This work is published under the responsibility of the Secretary-General of the OECD. The opinions expressed and arguments employed herein do not necessarily reflect the official views of OECD member countries.

This document and any map included herein are without prejudice to the status of or sovereignty over any territory, to the delimitation of international frontiers and boundaries and to the name of any territory, city or area.

You can copy, download or print OECD content for your own use, and you can include excerpts from OECD publications, databases and multimedia products in your own documents, presentations, blogs, websites and teaching materials, provided that suitable acknowledgment of OECD as source and copyright owner is given. All requests for public or commercial use and translation rights should be submitted to rights@oecd.org. Requests for permission to photocopy portions of this material for public or commercial use shall be addressed directly to the Copyright Clearance Center (CCC) at info@copyright.com or the Centre français d'exploitation du droit de copie (CFC) at contact@cfcopies.com.