OECD Series on Adverse Outcome Pathways No. 9

Adverse Outcome Pathway
on Androgen receptor
agonism leading
to reproductive dysfunction
(in repeat-spawning fish)

Daniel L. Villeneuve

<https://dx.doi.org/10.1787/b0c6838a-en>

*Adverse Outcome Pathway on Androgen
receptor agonism leading to reproductive dysfunction (in
repeat-spawning fish)*

Series on Adverse Outcome Pathways No. 9

Foreword

This Adverse Outcome Pathway (AOP) on Androgen receptor agonism leading to reproductive dysfunction (in repeat-spawning fish), has been developed under the auspices of the OECD AOP Development Programme, overseen by the Extended Advisory Group on Molecular Screening and Toxicogenomics (EAGMST), which is an advisory group under the Working Group of the National Coordinators for the Test Guidelines Programme (WNT). The AOP has been reviewed internally by the EAGMST, externally by experts nominated by the WNT, and has been endorsed by the WNT and the Working Party on Hazard Assessment (WPHA) in June 2018.

Through endorsement of this AOP, the WNT and the WPHA express confidence in the scientific review process that the AOP has undergone and accept the recommendation of the EAGMST that the AOP be disseminated publicly. Endorsement does not necessarily indicate that the AOP is now considered a tool for direct regulatory application.

The Joint Meeting of the Chemicals Committee and the Working Party on Chemicals, Pesticides and Biotechnology agreed to declassification of this AOP in September 2018.

This document is being published under the responsibility of the Joint Meeting of the Chemicals Committee and the Working Party on Chemicals, Pesticides and Biotechnology.

The outcome of the internal and external reviews are publicly available respectively in the [AOP Wiki](#) and the [eAOP Portal of the AOP Knowledge Base](#) at the following links: [\[internal review\]](#) [\[external review\]](#).

Table of contents

Foreword	1
Adverse Outcome Pathway on Androgen receptor agonism leading to reproductive dysfunction (in repeat-spawning fish)	5
Short Title: Androgen receptor agonism leading to reproductive dysfunction	5
Authors:	5
Abstract	6
Graphical Representation.....	7
Summary of the AOP	8
Molecular Initiating Events (MIEs), Key Events (KEs), Adverse Outcomes (AOs).....	8
Key Event Relationships (KERs).....	8
Stressors	9
17beta-Trenbolone	9
Overall Assessment of the AOP.....	10
Domain of Applicability	10
Life Stage Applicability	10
Taxonomic Applicability.....	10
Sex Applicability.....	10
Essentiality of the Key Events	11
Weight of Evidence Summary	11
Quantitative Consideration	14
References	15
Appendix 1: List of MIE(s), KE(s) and AO(s) in the AOP	23
List of MIE(s) in this AOP.....	23
Event: 25: Agonism, Androgen receptor.....	23
List of KE(s) in the AOP	30
Event: 129: Reduction, Gonadotropins, circulating concentrations	30
Event: 274: Reduction, Testosterone synthesis by ovarian theca cells	33
Event: 3: Reduction, 17beta-estradiol synthesis by ovarian granulosa cells	36
Event: 219: Reduction, Plasma 17beta-estradiol concentrations	39
Event: 285: Reduction, Vitellogenin synthesis in liver	42
Event: 221: Reduction, Plasma vitellogenin concentrations	45
Event: 309: Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development.....	47
Event: 78: Reduction, Cumulative fecundity and spawning	49
List of AO(s) in this AOP	51
Event: 360: Decrease, Population trajectory	51
Appendix 2: List of Key Event Relationships in the AOP	53
List of Adjacent Key Event Relationships.....	53
Relationship: 31: Agonism, Androgen receptor leads to Reduction, Gonadotropins, circulating concentrations	53

Relationship: 143: Reduction, Gonadotropins, circulating concentrations leads to Reduction, Testosterone synthesis by ovarian theca cells	56
Relationship: 302: Reduction, Testosterone synthesis by ovarian theca cells leads to Reduction, 17beta-estradiol synthesis by ovarian granulosa cells	58
Relationship: 5: Reduction, 17beta-estradiol synthesis by ovarian granulosa cells leads to Reduction, Plasma 17beta-estradiol concentrations	62
Relationship: 252: Reduction, Plasma 17beta-estradiol concentrations leads to Reduction, Vitellogenin synthesis in liver	67
Relationship: 315: Reduction, Vitellogenin synthesis in liver leads to Reduction, Plasma vitellogenin concentrations	71
Relationship: 255: Reduction, Plasma vitellogenin concentrations leads to Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development	74
Relationship: 337: Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development leads to Reduction, Cumulative fecundity and spawning	77
Relationship: 94: Reduction, Cumulative fecundity and spawning leads to Decrease, Population trajectory	80
List of Non Adjacent Key Event Relationships	83
Relationship: 32: Agonism, Androgen receptor leads to Reduction, Testosterone synthesis by ovarian theca cells	83
Relationship: 1384: Agonism, Androgen receptor leads to Reduction, 17beta-estradiol synthesis by ovarian granulosa cells	88
Relationship: 1385: Agonism, Androgen receptor leads to Reduction, Vitellogenin synthesis in liver	91
Relationship: 1386: Reduction, Plasma 17beta-estradiol concentrations leads to Reduction, Plasma vitellogenin concentrations	95
Appendix 3: Concordance Table.....	100

*Adverse Outcome Pathway on Androgen receptor agonism
leading to reproductive dysfunction (in repeat-spawning fish)*

Short Title: Androgen receptor agonism leading to reproductive dysfunction

Authors:

Dan Villeneuve, US EPA Mid-Continent Ecology Division

Abstract

This AOP details the linkage between binding and activation of androgen receptor (AR) in females and reductions in cumulative fecundity and spawning. AR-mediated activities are among the major concerns in endocrine disruptor screening programs. Cumulative fecundity is the apical endpoint considered in the OECD 229 Fish Short Term Reproduction Assay. It is also one of several variables with demographic significance in forecasting fish population trends. Therefore, this AOP supports the use of measures of AR activation as a means to identify chemicals with potential to adversely affect fish populations. At present, this AOP is largely supported by evidence from small laboratory model fish species. While many aspects of the biology underlying this AOP are largely conserved across oviparous vertebrates, its relevance to vertebrate classes other than fish, or to fish species employing different reproductive strategies has not been established. Thus, the applicability domain should be carefully considered when evaluating fit-for-purpose.

Graphical Representation

Summary of the AOP

Molecular Initiating Events (MIEs), Key Events (KEs), Adverse Outcomes (AOs)

Sequence	Type	Event ID	Title	Short name
1	MIE	25	Agonism, Androgen receptor	Agonism, Androgen receptor
2	KE	129	Reduction, Gonadotropins, circulating concentrations	Reduction, Gonadotropins, circulating concentrations
3	KE	274	Reduction, Testosterone synthesis by ovarian theca cells	Reduction, Testosterone synthesis by ovarian theca cells
4	KE	3	Reduction, 17beta-estradiol synthesis by ovarian granulosa cells	Reduction, 17beta-estradiol synthesis by ovarian granulosa cells
5	KE	219	Reduction, Plasma 17beta-estradiol concentrations	Reduction, Plasma 17beta-estradiol concentrations
6	KE	285	Reduction, Vitellogenin synthesis in liver	Reduction, Vitellogenin synthesis in liver
7	KE	221	Reduction, Plasma vitellogenin concentrations	Reduction, Plasma vitellogenin concentrations
8	KE	309	Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development	Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development
9	KE	78	Reduction, Cumulative fecundity and spawning	Reduction, Cumulative fecundity and spawning
10	AO	360	Decrease, Population trajectory	Decrease, Population trajectory

Key Event Relationships (KERs)

Upstream Event	Relationship Type	Downstream Event	Evidence	Quantitative Understanding
Agonism, Androgen receptor	adjacent	Reduction, Gonadotropins, circulating concentrations	Low	Low
Reduction, Gonadotropins, circulating concentrations	adjacent	Reduction, Testosterone synthesis by ovarian theca cells	High	Low
Reduction, Testosterone synthesis by ovarian theca cells	adjacent	Reduction, 17beta-estradiol synthesis by ovarian granulosa cells	High	Low
Reduction, 17beta-estradiol synthesis by ovarian granulosa cells	adjacent	Reduction, Plasma 17beta-estradiol concentrations	High	Low
Reduction, Plasma 17beta-estradiol concentrations	adjacent	Reduction, Vitellogenin synthesis in liver	High	Moderate

Reduction, Vitellogenin synthesis in liver	adjacent	Reduction, Plasma vitellogenin concentrations	High	Moderate
Reduction, Plasma vitellogenin concentrations	adjacent	Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development	Moderate	Low
Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development	adjacent	Reduction, Cumulative fecundity and spawning	Moderate	Moderate
Reduction, Cumulative fecundity and spawning	adjacent	Decrease, Population trajectory	Moderate	Moderate
Agonism, Androgen receptor	non-adjacent	Reduction, Testosterone synthesis by ovarian theca cells	Moderate	Low
Agonism, Androgen receptor	non-adjacent	Reduction, 17beta-estradiol synthesis by ovarian granulosa cells	Moderate	Low
Agonism, Androgen receptor	non-adjacent	Reduction, Vitellogenin synthesis in liver	High	Low
Reduction, Plasma 17beta-estradiol concentrations	non-adjacent	Reduction, Plasma vitellogenin concentrations	High	Moderate

Stressors

Name	Evidence
17beta-Trenbolone	High

17beta-Trenbolone

17β-trenbolone is a prototypical stressor for this AOP.

Overall Assessment of the AOP

Domain of Applicability

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Pimephales promelas	Pimephales promelas		NCBI

Sex Applicability

Sex	Evidence
Female	

Domain(s) of Applicability

Chemical: This AOP applies to non-aromatizable androgens. Compounds which can bind the AR in vitro, but are converted to high potency estrogens in vivo through aromatization do not produce the profile of effects described in the present AOP (e.g., methyltestosterone [Ankley et al. 2001; Pawlowski et al. 2004]; androstenedione [OECD 2007]).

Sex: The AOP applies to females only.

Life stages: The relevant life stages for this AOP are reproductively mature adults. This AOP does not apply to adult stages that lack a sexually mature ovary, for example as a result of seasonal or environmentally-induced gonadal senescence (i.e., through control of temperature, photo-period, etc. in a laboratory setting).

Taxonomic: At present, the assumed taxonomic applicability domain of this AOP is iteroparous teleost fish species.

- However, to date the majority of toxicological data on which this AOP is based has been limited to several small fish species, fathead minnow (*Pimephales promelas*), Japanese medaka (*Oryzias latipes*), and mummichog (*Fundulus heteroclitus*) with asynchronous oocyte development and a repeat spawning reproductive strategy.
- Species dependent differences in endocrine feedback responses, likely associated with different reproductive strategies, have been reported. Thus, the applicability domain may prove more restricted than currently assumed. In particular, the applicability to fish species with synchronous or group synchronous oocyte development patterns (see Wallace and Selman 1981) is unclear.
- European eel may be an exception to the generalizability of the negative feedback response to a non-aromatizable xenoandrogen (Huang et al. 1997).

- Reductions in plasma VTG concentrations and/or hepatic VTG mRNA abundance in females following exposure to 17 β -trenbolone has been observed in *Pimephales promelas*, *Oryzias latipes*, *Danio rerio*, (Seki et al. 2006), *Cyprinodon variegatus* (Hemmer et al. 2008), *Gambusia holbrooki* and *Gambusia affinis* (Brockmeier et al. 2013)

[Assessment provided by Ioanna Katsiadaki - reviewer]: This is restricted clearly to female fish only as adversity is linked to reduced oestrogen synthesis (via reduced androgen synthesis); it is also limited to fully reproductive mature fish (not fish entering puberty or juvenile fish) and importantly is limited to fish that once they reach sexual maturity they spawn constantly. The latter is a reproductive strategy employed by fish that tend to occupy tropical areas (around the equator). Unfortunately, most fish species have different reproductive strategies (annual life cycle) hence the level of gonadotropin expression (and consequently steroid production) is regulated by photoperiodic and temperature changes throughout the year. Even if a negative feedback mechanism operates in all of these species and in all life stages (which is certainly not the case) we still need to establish what is the relative strength of the AR agonist induced negative feedback to the environment-induced stimulation of gonadotropins! This link has never been studied and is critical if we really mean to protect wildlife.

Essentiality of the Key Events

- In general, few studies have directly addressed the essentiality of the proposed sequence of key events.
- Ekman et al. 2011 provide evidence that in fathead minnow, cessation of trenbolone exposure resulted in recovery of plasma E2 and VTG concentrations which were depressed by continuous exposure to 17 β trenbolone. This provides some support for the essentiality of these two key events.
- There are no known reports of 17 β -trenbolone directly inhibiting steroid biosynthesis. When tested in an in vitro steroidogenesis assay using H295R adrenal carcinoma cells, trenbolone caused a concentration-dependent increase in estradiol production, as opposed to any reductions in steroid hormone concentrations, an effect that was concurrent with increased transcription of CYP19 (aromatase) in the cell line (Gracia et al. 2007).

Weight of Evidence Summary

Biological Plausibility

- The biochemistry of steroidogenesis and the predominant role of the gonad in synthesis of the sex steroids are well established.
- Similarly, the role of E2 as the major regulator of hepatic vitellogenin production is widely documented in the literature.
- The direct link between reduced VTG concentrations in the plasma and reduced uptake into oocytes is highly plausible, as the plasma is the primary source of the VTG.
- The direct connection between reduced VTG uptake and impaired spawning/reduced cumulative fecundity is more tentative. It is not clear, for instance whether impaired VTG uptake limits oocyte growth and failure to reach a critical size in turn impairs physical or inter-cellular signaling processes that promote release of the oocyte from the surrounding follicles. In at least one

experiment, oocytes with similar size to vitellogenic oocytes, but lacking histological staining characteristic of vitellogenic oocytes was observed (R. Johnson, personal communication). At present, the link between reductions in circulating VTG concentrations and reduced cumulative fecundity are best supported by the correlation between those endpoints across multiple experiments, including those that impact VTG via other molecular initiating events (Miller et al. 2007).

- At present, negative feedback is the most biologically plausible explanation for the reductions in ex vivo T and E2 production following exposure to 17 β -trenbolone. There are no known reports of 17 β -trenbolone directly inhibiting steroid biosynthesis. When tested in an in vitro steroidogenesis assay using H295R adrenal carcinoma cells, trenbolone caused a concentration-dependent increase in estradiol production, as opposed to any reductions in steroid hormone concentrations, an effect that was concurrent with increased transcription of CYP19 (aromatase) in the cell line (Gracia et al. 2007). Given the lack of any established direct effect on steroidogenic enzyme activity, negative feedback is currently the most likely explanation for the consistent effects observed in vivo. That said, many uncertainties regarding the exact mechanisms through which an exogenous, non-aromatizable, AR agonist elicits negative feedback remain.

Concordance of dose-response relationships: See Concordance Table in [Appendix 3](#).

There are a limited number of studies in which multiple key events were considered in the same study following exposure to known, non-aromatizable, AR agonists. These were considered the most useful for evaluating the concordance of dose-response relationships. In general, effects on downstream key events occurred at concentrations equal to or greater than those at which upstream events occurred. For exposures to 17 β -trenbolone, key events related to steroid production and circulating estradiol and vitellogenin concentrations were impacted at the same dose at which effects on cumulative fecundity were observed. Effects on vitellogenin transcription were only observed at greater concentrations, but data for comparable species and dose ranges were unavailable at present. For two other AR agonists tested in fish, available studies examined a single time-point only. Consequently, it was unclear whether lower effect concentrations for certain downstream KEs, relative to upstream were due to a lack of dose-response concordance, or due to decreased sensitivity of the upstream later in the exposure time-course.

While not directly addressing dose-response concordance, the dependence of the key events on the concentration of the androgen agonist has been established for all key events starting at and down-stream of reduced T synthesis. However, to date we are not aware of any studies that have established a concentration-response relationship between exposure to non-endogenous AR agonists (e.g., xenobiotics, pharmaceuticals) and circulating gonadotropin concentrations in fish or other vertebrates.

- Exposure of female fathead minnows to the AR agonist 17 β -trenbolone for 21 d caused concentration-dependent reductions in circulating T, E2, and VTG concentrations over a range from 0.005 to 0.5 μ g/L. The concentration response for all three variables had a “U”-shaped concentration response curve which may indicate concentration-dependent differences in the feedback response and/or compensatory processes. Histological evidence of reduced VTG uptake and reduced gonad stage were evident, although the concentration-response of histological effects was not determined. Despite the “U”-shaped concentration-

response at the biochemical level, concentration-dependent reductions in cumulative fecundity were observed (Ankley et al. 2003). Effective concentrations were consistent with those causing phenotypic masculinisation in female fish.

- Jensen et al. (2006) also demonstrated concentration-dependent reductions in circulating T, E2, and VTG following 21 d of in vivo exposure to 17 α -trenbolone (Jensen et al. 2006).
- In a time-course experiment in which female fathead minnows were exposed to 33 or 472 ng 17 β -trenbolone/L ex vivo T, ex vivo E2, plasma E2, and plasma VTG all showed concentration-dependent reductions that were consistent with the AOP (Ekman et al. 2011).
- Exposure of female fathead minnows to spironolactone, a pharmaceutical that binds the fathead minnow AR, for 21 d caused concentration-dependent reductions in cumulative fecundity, plasma VTG and VTG mRNA expression, and plasma E2 concentrations. The frequency and severity of females with decreased yolk accumulation, and increased oocyte atresia was concentration-dependent. The chemical also induced phenotypic masculinisation in female fish. (Lalone et al. 2013).
- Exposure of female medaka to spironolactone caused concentration-dependent reductions in cumulative fecundity and VTG mRNA expression (impacts on steroid hormone concentrations were not measured). Spironolactone also caused phenotypic masculinisation of female medaka (Lalone et al. 2013).

Temporal concordance among the key events and adverse effect: Temporal concordance between activation of the AR as a nuclear transcription factor and onset of a negative feedback response resulting in decreased gonadotropin secretion has not been established. Temporal concordance of the key events starting with reduced T biosynthesis and proceeding through reductions in plasma vitellogenin has been established ([Appendix 3](#)). Temporal concordance beyond the key event of reductions in plasma vitellogenin has not been established, in large part due to disconnect in the time-scales over which the events can be measured. For example, most small fish used in reproductive toxicity testing can spawn anywhere from once daily to several days per week. Given the variability in daily spawning rates, it is neither practical nor effective to evaluate cumulative fecundity at a time scale shorter than roughly a week. Since the impacts at lower levels of biological organization can be detected within hours of exposure, lack of impact on cumulative fecundity before the other key events are impacted cannot be effectively measured. Overall, among those key events whose temporal concordance can reasonably be evaluated based on currently available data, the temporal profile observed is consistent with the AOP.

Consistency: We are aware of no cases where the pattern of key events described was observed without also observing a significant impact on cumulative fecundity. Due to variability in the cumulative fecundity endpoint and potential compensatory responses (Villeneuve et al. 2009; Villeneuve et al. 2013; Ankley et al. 2009b; Zhang et al. 2008; Ekman et al. 2012), the cumulative fecundity endpoint can be less sensitive than key events measured at lower levels of biological organization. Nonetheless, the occurrence of the final adverse outcome when the other key events are observed is very consistent. The final adverse effect is not specific to this AOP. Many of the key events included in this AOP overlap with AOPs linking other molecular initiating events to reproductive dysfunction in small fish.

- In general, there is a consistent body of evidence linking exposure to an AR agonist to decreased T synthesis, E2 synthesis, circulating E2 and VTG concentrations, and

cumulative fecundity in female fish. For example, the association between 17 β -trenbolone exposure and reduced vitellogenin concentrations in females has been replicated in over a dozen independent experiments (Ekman et al. 2011; Ankley et al. 2003; Jensen et al. 2006; Ankley et al. 2010; Hemmer et al. 2008; Seki et al. 2006; Brockmeier et al. 2013). However, relatively few exogenous, non-aromatizable, AR agonists have been tested. Other than recent work with spironolactone (Lalone et al. 2013), we are not aware of the profile of responses being demonstrated for other AR agonists.

Uncertainties, inconsistencies, and data gaps: There are three major areas of uncertainty and data gaps in the current AOP:

- First, there remains considerable uncertainty as to the specific mechanism(s) through which AR agonism elicits a negative feedback response at the level of the hypothalamus and/or pituitary. There is also a substantial data gap relative to establishing that exposure to an AR agonist like 17 β -trenbolone causes concentration-dependent reductions in circulating gonadotropins. That uncertainty is amplified further by the variation in feedback control along the endocrine axis for fish species employing different reproductive strategies. For example, gonadotropin regulation may be very different in species with synchronous oocyte maturation and annual or once per life-time reproductive strategies. Thus, there are considerable uncertainties related to the taxonomic relevance of this AOP to a broader range of fish species or other vertebrates.
- The second major uncertainty in this AOP relates to whether there is a direct biological linkage between impaired VTG uptake into oocytes and impaired spawning/reduced cumulative fecundity. Plausible biological connections have been hypothesised, but have not yet been tested experimentally.
- A third uncertainty pertains to the chemical domain of applicability. In vivo, a number of chemicals that are detected as androgens in in vitro screening assays such as receptor binding assays or ligand-activated transcriptional assay can be aromatized to functional estrogens. Thus, in vivo such compounds may produce a profile of effects more consistent with estrogen receptor activation than AR activation or may produced mixed effects characteristic of either estrogen or androgen exposures (e.g., Pawlowski et al. 2004; Hornung et al. 2004). Examples of such aromatizable androgens include, testosterone, methyltestosterone, and androstenedione. Consequently, caution is warranted in applying this AOP based on in vitro screening data alone, without consideration for possible conversion to estrogens.

Quantitative Consideration

Assessment of quantitative understanding of the AOP: At present, the quantitative understanding of the AOP is insufficient to directly link a measure of chemical potency as an AR agonist (e.g., as measured in a transcriptional activation assay) to a predicted effect concentration at the level of cumulative fecundity. However, a number of mechanistic and statistical models are sufficiently developed to facilitate predictions of cumulative outcomes based on intermediate key event measurements such as circulating vitellogenin concentrations. Because the current models were developed based on a fairly limited range of model compounds and species, the general applicability and degree of accuracy and precision in the model-derived predictions remain uncertain.

References

- Amano M, Iigo M, Ikuta K, Kitamura S, Yamada H, Yamamori K. 2000. Roles of melatonin in gonadal maturation of underyearling precocious male masu salmon. *General and comparative endocrinology* 120(2): 190-197.
- Ankley GT, Bencic D, Cavallin JE, Jensen KM, Kahl MD, Makynen EA, et al. 2009. Dynamic nature of alterations in the endocrine system of fathead minnows exposed to the fungicide prochloraz. *Toxicol Sci* 112(2): 344-353.
- Ankley GT, Cavallin JE, Durhan EJ, Jensen KM, Kahl MD, Makynen EA, et al. 2012. A time-course analysis of effects of the steroidogenesis inhibitor ketoconazole on components of the hypothalamic-pituitary-gonadal axis of fathead minnows. *Aquatic toxicology* 114-115: 88-95.
- Ankley GT, Jensen KM, Durhan EJ, Makynen EA, Butterworth BC, Kahl MD, et al. 2005. Effects of two fungicides with multiple modes of action on reproductive endocrine function in the fathead minnow (*Pimephales promelas*). *Toxicol Sci* 86(2): 300-308.
- Ankley GT, Jensen KM, Kahl MD, Durhan EJ, Makynen EA, Cavallin JE, et al. 2010. Use of chemical mixtures to differentiate mechanisms of endocrine action in a small fish model. *Aquatic toxicology* 99(3): 389-396.
- Ankley GT, Jensen KM, Kahl MD, Korte JJ, Makynen EA. 2001. Description and evaluation of a short-term reproduction test with the fathead minnow (*Pimephales promelas*). *Environ Toxicol Chem* 20:1276-1290.
- Ankley GT, Jensen KM, Kahl MD, Makynen EA, Blake LS, Greene KJ, et al. 2007. Ketoconazole in the fathead minnow (*Pimephales promelas*): reproductive toxicity and biological compensation. *Environ Toxicol Chem* 26(6): 1214-1223.
- Ankley GT, Jensen KM, Makynen EA, Kahl MD, Korte JJ, Hornung MW, et al. 2003. Effects of the androgenic growth promoter 17-b-trenbolone on fecundity and reproductive endocrinology of the fathead minnow. *Environmental Toxicology and Chemistry* 22(6): 1350-1360.
- Ankley GT, Kahl MD, Jensen KM, Hornung MW, Korte JJ, Makynen EA, et al. 2002. Evaluation of the aromatase inhibitor fadrozole in a short-term reproduction assay with the fathead minnow (*Pimephales promelas*). *Toxicological Sciences* 67: 121-130.
- Ankley GT, Miller DH, Jensen KM, Villeneuve DL, Martinovic D. 2008. Relationship of plasma sex steroid concentrations in female fathead minnows to reproductive success and population status. *Aquatic toxicology* 88(1): 69-74.
- Araki N, Ohno K, Nakai M, Takeyoshi M, Iida M. 2005. Screening for androgen receptor activities in 253 industrial chemicals by in vitro reporter gene assays using AR-EcoScreen cells. *Toxicology in vitro : an international journal published in association with BIBRA* 19(6): 831-842.
- Arukwe A, Goksøyr A. 2003. Eggshell and egg yolk proteins in fish: hepatic proteins for the next generation: oogenetic, population, and evolutionary implications of endocrine disruption. *Comparative Hepatology* 2(4): 1-21.

- Baker ME. 1997. Steroid receptor phylogeny and vertebrate origins. *Molecular and cellular endocrinology* 135(2): 101-107.
- Baker ME. 2011. Origin and diversification of steroids: co-evolution of enzymes and nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 14-20.
- Benninghoff AD, Thomas P. 2006. Gonadotropin regulation of testosterone production by primary cultured theca and granulosa cells of Atlantic croaker: I. Novel role of CaMKs and interactions between calcium- and adenylyl cyclase-dependent pathways. *General and comparative endocrinology* 147(3): 276-287.
- Biales AD, Bencic DC, Lazorchak JL, Lattier DL. 2007. A quantitative real-time polymerase chain reaction method for the analysis of vitellogenin transcripts in model and nonmodel fish species. *Environ Toxicol Chem* 26(12): 2679-2686.
- Bohl CE, Chang C, Mohler ML, Chen J, Miller DD, Swaan PW, et al. 2004. A ligand-based approach to identify quantitative structure-activity relationships for the androgen receptor. *Journal of medicinal chemistry* 47(15): 3765-3776.
- Bowman CJ, Kroll KJ, Hemmer MJ, Folmar LC, Denslow ND. 2000. Estrogen-induced vitellogenin mRNA and protein in sheepshead minnow (*Cyprinodon variegatus*). *General and comparative endocrinology* 120(3): 300-313.
- Breen MS, Villeneuve DL, Breen M, Ankley GT, Conolly RB. 2007. Mechanistic computational model of ovarian steroidogenesis to predict biochemical responses to endocrine active compounds. *Annals of biomedical engineering* 35(6): 970-981.
- Brockmeier EK, Ogino Y, Iguchi T, Barber DS, Denslow ND. 2013. Effects of 17beta-trenbolone on Eastern and Western mosquitofish (*Gambusia holbrooki* and *G. affinis*) anal fin growth and gene expression patterns. *Aquatic toxicology* 128-129: 163-170.
- Campbell BK, Baird DT, Webb R. 1998. Effects of dose of LH on androgen production and luteinization of ovine theca cells cultured in a serum-free system. *Journal of reproduction and fertility* 112(1): 69-77.
- Centenera MM, Harris JM, Tilley WD, Butler LM. 2008. The contribution of different androgen receptor domains to receptor dimerization and signaling. *Molecular endocrinology* 22(11): 2373-2382.
- Cheng GF, Yuen CW, Ge W. 2007. Evidence for the existence of a local activin follistatin negative feedback loop in the goldfish pituitary and its regulation by activin and gonadal steroids. *The Journal of endocrinology* 195(3): 373-384.
- Claessens F, Denayer S, Van Tilborgh N, Kerkhofs S, Helsen C, Haelens A. 2008. Diverse roles of androgen receptor (AR) domains in AR-mediated signaling. *Nuclear receptor signaling* 6: e008.
- Cutress ML, Whitaker HC, Mills IG, Stewart M, Neal DE. 2008. Structural basis for the nuclear import of the human androgen receptor. *Journal of cell science* 121(Pt 7): 957-968.
- Eick GN, Thornton JW. 2011. Evolution of steroid receptors from an estrogen-sensitive ancestral receptor. *Molecular and cellular endocrinology* 334(1-2): 31-38.
- Ekman DR, Hartig PC, Cardon M, Skelton DM, Teng Q, Durhan EJ, et al. 2012. Metabolite profiling and a transcriptional activation assay provide direct evidence of androgen receptor antagonism by bisphenol A in fish. *Environmental science & technology* 46(17): 9673-9680.

- Ekman DR, Villeneuve DL, Teng Q, Ralston-Hooper KJ, Martinovic-Weigelt D, Kahl MD, et al. 2011. Use of gene expression, biochemical and metabolite profiles to enhance exposure and effects assessment of the model androgen 17beta-trenbolone in fish. *Environmental toxicology and chemistry / SETAC* 30(2): 319-329.
- Eppig JJ. 1994. Further reflections on culture systems for the growth of oocytes in vitro. *Human reproduction* 9(6): 974-976.
- Genovese G, Regueira M, Piazza Y, Towle DW, Maggese MC, Lo Nostro F. 2012. Time-course recovery of estrogen-responsive genes of a cichlid fish exposed to waterborne octylphenol. *Aquatic toxicology* 114-115: 1-13.
- Gopurappilly R, Ogawa S, Parhar IS. 2013. Functional significance of GnRH and kisspeptin, and their cognate receptors in teleost reproduction. *Frontiers in endocrinology* 4: 24.
- Govoroun M, Chyb J, Breton B. 1998. Immunological cross-reactivity between rainbow trout GTH I and GTH II and their alpha and beta subunits: application to the development of specific radioimmunoassays. *General and comparative endocrinology* 111(1): 28-37.
- Gracia T, Hilscherova K, Jones PD, Newsted JL, Higley EB, Zhang X, et al. 2007. Modulation of steroidogenic gene expression and hormone production of H295R cells by pharmaceuticals and other environmentally active compounds. *Toxicology and applied pharmacology* 225(2): 142-153.
- Habibi HR, Huggard DL. 1998. Testosterone regulation of gonadotropin production in goldfish. *Comparative biochemistry and physiology Part C, Pharmacology, toxicology & endocrinology* 119(3): 339-344.
- Havelock JC, Rainey WE, Carr BR. 2004. Ovarian granulosa cell lines. *Molecular and cellular endocrinology* 228(1-2): 67-78.
- Heemers HV, Tindall DJ. 2007. Androgen receptor (AR) coregulators: a diversity of functions converging on and regulating the AR transcriptional complex. *Endocrine reviews* 28(7): 778-808.
- Hemmer MJ, Cripe GM, Hemmer BL, Goodman LR, Salinas KA, Fournie JW, et al. 2008. Comparison of estrogen-responsive plasma protein biomarkers and reproductive endpoints in sheepshead minnows exposed to 17beta-trenbolone. *Aquatic toxicology* 88(2): 128-136.
- Hong H, Fang H, Xie Q, Perkins R, Sheehan DM, Tong W. 2003. Comparative molecular field analysis (CoMFA) model using a large diverse set of natural, synthetic and environmental chemicals for binding to the androgen receptor. *SAR and QSAR in environmental research* 14(5-6): 373-388.
- Hornung MW, Jensen KM, Korte JJ, Kahl MD, Durhan EJ, Denny JS, Henry TR, Ankley GT. 2004. Mechanistic basis for estrogenic effects in fathead minnow (*Pimephales promelas*) following exposure to the androgen 17alpha-methyltestosterone: conversion of 17alpha-methyltestosterone to 17alpha-methylestradiol. 66: 15-23.
- Iguchi T, Irie F, Urushitani H, Tooi O, Kawashima Y, Roberts M, et al. 2006. Availability of in vitro vitellogenin assay for screening of estrogenic and anti-estrogenic activities of environmental chemicals. *Environ Sci* 13(3): 161-183.
- Jamnongjit M, Hammes SR. 2005. Oocyte maturation: the coming of age of a germ cell. *Seminars in reproductive medicine* 23(3): 234-241.

- Jensen K, Korte J, Kahl M, Pasha M, Ankley G. 2001. Aspects of basic reproductive biology and endocrinology in the fathead minnow (*Pimephales promelas*). *Comparative Biochemistry and Physiology Part C* 128: 127-141.
- Jensen KM, Makynen EA, Kahl MD, Ankley GT. 2006. Effects of the feedlot contaminant 17alpha-trenbolone on reproductive endocrinology of the fathead minnow. *Environmental science & technology* 40(9): 3112-3117.
- Jolly C, Katsiadaki I, Le Belle N, Mayer I, Dufour S. 2006. Development of a stickleback kidney cell culture assay for the screening of androgenic and anti-androgenic endocrine disrupters. *Aquatic toxicology* 79(2): 158-166.
- Kah O, Pontet A, Nunez Rodriguez J, Calas A, Breton B. 1989. Development of an enzyme-linked immunosorbent assay for goldfish gonadotropin. *Biology of reproduction* 41(1): 68-73.
- Kim TS, Yoon CY, Jung KK, Kim SS, Kang IH, Baek JH, et al. 2010. In vitro study of Organization for Economic Co-operation and Development (OECD) endocrine disruptor screening and testing methods- establishment of a recombinant rat androgen receptor (rrAR) binding assay. *The Journal of toxicological sciences* 35(2): 239-243.
- Korte JJ, Kahl MD, Jensen KM, Mumtaz SP, Parks LG, LeBlanc GA, et al. 2000. Fathead minnow vitellogenin: complementary DNA sequence and messenger RNA and protein expression after 17B-estradiol treatment. *Environmental Toxicology and Chemistry* 19(4): 972-981.
- Lalone CA, Villeneuve DL, Cavallin JE, Kahl MD, Durhan EJ, Makynen EA, et al. 2013. Cross species sensitivity to a novel androgen receptor agonist of environmental concern, spironolactone. *Environ. Toxicol. Chem.* 32: 2528-2541.
- Leino R, Jensen K, Ankley G. 2005. Gonadal histology and characteristic histopathology associated with endocrine disruption in the adult fathead minnow. *Environmental Toxicology and Pharmacology* 19: 85-98.
- Levavi-Sivan B, Bogerd J, Mananos EL, Gomez A, Lareyre JJ. 2010. Perspectives on fish gonadotropins and their receptors. *General and comparative endocrinology* 165(3): 412-437.
- Li Z, Kroll KJ, Jensen KM, Villeneuve DL, Ankley GT, Brian JV, et al. 2011a. A computational model of the hypothalamic: pituitary: gonadal axis in female fathead minnows (*Pimephales promelas*) exposed to 17alpha-ethynylestradiol and 17beta-trenbolone. *BMC systems biology* 5: 63.
- Li Z, Villeneuve DL, Jensen KM, Ankley GT, Watanabe KH. 2011b. A computational model for asynchronous oocyte growth dynamics in a batch-spawning fish. *Can J Fish Aquat Sci* 68: 1528-1538.
- Magoffin DA. 2005. Ovarian theca cell. *The international journal of biochemistry & cell biology* 37(7): 1344-1349.
- Mak P, Cruz FD, Chen S. 1999. A yeast screen system for aromatase inhibitors and ligands for androgen receptor: yeast cells transformed with aromatase and androgen receptor. *Environmental health perspectives* 107(11): 855-860.
- Markov GV, Laudet V. 2011. Origin and evolution of the ligand-binding ability of nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 21-30.

- McMaster ME MK, Jardine JJ, Robinson RD, Van Der Kraak GJ. 1995. Protocol for measuring in vitro steroid production by fish gonadal tissue. Canadian Technical Report of Fisheries and Aquatic Sciences 1961 1961: 1-78.
- Miller DH, Ankley GT. 2004. Modeling impacts on populations: fathead minnow (*Pimephales promelas*) exposure to the endocrine disruptor 17 β -trenbolone as a case study. *Ecotoxicology and Environmental Safety* 59: 1-9.
- Miller DH, Jensen KM, Villeneuve DL, Kahl MD, Makynen EA, Durhan EJ, et al. 2007. Linkage of biochemical responses to population-level effects: a case study with vitellogenin in the fathead minnow (*Pimephales promelas*). *Environ Toxicol Chem* 26(3): 521-527.
- Miller DH, Tietge JE, McMaster ME, Munkittrick KR, Xia X, Ankley GT. 2013. Assessment of Status of White Sucker (*Catostomus Commersoni*) Populations Exposed to Bleached Kraft Pulp Mill Effluent. *Environmental toxicology and chemistry / SETAC*.
- Miller WL, Strauss JF, 3rd. 1999. Molecular pathology and mechanism of action of the steroidogenic acute regulatory protein, StAR. *The Journal of steroid biochemistry and molecular biology* 69(1-6): 131-141.
- Miller WL. 1988. Molecular biology of steroid hormone synthesis. *Endocrine reviews* 9(3): 295-318.
- Murphy CA, Rose KA, Rahman MS, Thomas P. 2009. Testing and applying a fish vitellogenesis model to evaluate laboratory and field biomarkers of endocrine disruption in Atlantic croaker (*Micropogonias undulatus*) exposed to hypoxia. *Environmental toxicology and chemistry / SETAC* 28(6): 1288-1303.
- Murphy CA, Rose KA, Thomas P. 2005. Modeling vitellogenesis in female fish exposed to environmental stressors: predicting the effects of endocrine disturbance due to exposure to a PCB mixture and cadmium. *Reproductive toxicology* 19(3): 395-409.
- Nagahama Y, Yoshikumi M, Yamashita M, Sakai N, Tanaka M. 1993. Molecular endocrinology of oocyte growth and maturation in fish. *Fish Physiology and Biochemistry* 11: 3-14.
- Navas JM, Segner H. 2006. Vitellogenin synthesis in primary cultures of fish liver cells as endpoint for in vitro screening of the (anti)estrogenic activity of chemical substances. *Aquat Toxicol* 80(1): 1-22.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Norris JD, Joseph JD, Sherk AB, Juzumiene D, Turnbull PS, Rafferty SW, et al. 2009. Differential presentation of protein interaction surfaces on the androgen receptor defines the pharmacological actions of bound ligands. *Chemistry & biology* 16(4): 452-460.
- Oakley AE, Clifton DK, Steiner RA. 2009. Kisspeptin signaling in the brain. *Endocrine reviews* 30(6): 713-743.
- OECD. 2012. Test No. 229: Fish Short Term Reproduction Assay. Paris, France: Organization for Economic Cooperation and Development.
- Olsson P-E, Berg A, von Hofsten J, Grahn B, Hellqvist A, Larsson A, et al. 2005. Molecular cloning and characterization of a nuclear androgen receptor activated by 11-ketotestosterone. *Reproductive Biology and Endocrinology* 3: 1-17.

- Organization for Economic Cooperation and Development. 2007. Report of Eight 21-day Fish Endocrine Screening Assays With Additional Test Substances for Phase-3 of the OECD Validation Program: Studies with Octylphenol in the Fathead Minnow (*Pimephales promelas*) and Zebrafish (*Danio rerio*) and with Sodium Pentachlorophenol and Androstenedione in the Fathead Minnow (*Pimephales promelas*). Phase-3 OECD 21-day Fish Screening Assay Validation Report Additional Test Substances Studies. Paris, France.
- Pawlowski S, Sauer A, Shears JA, Tyler CR, Braunbeck T. 2004. Androgenic and estrogenic effects of the synthetic androgen 17 α -methyltestosterone on sexual development and reproductive performance in the fathead minnow (*Pimephales promelas*) determined using the gonadal recrudescence assay. *Aquat Toxicol* 68:277-291.
- Payne AH, Hales DB. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine reviews* 25(6): 947-970.
- Prat F, Sumpter JP, Tyler CR. 1996. Validation of radioimmunoassays for two salmon gonadotropins (GTH I and GTH II) and their plasma concentrations throughout the reproductive cycle in male and female rainbow trout (*Oncorhynchus mykiss*). *Biology of reproduction* 54(6): 1375-1382.
- Prescott J, Coetzee GA. 2006. Molecular chaperones throughout the life cycle of the androgen receptor. *Cancer letters* 231(1): 12-19.
- Quignot N, Bois FY. 2013. A computational model to predict rat ovarian steroid secretion from in vitro experiments with endocrine disruptors. *PloS one* 8(1): e53891.
- Schmid T, Gonzalez-Valero J, Ruffli H, Dietrich DR. 2002. Determination of vitellogenin kinetics in male fathead minnows (*Pimephales promelas*). *Toxicol Lett* 131(1-2): 65-74.
- Schmieder P, Tapper M, Linnun A, Denny J, Kolanczyk R, Johnson R. 2000. Optimization of a precision-cut trout liver tissue slice assay as a screen for vitellogenin induction: comparison of slice incubation techniques. *Aquat Toxicol* 49(4): 251-268.
- Schultz IR, Orner G, Merdink JL, Skillman A. 2001. Dose-response relationships and pharmacokinetics of vitellogenin in rainbow trout after intravascular administration of 17 α -ethynylestradiol. *Aquatic toxicology* 51(3): 305-318.
- Seki M, Fujishima S, Nozaka T, Maeda M, Kobayashi K. 2006. Comparison of response to 17 beta-estradiol and 17 beta-trenbolone among three small fish species. *Environmental toxicology and chemistry / SETAC* 25(10): 2742-2752.
- Serafimova R, Walker J, Mekenyan O. 2002. Androgen receptor binding affinity of pesticide "active" formulation ingredients. QSAR evaluation by COREPA method. SAR and QSAR in environmental research 13(1): 127-134.
- Shoemaker JE, Gayen K, Garcia-Reyero N, Perkins EJ, Villeneuve DL, Liu L, et al. 2010. Fathead minnow steroidogenesis: in silico analyses reveals tradeoffs between nominal target efficacy and robustness to cross-talk. *BMC systems biology* 4: 89.
- Skolness SY, Blanksma CA, Cavallin JE, Churchill JJ, Durhan EJ, Jensen KM, et al. 2013. Propiconazole Inhibits Steroidogenesis and Reproduction in the Fathead Minnow (*Pimephales promelas*). *Toxicological sciences: an official journal of the Society of Toxicology* 132(2): 284-297.
- Skolness SY, Durhan EJ, Garcia-Reyero N, Jensen KM, Kahl MD, Makynen EA, et al. 2011. Effects of a short-term exposure to the fungicide prochloraz on endocrine function

and gene expression in female fathead minnows (*Pimephales promelas*). *Aquat Toxicol* 103(3-4): 170-178.

- Sower SA, Freamat M, Kavanaugh SI. 2009. The origins of the vertebrate hypothalamic-pituitary-gonadal (HPG) and hypothalamic-pituitary-thyroid (HPT) endocrine systems: new insights from lampreys. *General and comparative endocrinology* 161(1): 20-29.
- Sperry TS, Thomas P. 1999. Identification of two nuclear androgen receptors in kelp bass (*Paralabrax clathratus*) and their binding affinities for xenobiotics: comparison with Atlantic croaker (*Micropogonias undulatus*) androgen receptors. *Biology of reproduction* 61(4): 1152-1161.
- Sun L, Wen L, Shao X, Qian H, Jin Y, Liu W, et al. 2010. Screening of chemicals with anti-estrogenic activity using in vitro and in vivo vitellogenin induction responses in zebrafish (*Danio rerio*). *Chemosphere* 78(7): 793-799.
- Sun L, Zha J, Spear PA, Wang Z. 2007. Toxicity of the aromatase inhibitor letrozole to Japanese medaka (*Oryzias latipes*) eggs, larvae and breeding adults. *Comp Biochem Physiol C Toxicol Pharmacol* 145(4): 533-541.
- Thornton JW. 2001. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by ligand exploitation and serial genome expansions. *Proceedings of the National Academy of Sciences of the United States of America* 98(10): 5671-5676.
- Tilley WD, Marcelli M, Wilson JD, McPhaul MJ. 1989. Characterization and expression of a cDNA encoding the human androgen receptor. *Proceedings of the National Academy of Sciences of the United States of America* 86(1): 327-331.
- Todorov M, Mombelli E, Ait-Aissa S, Mekenyan O. 2011. Androgen receptor binding affinity: a QSAR evaluation. *SAR and QSAR in environmental research* 22(3): 265-291.
- Trudeau VL, Spanswick D, Fraser EJ, Larivière K, Crump D, Chiu S, et al. 2000. The role of amino acid neurotransmitters in the regulation of pituitary gonadotropin release in fish. *Biochemistry and Cell Biology* 78: 241-259.
- Trudeau VL. 1997. Neuroendocrine regulation of gonadotropin II release and gonadal growth in the goldfish, *Carassius auratus*. *Reviews of Reproduction* 2: 55-68.
- Tyler C, Sumpter J. 1996. Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287-318.
- Tyler C, van der Eerden B, Jobling S, Panter G, Sumpter J. 1996. Measurement of vitellogenin, a biomarker for exposure to oestrogenic chemicals, in a wide variety of cyprinid fish. *Journal of Comparative Physiology and Biology* 166: 418-426.
- van der Burg B, Winter R, Man HY, Vangenechten C, Berckmans P, Weimer M, et al. 2010. Optimization and prevalidation of the in vitro AR CALUX method to test androgenic and antiandrogenic activity of compounds. *Reproductive toxicology* 30(1): 18-24.
- Villeneuve DL, Ankley GT, Makynen EA, Blake LS, Greene KJ, Higley EB, et al. 2007. Comparison of fathead minnow ovary explant and H295R cell-based steroidogenesis assays for identifying endocrine-active chemicals. *Ecotoxicol Environ Saf* 68(1): 20-32.
- Villeneuve DL, Breen M, Bencic DC, Cavallin JE, Jensen KM, Makynen EA, et al. 2013. Developing Predictive Approaches to Characterize Adaptive Responses of the Reproductive Endocrine Axis to Aromatase Inhibition: I. Data Generation in a Small Fish Model. *Toxicological sciences : an official journal of the Society of Toxicology*.

- Villeneuve DL, Mueller ND, Martinovic D, Makynen EA, Kahl MD, Jensen KM, et al. 2009. Direct effects, compensation, and recovery in female fathead minnows exposed to a model aromatase inhibitor. *Environ Health Perspect* 117(4): 624-631.
- Waller CL, Juma BW, Gray EJ, Kelce WR. 1996. Three-dimensional quantitative structure-activity relationships for androgen receptor ligands. *Toxicology and Applied Pharmacology* 137: 219-227.
- Wilson VS, Bobseine K, Lambright CR, Gray LE. 2002. A novel cell line, MDA-kb2, that stably expresses an androgen- and glucocorticoid-responsive reporter for the detection of hormone receptor agonists and antagonists. *Toxicological Sciences* 66: 69-81.
- Wilson VS, Cardon MC, Gray LE, Jr., Hartig PC. 2007. Competitive binding comparison of endocrine-disrupting compounds to recombinant androgen receptor from fathead minnow, rainbow trout, and human. *Environmental toxicology and chemistry / SETAC* 26(9): 1793-1802.
- Wolf JC, Dietrich DR, Friederich U, Caunter J, Brown AR. 2004. Qualitative and quantitative histomorphologic assessment of fathead minnow *Pimephales promelas* gonads as an endpoint for evaluating endocrine-active compounds: a pilot methodology study. *Toxicol Pathol* 32(5): 600-612.
- Yaron Z. 1995. Endocrine control of gametogenesis and spawning induction in the carp. *Aquaculture* 129: 49-73.
- Yin D, He Y, Perera MA, Hong SS, Marhefka C, Stourman N, et al. 2003. Key structural features of nonsteroidal ligands for binding and activation of the androgen receptor. *Molecular pharmacology* 63(1): 211-223.
- Young JM, McNeilly AS. 2010. Theca: the forgotten cell of the ovarian follicle. *Reproduction* 140(4): 489-504.
- Zhang X, Hecker M, Tompsett AR, Park JW, Jones PD, Newsted J, et al. 2008. Responses of the medaka HPG axis PCR array and reproduction to prochloraz and ketoconazole. *Environ Sci Technol* 42(17): 6762-6769.

Appendix 1: List of MIE(s), KE(s) and AO(s) in the AOP

List of MIE(s) in this AOP

Event: 25: Agonism, Androgen receptor

Short Name: Agonism, Androgen receptor

Key Event Component

Process	Object	Action
androgen receptor activity	androgen receptor	increased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	MolecularInitiatingEvent

Stressors

Name
17beta-Trenbolone
Spironolactone
5alpha-Dihydrotestosterone

Biological Context

Level of Biological Organization
Molecular

Evidence for Perturbation by Stressor

Overview for Molecular Initiating Event

Characterization of chemical properties: Androgen receptor binding chemicals can be grouped into two broad structural domains, steroidal and non-steroidal (Yin et al. 2003). Steroidal androgens consist primarily of testosterone and its derivatives (Yin et al. 2003). Many of the non-steroidal AR binding chemicals studied are derivatives of well known non-steroidal AR antagonists like bicalutamide, hydroxyflutamide, and nilutamide (Yin et al. 2003). Nonetheless, a number of QSARs and SARs that consider AR binding of both these pharmaceutical agents as well as environmental chemicals have been developed

(Waller et al. 1996; Serafimova et al. 2002; Todorov et al. 2011; Hong et al. 2003; Bohl et al. 2004). However, it has been shown that very minor structural differences can dramatically impact function as either an agonist or antagonist (Yin et al. 2003; Bohl et al. 2004; Norris et al. 2009), making it difficult at present to predict agonist versus antagonist activity based on chemical structure alone.

In vivo considerations: A variety of steroidal androgens can be converted to estrogens in vitro through the action of cytochrome P450 19 (aromatase). Structures subject to aromatization may behave in vivo as estrogens despite exhibiting potent androgen receptor agonism in vitro.

5alpha-Dihydrotestosterone

Chemical is a non-aromatizable androgen.

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	High	NCBI
medaka	Oryzias latipes	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

Taxonomic applicability: Androgen receptor orthologs are primarily limited to vertebrates (Baker 1997; Thornton 2001; Eick and Thornton 2011; Markov and Laudet 2011). Therefore, this MIE would generally be viewed as relevant to vertebrates, but not invertebrates.

Key Event Description

Site of action: The molecular site of action is the ligand binding domain of the AR. This particular key event specifically refers to interaction with nuclear AR. Downstream KE responses to activation of membrane ARs may be different. The cellular site of action for the molecular initiating event is undefined.

Responses at the macromolecular level: Binding of a ligand, including xenobiotics that act as AR agonists, to the cytosolic AR mediates a conformational shift that facilitates dissociation from accompanying heat shock proteins and dimerization with another AR (Prescott and Coetzee 2006; Claessens et al. 2008; Centenera et al. 2008). Homodimerization unveils a nuclear localisation sequence, allowing the AR-ligand complex to translocate to the nucleus and bind to androgen-response elements (AREs) (Claessens et al. 2008; Cutress et al. 2008). This elicits recruitment of additional transcription factors and transcriptional activation of androgen-responsive genes (Heemers and Tindall 2007).

AR paralogs:

- Most vertebrates have a single gene coding for nuclear AR. However, most fish have two AR genes (AR-A, AR-B) as a result of a whole genome duplication event after the split of Acipenseriformes from teleosts but before the divergence of Osteoglossiformes (Douard et al. 2008).
- AR-B has been lost in Cypriniformes, Siluriformes, Characiformes, and Salmoniformes (Douard et al. 2008).
- In Percomorphs, AR-B has accumulated significant substitutions in the both ligand binding and DNA binding domains (Douard et al. 2008).
- Differential ligand selectivity and subcellular localisation has been reported for AR paralogs in some fish species (e.g., Bain et al. 2015), but the difference is not easily generalized based on available data in the literature.

How it is Measured or Detected

Measurement/detection:

- **In vitro methods:**
 - OECD Test No. 458: Stably transfected human androgen receptor transcriptional activation assay for detection of androgen agonists and antagonists has been reviewed and validated by OECD and is well suited for detection of this key event ([OECD 2016](#)).
 - Binding to the androgen receptor can be directly measured in cell free systems based on displacement of a radio-labeled standard (generally testosterone or DHT) in a competitive binding assay (e.g., (Olsson et al. 2005; Sperry and Thomas 1999; Wilson et al. 2007; Tilley et al. 1989; Kim et al. 2010).
 - Cell based transcriptional activation assays are typically required to differentiate agonists from antagonists, in vitro. A number of reporter gene assays have been developed and used to screen chemicals for AR agonist and/or antagonist activity (e.g., Wilson et al. 2002; van der Burg et al. 2010; Mak et al. 1999; Araki et al. 2005).
 - Expression of androgen responsive proteins like spiggin in primary cell cultures has also been used to detect AR agonist activity (Jolly et al. 2006).
- **In vivo methods:**
 - In fish, phenotypic masculinisation of females has frequently been used as an indirect measurement of in vivo androgen receptor agonism.

- Development of nuptial tubercles, a dorsal fatpad, and a characteristic banding pattern has been observed in female fathead minnows exposed to androgen agonists (Ankley et al. 2003; Jensen et al. 2006; Ankley et al. 2010; LaLone et al. 2013; [OECD 2012](#)).
- Anal fin elongation in female western mosquitofish (*Gambusia affinis*) has similarly been viewed as evidence of AR activation (Raut et al. 2011; Sone et al. 2005).
- In medaka, development of papillary processes, which normally only appear on the second to seventh or eighth fin ray of the anal fin, has also been used as an indirect measure of androgen receptor agonism ([OECD 2012](#)).
- Production of the nest building glue, spiggin, in three female 3-spined sticklebacks (*Gasterosteus aculeatus*) has also been well documented as an indicator of androgen receptor agonism (Jakobsson et al. 1999; Hahlbeck et al. 2004). Quantification of the spiggin protein in exposed female 3-spined stickleback or green fluorescence protein expression in a transgenic spg1-gfp medaka line (Sébillot et al. 2014) can be used to detect androgen receptor agonism.
- **High Throughput Screening**
 - Measures of AR agonism have been included in high throughput screening programs, such as US EPA's Toxcast program. Toxcast assays relevant for screening chemicals for their ability to bind and/or activate the AR include:
 - ATG_AR_TRANS A cell based assay that can differentiate agonism from antagonism
 - NVS_NR_hAR A cell free assay using recombinant human AR. Can detect binding, but cannot distinguish agonism from antagonism.
 - NVS_NR_rAR A cell free assay using recombinant rat AR. Can detect binding, but cannot distinguish agonism from antagonism.
 - OT_AR_ARELUC_AG_1440 A cell based assay that measures expression of a reporter gene under control of androgen-responsive elements. Can distinguish agonism from antagonism.
 - Tox21_AR_BLA_Agonist_ratio A cell based assay with an inducible reporter. Can distinguish agonists from antagonists.
 - Tox21_AR_LUC_MDAKB2_agonist A cell based assay with an inducible reporter. Can distinguish agonists from antagonists.
 - [Assay descriptions](#)

References

- Ankley GT, Gray LE. Cross-species conservation of endocrine pathways: a critical analysis of tier 1 fish and rat screening assays with 12 model chemicals. Environ Toxicol Chem. 2013 Apr;32(5):1084-7. doi: 10.1002/etc.2151. Epub 2013 Mar 19. PubMed PMID: 23401061.

- Ankley GT, Jensen KM, Kahl MD, Durhan EJ, Makynen EA, Cavallin JE, Martinović D, Wehmas LC, Mueller ND, Villeneuve DL. Use of chemical mixtures to differentiate mechanisms of endocrine action in a small fish model. *Aquat Toxicol*. 2010 Sep 1;99(3):389-96. doi: 10.1016/j.aquatox.2010.05.020. Epub 2010 Jun 4. PubMed PMID: 20573408.
- Ankley GT, Jensen KM, Makynen EA, Kahl MD, Korte JJ, Hornung MW, Henry TR, Denny JS, Leino RL, Wilson VS, Cardon MC, Hartig PC, Gray LE. Effects of the androgenic growth promoter 17-beta-trenbolone on fecundity and reproductive endocrinology of the fathead minnow. *Environ Toxicol Chem*. 2003 Jun;22(6):1350-60. PubMed PMID: 12785594.
- Araki N, Ohno K, Nakai M, Takeyoshi M, Iida M. 2005. Screening for androgen receptor activities in 253 industrial chemicals by in vitro reporter gene assays using AR-EcoScreen cells. *Toxicology in vitro : an international journal published in association with BIBRA* 19(6): 831-842.
- Bain PA, Ogino Y, Miyagawa S, Iguchi T, Kumar A. Differential ligand selectivity of androgen receptors α and β from Murray-Darling rainbowfish (*Melanotaenia fluviatilis*). *Gen Comp Endocrinol*. 2015 Feb 1;212:84-91. doi: 10.1016/j.ygcen.2015.01.024. PubMed PMID: 25644213.
- Baker ME. 1997. Steroid receptor phylogeny and vertebrate origins. *Molecular and cellular endocrinology* 135(2): 101-107.
- Bohl CE, Chang C, Mohler ML, Chen J, Miller DD, Swaan PW, et al. 2004. A ligand-based approach to identify quantitative structure-activity relationships for the androgen receptor. *Journal of medicinal chemistry* 47(15): 3765-3776.
- Centenera MM, Harris JM, Tilley WD, Butler LM. 2008. The contribution of different androgen receptor domains to receptor dimerization and signaling. *Molecular endocrinology* 22(11): 2373-2382.
- Claessens F, Denayer S, Van Tilborgh N, Kerkhofs S, Helsen C, Haelens A. 2008. Diverse roles of androgen receptor (AR) domains in AR-mediated signaling. *Nuclear receptor signaling* 6: e008.
- Cutress ML, Whitaker HC, Mills IG, Stewart M, Neal DE. 2008. Structural basis for the nuclear import of the human androgen receptor. *Journal of cell science* 121(Pt 7): 957-968.
- Douard V, Brunet F, Boussau B, Ahrens-Fath I, Vlaeminck-Guillem V, Haendler B, Laudet V, Guiguen Y. The fate of the duplicated androgen receptor in fishes: a late neofunctionalization event? *BMC Evol Biol*. 2008 Dec 18;8:336. doi: 10.1186/1471-2148-8-336. PubMed PMID: 19094205
- Eick GN, Thornton JW. 2011. Evolution of steroid receptors from an estrogen-sensitive ancestral receptor. *Molecular and cellular endocrinology* 334(1-2): 31-38.
- Hahlbeck E, Katsiadaki I, Mayer I, Adolfsson-Erici M, James J, Bengtsson BE. The juvenile three-spined stickleback (*Gasterosteus aculeatus* L.) as a model organism for endocrine disruption II--kidney hypertrophy, vitellogenin and spiggin induction. *Aquat Toxicol*. 2004 Dec 20;70(4):311-26
- Hong H, Fang H, Xie Q, Perkins R, Sheehan DM, Tong W. 2003. Comparative molecular field analysis (CoMFA) model using a large diverse set of natural, synthetic and environmental chemicals for binding to the androgen receptor. *SAR and QSAR in environmental research* 14(5-6): 373-388.
- Jakobsson, S., Borg, B., Haux, C. et al. *Fish Physiology and Biochemistry* (1999) 20: 79. doi:10.1023/A:1007776016610

- Jensen KM, Makynen EA, Kahl MD, Ankley GT. Effects of the feedlot contaminant 17alpha-trenbolone on reproductive endocrinology of the fathead minnow. *Environ Sci Technol*. 2006 May 1;40(9):3112-7. PubMed PMID: 16719119.
- Jolly C, Katsiadaki I, Le Belle N, Mayer I, Dufour S. 2006. Development of a stickleback kidney cell culture assay for the screening of androgenic and anti-androgenic endocrine disrupters. *Aquatic toxicology* 79(2): 158-166.
- Kim TS, Yoon CY, Jung KK, Kim SS, Kang IH, Baek JH, et al. 2010. In vitro study of Organization for Economic Co-operation and Development (OECD) endocrine disruptor screening and testing methods- establishment of a recombinant rat androgen receptor (rrAR) binding assay. *The Journal of toxicological sciences* 35(2): 239-243.
- LaLone CA, Villeneuve DL, Cavallin JE, Kahl MD, Durhan EJ, Makynen EA, Jensen KM, Stevens KE, Severson MN, Blanksma CA, Flynn KM, Hartig PC, Woodard JS, Berninger JP, Norberg-King TJ, Johnson RD, Ankley GT. Cross-species sensitivity to a novel androgen receptor agonist of potential environmental concern, spironolactone. *Environ Toxicol Chem*. 2013 Nov;32(11):2528-41. doi: 10.1002/etc.2330. Epub 2013 Sep 6. PubMed PMID: 23881739.
- Mak P, Cruz FD, Chen S. 1999. A yeast screen system for aromatase inhibitors and ligands for androgen receptor: yeast cells transformed with aromatase and androgen receptor. *Environmental health perspectives* 107(11): 855-860.
- Markov GV, Laudet V. 2011. Origin and evolution of the ligand-binding ability of nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 21-30.
- Norris JD, Joseph JD, Sherk AB, Juzumiene D, Turnbull PS, Rafferty SW, et al. 2009. Differential presentation of protein interaction surfaces on the androgen receptor defines the pharmacological actions of bound ligands. *Chemistry & biology* 16(4): 452-460.
- OECD (2012), *Test No. 229: Fish Short Term Reproduction Assay*, OECD Publishing, Paris. DOI: <http://dx.doi.org/10.1787/9789264185265-en>
- OECD (2016), *Test No. 458: Stably Transfected Human Androgen Receptor Transcriptional Activation Assay for Detection of Androgenic Agonist and Antagonist Activity of Chemicals*, OECD Publishing, Paris. DOI: <http://dx.doi.org/10.1787/9789264264366-en>
- Olsson P-E, Berg A, von Hofsten J, Grahm B, Hellqvist A, Larsson A, et al. 2005. Molecular cloning and characterization of a nuclear androgen receptor activated by 11-ketotestosterone. *Reproductive Biology and Endocrinology* 3: 1-17.
- Prescott J, Coetzee GA. 2006. Molecular chaperones throughout the life cycle of the androgen receptor. *Cancer letters* 231(1): 12-19.
- Serafimova R, Walker J, Mekenyan O. 2002. Androgen receptor binding affinity of pesticide "active" formulation ingredients. QSAR evaluation by COREPA method. *SAR and QSAR in environmental research* 13(1): 127-134.
- Sone K, Hinago M, Itamoto M, Katsu Y, Watanabe H, Urushitani H, Tooi O, Guillelte LJ Jr, Iguchi T. Effects of an androgenic growth promoter 17beta-trenbolone on masculinization of Mosquitofish (*Gambusia affinis affinis*). *Gen Comp Endocrinol*. 2005 Sep 1;143(2):151-60. Epub 2005 Apr 13. PubMed PMID: 16061073.
- Sperry TS, Thomas P. 1999. Identification of two nuclear androgen receptors in kelp bass (*Paralabrax clathratus*) and their binding affinities for xenobiotics:

comparison with Atlantic croaker (*Micropogonias undulatus*) androgen receptors. *Biology of reproduction* 61(4): 1152-1161.

- Stanko JP, Angus RA. In vivo assessment of the capacity of androstenedione to masculinize female mosquitofish (*Gambusia affinis*) exposed through dietary and static renewal methods. *Environ Toxicol Chem.* 2007 May;26(5):920-6. PubMed PMID: 17521138.
- Thornton JW. 2001. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by ligand exploitation and serial genome expansions. *Proceedings of the National Academy of Sciences of the United States of America* 98(10): 5671-5676.
- Tilley WD, Marcelli M, Wilson JD, McPhaul MJ. 1989. Characterization and expression of a cDNA encoding the human androgen receptor. *Proceedings of the National Academy of Sciences of the United States of America* 86(1): 327-331.
- Todorov M, Mombelli E, Ait-Aissa S, Mekenyan O. 2011. Androgen receptor binding affinity: a QSAR evaluation. *SAR and QSAR in environmental research* 22(3): 265-291.
- van der Burg B, Winter R, Man HY, Vangenechten C, Berckmans P, Weimer M, et al. 2010. Optimization and prevalidation of the in vitro AR CALUX method to test androgenic and antiandrogenic activity of compounds. *Reproductive toxicology* 30(1): 18-24.
- Waller CL, Juma BW, Gray EJ, Kelce WR. 1996. Three-dimensional quantitative structure-activity relationships for androgen receptor ligands. *Toxicology and Applied Pharmacology* 137: 219-227.
- Wilson VS, Bobseine K, Lambright CR, Gray LE. 2002. A novel cell line, MDA-kb2, that stably expresses an androgen- and glucocorticoid-responsive reporter for the detection of hormone receptor agonists and antagonists. *Toxicological Sciences* 66: 69-81.
- Wilson VS, Cardon MC, Gray LE, Jr., Hartig PC. 2007. Competitive binding comparison of endocrine-disrupting compounds to recombinant androgen receptor from fathead minnow, rainbow trout, and human. *Environmental toxicology and chemistry / SETAC* 26(9): 1793-1802.
- Yin D, He Y, Perera MA, Hong SS, Marhefka C, Stourman N, et al. 2003. Key structural features of nonsteroidal ligands for binding and activation of the androgen receptor. *Molecular pharmacology* 63(1): 211-223.

List of KE(s) in the AOP

Event: 129: Reduction, Gonadotropins, circulating concentrations

Short Name: Reduction, Gonadotropins, circulating concentrations

Key Event Component

Process	Object	Action
	Luteinizing hormone	decreased
	Follicle stimulating hormone	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	KeyEvent

Stressors

Name
n/a

Biological Context

Level of Biological Organization
Organ

Organ term

Organ term
blood plasma

Domain of Applicability

Life Stage Applicability

Life Stage	Evidence
Not Otherwise Specified	Not Specified

Sex Applicability

Sex	Evidence
Unspecific	Not Specified

A functional hypothalamic-pituitary-gonadal axis involving GnRH and gonadotropin-mediated regulation of reproductive functions is a vertebrate trait (Sower et al. 2009). The taxonomic applicability of this key event is limited to chordates.

Key Event Description

Gonadotropin (luteinizing hormone [LH] and follicle-stimulating hormone [FSH]) secretion from the pituitary is a key regulator of gonadal steroid biosynthesis. LH and FSH are heterodimeric glycoproteins composed of a hormone-specific beta subunit and a common alpha subunit (Norris 2007). The subunits are synthesised in pituitary gonadotropes and stored in secretory vesicles. Gonadotropin secretion by pituitary gonadotropes is regulated via gonadotropin releasing hormone (GnRH) signaling from the hypothalamus as well as by intrapituitary regulators of gonadotropin expression (e.g., activin, follistatin, inhibin) (Norris 2007; Habibi and Huggard 1998).

How it is Measured or Detected

- Circulating concentrations of gonadotropins in humans and common mammalian models (e.g., rodents, many livestock species) can be directly measured using either commercial or custom immunoassays (e.g., enzyme-linked immunosorbent assays, radioimmunoassays, etc.).
- Similar immunoassay-based methods have been developed for quantifying gonadotropins in fish (e.g., Govoroun et al. 1998; Amano et al. 2000; Kah et al. 1989; Prat et al. 1996). However, at present, antibodies specific for distinguishing LH and FSH are only available for a limited number of species, primarily salmonids (Levavi-Sivan et al. 2010).
- Expression of mRNAs coding for luteinizing hormone beta subunit (lhb) and follicle-stimulating hormone beta subunit (fshb) tend to fluctuate in parallel in repeat-spawning fish and plasma concentrations LH and FSH in tilapia were also shown to fluctuate in parallel (reviewed in Levavi-Sivan et al. 2010). Consequently, the two gonadotropins are treated non-specifically for the purposes of the current key event.
- For small fish species limited plasma volumes relative to the sensitivity of the available immunoassay methods may impose limits on the ability to measure this key event directly and reliably.

References

- Amano M, Iigo M, Ikuta K, Kitamura S, Yamada H, Yamamori K. 2000. Roles of melatonin in gonadal maturation of underyearling precocious male masu salmon. *General and comparative endocrinology* 120(2): 190-197.

- Cheng GF, Yuen CW, Ge W. 2007. Evidence for the existence of a local activin follistatin negative feedback loop in the goldfish pituitary and its regulation by activin and gonadal steroids. *The Journal of endocrinology* 195(3): 373-384.
- Govoroun M, Chyb J, Breton B. 1998. Immunological cross-reactivity between rainbow trout GTH I and GTH II and their alpha and beta subunits: application to the development of specific radioimmunoassays. *General and comparative endocrinology* 111(1): 28-37.
- Habibi HR, Huggard DL. 1998. Testosterone regulation of gonadotropin production in goldfish. *Comparative biochemistry and physiology Part C, Pharmacology, toxicology & endocrinology* 119(3): 339-344.
- Kah O, Pontet A, Nunez Rodriguez J, Calas A, Breton B. 1989. Development of an enzyme-linked immunosorbent assay for goldfish gonadotropin. *Biology of reproduction* 41(1): 68-73.
- Levavi-Sivan B, Bogerd J, Mananos EL, Gomez A, Lareyre JJ. 2010. Perspectives on fish gonadotropins and their receptors. *General and comparative endocrinology* 165(3): 412-437.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Oakley AE, Clifton DK, Steiner RA. 2009. Kisspeptin signaling in the brain. *Endocrine reviews* 30(6): 713-743.
- Prat F, Sumpter JP, Tyler CR. 1996. Validation of radioimmunoassays for two salmon gonadotropins (GTH I and GTH II) and their plasma concentrations throughout the reproductive cycle in male and female rainbow trout (*Oncorhynchus mykiss*). *Biology of reproduction* 54(6): 1375-1382.
- Sower SA, Freamat M, Kavanaugh SI. 2009. The origins of the vertebrate hypothalamic-pituitary-gonadal (HPG) and hypothalamic-pituitary-thyroid (HPT) endocrine systems: new insights from lampreys. *General and comparative endocrinology* 161(1): 20-29.
- Trudeau VL, Spanswick D, Fraser EJ, Larivière K, Crump D, Chiu S, et al. 2000. The role of amino acid neurotransmitters in the regulation of pituitary gonadotropin release in fish. *Biochemistry and Cell Biology* 78: 241-259.
- Trudeau VL. 1997. Neuroendocrine regulation of gonadotropin II release and gonadal growth in the goldfish, *Carassius auratus*. *Reviews of Reproduction* 2: 55-68.

Event: 274: Reduction, Testosterone synthesis by ovarian theca cells

Short Name: Reduction, Testosterone synthesis by ovarian theca cells

Key Event Component

Process	Object	Action
testosterone biosynthetic process	testosterone	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	KeyEvent

Biological Context

Level of Biological Organization
Cellular

Cell term

Cell term
theca cell

Domain of Applicability**Taxonomic Applicability**

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	High	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	Not Specified

This key event is relevant to vertebrates and amphioxus, but not invertebrates.

- Cytochrome P45011a (Cyp11a), a rate limiting enzyme for the production of testosterone, is specific to vertebrates and amphioxus (Markov et al. 2009; Baker et al. 2011; Payne and Hales, 2004).
- Cyp11a does not occur in invertebrates, as a result, they do not synthesise testosterone, nor other steroid intermediates required for testosterone synthesis (Markov et al. 2009; Payne and Hales, 2004).

Key Event Description

Testosterone is synthesised in ovarian theca cells through a series of enzyme catalyzed reactions that convert cholesterol to androgens (see KEGG reference pathway 00140 for details; www.genome.jp/kegg; (Payne and Hales 2004; Magoffin 2005; Young and McNeilly 2010). Binding of luteinizing hormone to luteinizing hormone receptors located on the surface of theca cell membranes leads to increased expression of steroidogenic cytochrome P450s, steroidogenic acute regulatory protein, and consequent increases in androgen production (Payne and Hales 2004; Miller 1988; Miller and Strauss 1999).

How it is Measured or Detected

Steroid production by isolated primary theca cells can be measured using radioimmunoassay or enzyme linked immunosorbent assay approaches (e.g., (Benninghoff and Thomas 2006; Campbell et al. 1998). However, the isolation and culture methods are not trivial. Similarly, development of immortalized theca cell lines has proven challenging (Havelock et al. 2004). Consequently, this key event is perhaps best evaluated by examining T production by intact ovarian tissue explants either exposed to chemicals in vitro (e.g., Villeneuve et al. 2007; McMaster ME 1995) or in vivo (i.e., via ex vivo steroidogenesis assay; e.g., Ankley et al. 2007). Reductions in T production by ovarian tissue explants can indicate either direct inhibition of steroidogenic enzymes involved in T synthesis, or indirect impacts due to feedback along the hypothalamic-pituitary-gonadal axis (in cases where chemical exposures occur in vivo). However, because T synthesis in the theca cells is closely linked to estradiol (E2) synthesis by granulosa cells, reductions in T production by intact ovary tissue can also be due to increased aromatase activity and the resulting increased rate of converting T to E2.

References

- Ankley GT, Jensen KM, Kahl MD, Makynen EA, Blake LS, Greene KJ, et al. 2007. Ketoconazole in the fathead minnow (*Pimephales promelas*): reproductive toxicity and biological compensation. *Environ Toxicol Chem* 26(6): 1214-1223.
- Baker ME. 2011. Origin and diversification of steroids: Co-evolution of enzymes and nuclear receptors. *Mol Cell Endocrinol* 334: 14-20.

- Benninghoff AD, Thomas P. 2006. Gonadotropin regulation of testosterone production by primary cultured theca and granulosa cells of Atlantic croaker: I. Novel role of CaMKs and interactions between calcium- and adenylyl cyclase-dependent pathways. *General and comparative endocrinology* 147(3): 276-287.
- Campbell BK, Baird DT, Webb R. 1998. Effects of dose of LH on androgen production and luteinization of ovine theca cells cultured in a serum-free system. *Journal of reproduction and fertility* 112(1): 69-77.
- Havelock JC, Rainey WE, Carr BR. 2004. Ovarian granulosa cell lines. *Molecular and cellular endocrinology* 228(1-2): 67-78.
- Magoffin DA. 2005. Ovarian theca cell. *The international journal of biochemistry & cell biology* 37(7): 1344-1349.
- Markov GV, Tavares R, Dauphin-Villemant C, Demeneix BA, Baker ME, Laudet V. Independent elaboration of steroid hormone signaling pathways in metazoans. *Proc Natl Acad Sci U S A*. 2009 Jul 21;106(29):11913-8. doi: 10.1073/pnas.0812138106.
- McMaster ME MK, Jardine JJ, Robinson RD, Van Der Kraak GJ. 1995. Protocol for measuring in vitro steroid production by fish gonadal tissue. *Canadian Technical Report of Fisheries and Aquatic Sciences* 1961 1961: 1-78.
- Miller WL, Strauss JF, 3rd. 1999. Molecular pathology and mechanism of action of the steroidogenic acute regulatory protein, StAR. *The Journal of steroid biochemistry and molecular biology* 69(1-6): 131-141.
- Miller WL. 1988. Molecular biology of steroid hormone synthesis. *Endocrine reviews* 9(3): 295-318.
- Payne AH, Hales DB. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine reviews* 25(6): 947-970
- Villeneuve DL, Ankley GT, Makynen EA, Blake LS, Greene KJ, Higley EB, et al. 2007. Comparison of fathead minnow ovary explant and H295R cell-based steroidogenesis assays for identifying endocrine-active chemicals. *Ecotoxicol Environ Saf* 68(1): 20-32.
- Young JM, McNeilly AS. 2010. Theca: the forgotten cell of the ovarian follicle. *Reproduction* 140(4): 489-504.

Event: 3: Reduction, 17beta-estradiol synthesis by ovarian granulosa cells

Short Name: Reduction, 17beta-estradiol synthesis by ovarian granulosa cells

Key Event Component

Process	Object	Action
estrogen biosynthetic process	17beta-estradiol	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:7 - Aromatase (Cyp19a1) reduction leading to impaired fertility in adult female	KeyEvent
Aop:25 - Aromatase inhibition leading to reproductive dysfunction	KeyEvent
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	KeyEvent
Aop:122 - Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	KeyEvent
Aop:123 - Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	KeyEvent

Biological Context

Level of Biological Organization
Cellular

Cell term

Cell term
granulosa cell

Domain of Applicability**Taxonomic Applicability**

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	High	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

Key enzymes needed to synthesise 17 β -estradiol first appear in the common ancestor of amphioxus and vertebrates (Markov et al. 2009; Baker 2011). Consequently, it is plausible that this key event is applicable to most vertebrates. This key event is not applicable to invertebrates, which lack the enzymes required to synthesise 17 β -estradiol.

Key Event Description

Like all steroids, estradiol is a cholesterol derivative. Estradiol synthesis in ovary is mediated by a number of enzyme catalyzed reactions involving cyp11 (cholesterol side chain cleavage enzyme), cyp 17 (17 α -hydroxylase/17,20-lyase), 3 β hydroxysteroid dehydrogenase, 17 β hydroxysteroid dehydrogenase, and cyp19 (aromatase). Among those enzyme catalyzed reactions, conversion of testosterone to estradiol, catalyzed by aromatase, is considered to be rate limiting for estradiol synthesis. Within the ovary, aromatase expression and activity is primarily localised in the granulosa cells (reviewed in (Norris 2007; Yaron 1995; Havelock et al. 2004) and others). Reactions involved in synthesis of C-19 androgens are primarily localised in the theca cells and C-19 androgens diffuse from the theca into granulosa cells where aromatase can catalyze their conversion to C-18 estrogens.

How it is Measured or Detected

Due to the importance of both theca and granulosa cells in ovarian steroidogenesis, it is generally impractical to measure E2 production by isolated granulosa cells (Havelock et al. 2004). However, this key event can be evaluated by examining E2 production by intact ovarian tissue explants either exposed to chemicals in vitro (e.g., Villeneuve et al. 2007; McMaster ME 1995) or in vivo (i.e., via ex vivo steroidogenesis assay; e.g., Ankley et al. 2007). Estradiol released by ovarian tissue explants into media can be quantified by radioimmunoassay (e.g., Jensen et al. 2001), ELISA, or analytical methods such as LC-MS (e.g., Owen et al. 2014).

OECD TG 456 ([OECD 2011](#)) is the validated test guideline for an in vitro screen for chemical effects on steroidogenesis, specifically the production of 17 β -estradiol (E2) and testosterone (T).

The synthesis of E2 can be measured in vitro cultured ovarian cells. The methods for culturing mammalian ovarian cells can be found in the Database Service on Alternative Methods to animal experimentation (DB-ALM): Culture of Human Cumulus Granulosa

Cells ([EURL ECVAM Protocol No. 92](#)), Granulosa and Theca Cell Culture Systems ([EURL ECVAM Method Summary No. 92](#)).

References

- Ankley GT, Jensen KM, Kahl MD, Makynen EA, Blake LS, Greene KJ, et al. 2007. Ketoconazole in the fathead minnow (*Pimephales promelas*): reproductive toxicity and biological compensation. *Environ Toxicol Chem* 26(6): 1214-1223.
- Baker ME. 2011. Origin and diversification of steroids: co-evolution of enzymes and nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 14-20.
- EURL ECVAM Method Summary no 92. Granulosa and Theca Cell Culture Systems - Summary
- EURL ECVAM Protocol no 92 Culture of Human Cumulus Granulosa Cells. Primary cell culture method. Contact Person: Dr. Mahadevan Maha M.
- Havelock JC, Rainey WE, Carr BR. 2004. Ovarian granulosa cell lines. *Molecular and cellular endocrinology* 228(1-2): 67-78.
- Jensen K, Korte J, Kahl M, Pasha M, Ankley G. 2001. Aspects of basic reproductive biology and endocrinology in the fathead minnow (*Pimephales promelas*). *Comparative Biochemistry and Physiology Part C* 128: 127-141.
- McMaster ME MK, Jardine JJ, Robinson RD, Van Der Kraak GJ. 1995. Protocol for measuring in vitro steroid production by fish gonadal tissue. *Canadian Technical Report of Fisheries and Aquatic Sciences* 1961 1961: 1-78.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- OECD (2011), Test No. 456: H295R Steroidogenesis Assay, OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris. DOI: <http://dx.doi.org/10.1787/9789264122642-en>
- Owen LJ, Wu FC, Keevil BG. 2014. A rapid direct assay for the routine measurement of oestradiol and oestrone by liquid chromatography tandem mass spectrometry. *Ann. Clin. Biochem.* 51(pt 3):360-367.
- Villeneuve DL, Ankley GT, Makynen EA, Blake LS, Greene KJ, Higley EB, et al. 2007. Comparison of fathead minnow ovary explant and H295R cell-based steroidogenesis assays for identifying endocrine-active chemicals. *Ecotoxicol Environ Saf* 68(1): 20-32.
- Villeneuve DL, Mueller ND, Martinovic D, Makynen EA, Kahl MD, Jensen KM, et al. 2009. Direct effects, compensation, and recovery in female fathead minnows exposed to a model aromatase inhibitor. *Environ Health Perspect* 117(4): 624-631.
- Yaron Z. 1995. Endocrine control of gametogenesis and spawning induction in the carp. *Aquaculture* 129: 49-73.

Event: 219: Reduction, Plasma 17beta-estradiol concentrations

Short Name: Reduction, Plasma 17beta-estradiol concentrations

Key Event Component

Process	Object	Action
	17beta-estradiol	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:7 - Aromatase (Cyp19a1) reduction leading to impaired fertility in adult female	KeyEvent
Aop:25 - Aromatase inhibition leading to reproductive dysfunction	KeyEvent
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	KeyEvent
Aop:122 - Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	KeyEvent
Aop:123 - Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	KeyEvent

Biological Context

Level of Biological Organization
Organ

Organ term

Organ term
blood plasma

Domain of Applicability**Taxonomic Applicability**

Term	Scientific Term	Evidence	Links
rat	Rattus norvegicus	High	NCBI
human	Homo sapiens	High	NCBI
fathead minnow	Pimephales promelas	High	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult	High

Sex Applicability

Sex	Evidence
Unspecific	Not Specified

Key enzymes needed to synthesise 17β -estradiol first appear in the common ancestor of amphioxus and vertebrates (Baker 2011). Consequently, this key event is applicable to most vertebrates.

Key Event Description

Estradiol synthesised by the gonads is transported to other tissues via blood circulation. The gonads are generally considered to be the primary source of estrogens in systemic circulation.

How it is Measured or Detected

Total concentrations of 17β -estradiol in plasma can be measured by radioimmunoassay (e.g., Jensen et al. 2001), enzyme-linked immunosorbent assay (available through many commercial vendors), or by analytical chemistry (e.g., LC/MS; Owen et al. 2014). Total steroid hormones are typically extracted from plasma or serum via liquid-liquid or solid phase extraction prior to analysis.

Given that there are numerous genes, like those coding for vertebrate vitellogenins, choriongenins, cyp19a1b, etc. which are known to be regulated by estrogen response elements, targeted qPCR or proteomic analysis of appropriate targets could also be used as an indirect measure of reduced circulating estrogen concentrations. However, further support for the specificity of the individual gene targets for estrogen-dependent regulation should be established in order to support their use.

A line of transgenic zebrafish employing green fluorescence protein under control of estrogen response elements could also be used to provide direct evidence of altered estrogen, with decreased GFP signal in estrogen responsive tissues like liver, ovary, pituitary, and brain indicating a reduction in circulating estrogens (Gorelick and Halpern 2011).

References

- Jensen K, Korte J, Kahl M, Pasha M, Ankley G. 2001. Aspects of basic reproductive biology and endocrinology in the fathead minnow (*Pimephales promelas*). Comparative Biochemistry and Physiology Part C 128: 127-141.

-
- Baker ME. 2011. Origin and diversification of steroids: co-evolution of enzymes and nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 14-20.
 - Owen LJ, Wu FC, Keevil BG. 2014. A rapid direct assay for the routine measurement of oestradiol and oestrone by liquid chromatography tandem mass spectrometry. *Ann. Clin. Biochem.* 51(pt 3):360-367.
 - Gorelick DA, Halpern ME. Visualization of estrogen receptor transcriptional activation in zebrafish. *Endocrinology*. 2011 Jul;152(7):2690-703. doi: 10.1210/en.2010-1257. Epub 2011 May 3. PubMed PMID: 21540282.

Event: 285: Reduction, Vitellogenin synthesis in liver

Short Name: Reduction, Vitellogenin synthesis in liver

Key Event Component

Process	Object	Action
gene expression	vitellogenins	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:25 - Aromatase inhibition leading to reproductive dysfunction	KeyEvent
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	KeyEvent
Aop:30 - Estrogen receptor antagonism leading to reproductive dysfunction	KeyEvent
Aop:122 - Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	KeyEvent
Aop:123 - Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	KeyEvent

Biological Context

Level of Biological Organization
Tissue

Cell term

Cell term
hepatocyte

Organ term

Organ term
liver

Domain of Applicability**Taxonomic Applicability**

Term	Scientific Term	Evidence	Links
Pimephales promelas	Pimephales promelas	High	NCBI

Fundulus heteroclitus	Fundulus heteroclitus	High	NCBI
Oryzias latipes	Oryzias latipes	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Unspecific	Not Specified

Oviparous vertebrates. Although vitellogenin is conserved among oviparous vertebrates and many invertebrates, liver is not a relevant tissue for the production of vitellogenin in invertebrates (Wahli 1988)

Key Event Description

Vitellogenin is an egg yolk precursor protein synthesised by hepatocytes of oviparous vertebrates. In vertebrates, transcription of vitellogenin genes is predominantly regulated by estrogens via their action on nuclear estrogen receptors. During vitellogenic periods of the reproductive cycle, when circulating estrogen concentrations are high, vitellogenin transcription and synthesis are typically orders of magnitude greater than during non-reproductive conditions.

How it is Measured or Detected

Relative abundance of vitellogenin transcripts or protein can be readily measured in liver tissue from organisms exposed in vivo (e.g., Biales et al. 2007), or in liver slices (e.g., Schmieder et al. 2000) or hepatocytes (e.g., Navas and Segner 2006) exposed in vitro, using real-time quantitative polymerase chain reaction (PCR; transcripts) or enzyme linked immunosorbent assay (ELISA; protein).

References

- Biales AD, Bencic DC, Lazorchak JL, Lattier DL. 2007. A quantitative real-time polymerase chain reaction method for the analysis of vitellogenin transcripts in model and nonmodel fish species. *Environ Toxicol Chem* 26(12): 2679-2686.
- Navas JM, Segner H. 2006. Vitellogenin synthesis in primary cultures of fish liver cells as endpoint for in vitro screening of the (anti)estrogenic activity of chemical substances. *Aquat Toxicol* 80(1): 1-22.
- Schmieder P, Tapper M, Linnun A, Denny J, Kolanczyk R, Johnson R. 2000. Optimization of a precision-cut trout liver tissue slice assay as a screen for

vitellogenin induction: comparison of slice incubation techniques. *Aquat Toxicol* 49(4): 251-268.

- Wahli W. 1988. Evolution and expression of vitellogenin genes. *Trends in Genetics*. 4:227-232.

Event: 221: Reduction, Plasma vitellogenin concentrations

Short Name: Reduction, Plasma vitellogenin concentrations

Key Event Component

Process	Object	Action
	vitellogenins	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:25 - Aromatase inhibition leading to reproductive dysfunction	KeyEvent
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	KeyEvent
Aop:30 - Estrogen receptor antagonism leading to reproductive dysfunction	KeyEvent
Aop:122 - Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	KeyEvent
Aop:123 - Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	KeyEvent

Biological Context

Level of Biological Organization
Organ

Organ term

Organ term
blood plasma

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	High	NCBI
Oryzias latipes	Oryzias latipes	High	NCBI
Danio rerio	Danio rerio	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Oviparous vertebrates synthesise yolk precursor proteins that are transported in the circulation for uptake by developing oocytes. Many invertebrates also synthesise vitellogenins that are taken up into developing oocytes via active transport mechanisms. However, invertebrate vitellogenins are transported in hemolymph or via other transport mechanisms rather than plasma.

Key Event Description

Vitellogenin synthesised in the liver is secreted into the blood and circulates to the ovaries for uptake.

How it is Measured or Detected

Vitellogenin concentrations in plasma are typically detected using enzyme linked Immunosorbent assay (ELISA; e.g., Korte et al. 2000; Tyler et al. 1996; Holbech et al. 2001; Fenske et al. 2001). Although less specific and/or sensitive, determination of alkaline-labile phosphate or Western blotting has also been employed.

References

- Fenske M, van Aerle R, Brack S, Tyler CR, Segner H. Development and validation of a homologous zebrafish (*Danio rerio* Hamilton-Buchanan) vitellogenin enzyme-linked immunosorbent assay (ELISA) and its application for studies on estrogenic chemicals. *Comp Biochem Physiol C Toxicol Pharmacol*. 2001. Jul;129(3):217-32.
- Holbech H, Andersen L, Petersen GI, Korsgaard B, Pedersen KL, Bjerregaard P. Development of an ELISA for vitellogenin in whole body homogenate of zebrafish (*Danio rerio*). *Comp Biochem Physiol C Toxicol Pharmacol*. 2001 Sep;130(1):119-31.
- Korte JJ, Kahl MD, Jensen KM, Mumtaz SP, Parks LG, LeBlanc GA, et al. 2000. Fathead minnow vitellogenin: complementary DNA sequence and messenger RNA and protein expression after 17 β -estradiol treatment. *Environmental Toxicology and Chemistry* 19(4): 972-981.
- Tyler C, van der Eerden B, Jobling S, Panter G, Sumpter J. 1996. Measurement of vitellogenin, a biomarker for exposure to oestrogenic chemicals, in a wide variety of cyprinid fish. *Journal of Comparative Physiology and Biology* 166: 418-426.
- Wahli W. 1988. Evolution and expression of vitellogenin genes. *Trends in Genetics*. 4:227-232.

Event: 309: Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development

Short Name: Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development

Key Event Component

Process	Object	Action
receptor-mediated endocytosis	vitellogenins	decreased
oocyte growth		decreased
oocyte development		decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:25 - Aromatase inhibition leading to reproductive dysfunction	KeyEvent
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	KeyEvent
Aop:30 - Estrogen receptor antagonism leading to reproductive dysfunction	KeyEvent
Aop:122 - Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	KeyEvent
Aop:123 - Unknown MIE leading to reproductive dysfunction via increased HIF-1 alpha transcription	KeyEvent

Biological Context

Level of Biological Organization
Cellular

Cell term

Cell term
oocyte

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Moderate	NCBI
Oryzias latipes	Oryzias latipes	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

Oviparous vertebrates and invertebrates. Although hormonal regulation of vitellogenin synthesis and mechanisms of vitellogenin transport from the site of synthesis to the ovary vary between vertebrates and invertebrates (Wahli 1988), in both vertebrates and invertebrates, vitellogenin is incorporated into oocytes and cleaved to form yolk proteins.

Key Event Description

Vitellogenin from the blood is selectively taken up by competent oocytes via receptor-mediated endocytosis. Although vitellogenin receptors mediate the uptake, opening of intercellular channels through the follicular layers to the oocyte surface as the oocyte reaches a “critical” size is thought to be a key trigger in allowing vitellogenin uptake (Tyler and Sumpter 1996). Once critical size is achieved, concentrations in the plasma and temperature are thought to impose the primary limits on uptake (Tyler and Sumpter 1996). Uptake of vitellogenin into oocytes causes considerable oocyte growth during vitellogenesis, accounting for up to 95% of the final egg size in many fish (Tyler and Sumpter 1996). Given the central role of vitellogenesis in oocyte maturation, vitellogenin accumulation is a prominent feature used in histological staging of oocytes (e.g., Leino et al. 2005; Wolf et al. 2004).

How it is Measured or Detected

Relative vitellogenin accumulation can be evaluated qualitatively using routine histological approaches (Leino et al. 2005; Wolf et al. 2004). Oocyte size can be evaluated qualitatively or quantitatively using routine histological and light microscopy and/or imaging approaches.

References

- Leino R, Jensen K, Ankley G. 2005. Gonadal histology and characteristic histopathology associated with endocrine disruption in the adult fathead minnow. *Environmental Toxicology and Pharmacology* 19: 85-98.
- Tyler C, Sumpter J. 1996. Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287-318.
- Wolf JC, Dietrich DR, Friederich U, Caunter J, Brown AR. 2004. Qualitative and quantitative histomorphologic assessment of fathead minnow *Pimephales promelas* gonads as an endpoint for evaluating endocrine-active compounds: a pilot methodology study. *Toxicol Pathol* 32(5): 600-612.

Event: 78: Reduction, Cumulative fecundity and spawning

Short Name: Reduction, Cumulative fecundity and spawning

Key Event Component

Process	Object	Action
egg quantity		decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:29 - Estrogen receptor agonism leading to reproductive dysfunction	KeyEvent
Aop:25 - Aromatase inhibition leading to reproductive dysfunction	KeyEvent
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	KeyEvent
Aop:30 - Estrogen receptor antagonism leading to reproductive dysfunction	KeyEvent
Aop:122 - Prollyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	AdverseOutcome
Aop:123 - Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	AdverseOutcome

Biological Context

Level of Biological Organization
Individual

Domain of Applicability**Taxonomic Applicability**

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	High	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	High	NCBI
Oryzias latipes	Oryzias latipes	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

Cumulative fecundity and spawning can, in theory, be evaluated for any egg laying animal.

Key Event Description

Spawning refers to the release of eggs. Cumulative fecundity refers to the total number of eggs deposited by a female, or group of females over a specified period of time.

How it is Measured or Detected

In laboratory-based reproduction assays (e.g., OECD Test No. 229; OECD Test No. 240), spawning and cumulative fecundity can be directly measured through daily observation of egg deposition and egg counts.

In some cases, fecundity may be estimated based on gonado-somatic index ([OECD 2008](#)).

Regulatory Significance of the AO

Cumulative fecundity is the most apical endpoint considered in the OECD 229 Fish Short Term Reproduction Assay. The OECD 229 assay serves as screening assay for endocrine disruption and associated reproductive impairment ([OECD 2012](#)). Fecundity is also an important apical endpoint in the Medaka Extended One Generation Reproduction Test (MEOGRT; [OECD Test Guideline 240](#); OECD 2015).

A variety of fish life cycle tests also include cumulative fecundity as an endpoint ([OECD 2008](#)).

References

- OECD 2008. Series on testing and assessment, Number 95. Detailed Review Paper on Fish Life-cycle Tests. OECD Publishing, Paris. [ENV/JM/MONO\(2008\)22](#).
- OECD (2015), *Test No. 240: Medaka Extended One Generation Reproduction Test (MEOGRT)*, OECD Publishing, Paris. DOI: <http://dx.doi.org/10.1787/9789264242258-en>
- OECD. 2012a. Test no. 229: Fish short term reproduction assay. Paris, France:Organization for Economic Cooperation and Development.

List of AO(s) in this AOP

Event: 360: Decrease, Population trajectory

Short Name: Decrease, Population trajectory

Key Event Component

Process	Object	Action
population growth rate		decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:23 - Androgen receptor agonism leading to reproductive dysfunction	AdverseOutcome
Aop:25 - Aromatase inhibition leading to reproductive dysfunction	AdverseOutcome
Aop:29 - Estrogen receptor agonism leading to reproductive dysfunction	AdverseOutcome
Aop:30 - Estrogen receptor antagonism leading to reproductive dysfunction	AdverseOutcome
Aop:100 - Cyclooxygenase inhibition leading to reproductive dysfunction via inhibition of female spawning behavior	AdverseOutcome
Aop:122 - Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	AdverseOutcome
Aop:123 - Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	AdverseOutcome
Aop:155 - Deiodinase 2 inhibition leading to reduced young of year survival via posterior swim bladder inflation	AdverseOutcome
Aop:156 - Deiodinase 2 inhibition leading to reduced young of year survival via anterior swim bladder inflation	AdverseOutcome
Aop:157 - Deiodinase 1 inhibition leading to reduced young of year survival via posterior swim bladder inflation	AdverseOutcome
Aop:158 - Deiodinase 1 inhibition leading to reduced young of year survival via anterior swim bladder inflation	AdverseOutcome
Aop:159 - Thyroperoxidase inhibition leading to reduced young of year survival via anterior swim bladder inflation	AdverseOutcome

Biological Context

Level of Biological Organization
Population

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
all species	all species		NCBI

Life Stage Applicability

Life Stage	Evidence
All life stages	Not Specified

Sex Applicability

Sex	Evidence
Unspecific	Not Specified

Consideration of population size and changes in population size over time is potentially relevant to all living organisms.

Key Event Description

Maintenance of sustainable fish and wildlife populations (i.e., adequate to ensure long-term delivery of valued ecosystem services) is an accepted regulatory goal upon which risk assessments and risk management decisions are based.

How it is Measured or Detected

Population trajectories, either hypothetical or site specific, can be estimated via population modeling based on measurements of vital rates or reasonable surrogates measured in laboratory studies. As an example, Miller and Ankley 2004 used measures of cumulative fecundity from laboratory studies with repeat spawning fish species to predict population-level consequences of continuous exposure.

Regulatory Significance of the AO

Maintenance of sustainable fish and wildlife populations (i.e., adequate to ensure long-term delivery of valued ecosystem services) is a widely accepted regulatory goal upon which risk assessments and risk management decisions are based.

References

- Miller DH, Ankley GT. 2004. Modeling impacts on populations: fathead minnow (*Pimephales promelas*) exposure to the endocrine disruptor 17 β -trenbolone as a case study. *Ecotoxicology and Environmental Safety* 59: 1-9.

Appendix 2: List of Key Event Relationships in the AOP

List of Adjacent Key Event Relationships

Relationship: 31: Agonism, Androgen receptor leads to Reduction, Gonadotropins, circulating concentrations

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor agonism leading to reproductive dysfunction	adjacent	Low	Low

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Low	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	Not Specified

Sex Applicability

Sex	Evidence
Female	Not Specified

At present, this relationship is assumed to operate in repeat spawning fish species with asynchronous oocyte development.

The extent to which the negative feedback mechanism proposed here is operable for fish species employing other reproductive strategies and/or for other vertebrates has not been extensively examined, to date.

Key Event Relationship Description

See biological plausibility, below.

Evidence Supporting this KER

Biological Plausibility

- Circulating concentrations of steroid hormones are tightly regulated via positive and negative feedback loops that operate through endocrine, autocrine, and/or paracrine mechanisms within the hypothalamic-pituitary-gonadal axis (Norris 2007).
- Gonadotropin (luteinizing hormone [LH] and follicle-stimulating hormone [FSH]) secretion from the pituitary is a key regulator of gonadal steroid biosynthesis.
- Negative feedback of androgens or estrogens at the level of the hypothalamus and/or pituitary can reduce gonadotropin secretion by pituitary gonadotropes either indirectly due to decreased GnRH signaling from the hypothalamus or directly through intrapituitary regulators of gonadotropin expression (e.g., activin, follistatin, inhibin) (Norris 2007; Habibi and Huggard 1998).
- While similar processes of negative feedback of sex steroids on gonadotropin expression and release have been established in fish (Levavi-Sivan et al. 2010), there are many remaining uncertainties about the exact mechanisms through which feedback takes place in fish as well as other vertebrates. For example, feedback is thought to involve a complex interplay of neurotransmitter signaling, kisspeptins, and the follistatin/inhibin/activin system (Trudeau et al. 2000; Trudeau 1997; Oakley et al. 2009; Cheng et al. 2007).
- In addition, the nature of the feedback produced by androgens is dependent on the concentration, form of the androgen (e.g., aromatizable versus non-aromatizable), life-stage and likely species (Habibi and Huggard 1998; Trudeau et al. 2000; Gopurappilly et al. 2013).
- The specific mechanisms through which negative feedback of AR agonists on the hypothalamus and pituitary are mediated in fish are not fully understood.
 - It is thought that GABAergic and dopaminergic neurons may be important mediators of sex steroid feedback on gonadotropin releasing hormone (GnRH) release from the hypothalamus (Trudeau et al. 2000; Trudeau 1997).
 - More recent evidence also suggests an important role of kisspeptin neurons, which have been shown to express both AR and ER α are important mediators of feedback response to circulating androgen concentrations (Oakley et al. 2009).
 - Follistatin expression in the pituitary has also been cited as a key regulator of gonadotropin expression that is directly regulated by androgens and estrogens (Cheng et al. 2007).
- Regardless of the exact mechanisms, negative feedback of androgens on GnRH and/or gonadotropin release from the hypothalamus and/or pituitary is a well established endocrine phenomenon.

Empirical Evidence

- There is a relatively strong body of evidence demonstrating that gonadectomy and/or treatment with potent AR antagonists can increase circulating concentrations of gonadotropins in fish and that those effects can be reversed by treatment with testosterone (reviewed in Habibi and Huggard 1998; Levavi-Sivan et al. 2010).

- However, we are currently unaware of any studies conducted with xenobiotic or pharmaceutical androgen agonists that measured effects on circulating gonadotropins.
- Empirical support for this linkage is largely lacking for most fish species, as antibodies capable of specifically detecting and distinguishing luteinizing hormone and follicle stimulating hormone have not yet been developed, despite many attempts.
- FSH and LH can be specifically measured for salmonids, but measurement methods for most other species are lacking.

Uncertainties and Inconsistencies

Due to uncertainties regarding the exact mechanisms through which exogenous androgens mediate a negative feedback response this initiation of a negative feedback response is not directly observable. Negative feedback would generally be inferred through a decrease in gonadotropin release and associated declines in circulating gonadotropin concentrations.

References

- Cheng GF, Yuen CW, Ge W. 2007. Evidence for the existence of a local activin follistatin negative feedback loop in the goldfish pituitary and its regulation by activin and gonadal steroids. *The Journal of endocrinology* 195(3): 373-384.
- Gopurappilly R, Ogawa S, Parhar IS. 2013. Functional significance of GnRH and kisspeptin, and their cognate receptors in teleost reproduction. *Frontiers in endocrinology* 4: 24.
- Habibi HR, Huggard DL. 1998. Testosterone regulation of gonadotropin production in goldfish. *Comparative biochemistry and physiology Part C, Pharmacology, toxicology & endocrinology* 119(3): 339-344.
- Levavi-Sivan B, Bogerd J, Mananos EL, Gomez A, Lareyre JJ. 2010. Perspectives on fish gonadotropins and their receptors. *General and comparative endocrinology* 165(3): 412-437.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Oakley AE, Clifton DK, Steiner RA. 2009. Kisspeptin signaling in the brain. *Endocrine reviews* 30(6): 713-743.
- Trudeau VL, Spanswick D, Fraser EJ, Larivière K, Crump D, Chiu S, et al. 2000. The role of amino acid neurotransmitters in the regulation of pituitary gonadotropin release in fish. *Biochemistry and Cell Biology* 78: 241-259.
- Trudeau VL. 1997. Neuroendocrine regulation of gonadotropin II release and gonadal growth in the goldfish, *Carassius auratus*. *Reviews of Reproduction* 2: 55-68.

Relationship: 143: Reduction, Gonadotropins, circulating concentrations leads to Reduction, Testosterone synthesis by ovarian theca cells

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor agonism leading to reproductive dysfunction	adjacent	High	Low

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Low	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	Not Specified

Sex Applicability

Sex	Evidence
Female	Not Specified

- A functional hypothalamic-pituitary-gonadal axis involving GnRH and gonadotropin-mediated regulation of reproductive functions is a vertebrate trait (Sower et al. 2009).
- The taxonomic applicability of this key event is limited to chordates.
- CYP11a, one of the critical enzymes for testosterone synthesis has only been found in amphioxus or vertebrates (Baker 2011). Consequently, taxonomic relevance of this KER is likely restricted to that domain.

Key Event Relationship Description

See biological plausibility below.

Evidence Supporting this KER

Biological Plausibility

- In mammals androgen production by theca cells is largely under control of LH (Norris 2007; Young and McNeilly 2010).
- In fish, the differential role of LH versus FSH has been more difficult to define, in part due to the inability to specifically measure LH and FSH in most fish species and the parallel fluctuations of LH and FSH expression in many species.
- Regardless of the differential effects of the two gonadotropins there is little dispute that gonadotropins stimulate gonadal steroid production and that the production of androgens (e.g., androstenedione, testosterone), which are the precursors for estrogen synthesis occurs in the theca cells (Payne and Hales 2004; Young and McNeilly 2010; Miller 1988; Nagahama et al. 1993).

Empirical Evidence

There is a strong weight of evidence establishing the role of gonadotropins in stimulating gonadal steroidogenesis. This relationship is widely accepted.

Uncertainties and Inconsistencies

No significant inconsistencies identified to date, although comprehensive literature review as not conducted.

References

- Baker ME. Origin and diversification of steroids: co-evolution of enzymes and nuclear receptors. *Mol Cell Endocrinol.* 2011 Mar 1;334(1-2):14-20. doi:10.1016/j.mce.2010.07.013.
- Miller WL. 1988. Molecular biology of steroid hormone synthesis. *Endocrine reviews* 9(3): 295-318.
- Nagahama Y, Yoshikumi M, Yamashita M, Sakai N, Tanaka M. 1993. Molecular endocrinology of oocyte growth and maturation in fish. *Fish Physiology and Biochemistry* 11: 3-14.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Payne AH, Hales DB. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine reviews* 25(6): 947-970.
- Young JM, McNeilly AS. 2010. Theca: the forgotten cell of the ovarian follicle. *Reproduction* 140(4): 489-504.

Relationship: 302: Reduction, Testosterone synthesis by ovarian theca cells leads to Reduction, 17beta-estradiol synthesis by ovarian granulosa cells

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor agonism leading to reproductive dysfunction	adjacent	High	Low

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Moderate	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	Moderate

Sex Applicability

Sex	Evidence
Unspecific	Not Specified

Enzymes required for testosterone and 17 β -estradiol synthesis are only found in vertebrates and amphioxus (Markov et al. 2009; Baker 2011). They are not present in invertebrates. Consequently, this KER is not applicable to invertebrates.

24 or 48 h of exposure (Ekman et al. 2011). Reductions in ex vivo T production preceded significant reductions in ex vivo E2 production.

- Ketoconazole is a fungicide thought to inhibit CYP11A and CYP17 (both involved in theca cell androgen production) with greater potency than it inhibits CYP19 (aromatase) (Villeneuve et al. 2007). Ex vivo E2 and T production were significantly reduced following exposure to 30 or 300 µg ketoconazole/L (Ankley et al. 2012).

Uncertainties and Inconsistencies

No significant inconsistencies identified to date. However, the literature review on this topic has not been comprehensive.

References

- Ankley GT, Cavallin JE, Durhan EJ, Jensen KM, Kahl MD, Makynen EA, et al. 2012. A time-course analysis of effects of the steroidogenesis inhibitor ketoconazole on components of the hypothalamic-pituitary-gonadal axis of fathead minnows. *Aquatic toxicology* 114-115: 88-95.
- Baker ME. 2011. Origin and diversification of steroids: Co-evolution of enzymes and nuclear receptors. *Mol Cell Endocrinol* 334: 14-20.
- Breen MS, Villeneuve DL, Breen M, Ankley GT, Conolly RB. 2007. Mechanistic computational model of ovarian steroidogenesis to predict biochemical responses to endocrine active compounds. *Annals of biomedical engineering* 35(6): 970-981.
- Ekman DR, Villeneuve DL, Teng Q, Ralston-Hooper KJ, Martinovic-Weigelt D, Kahl MD, et al. 2011. Use of gene expression, biochemical and metabolite profiles to enhance exposure and effects assessment of the model androgen 17beta-trenbolone in fish. *Environmental toxicology and chemistry / SETAC* 30(2): 319-329.
- Markov GV, Tavares R, Dauphin-Villemant C, Demeneix BA, Baker ME, Laudet V. Independent elaboration of steroid hormone signaling pathways in metazoans. *Proc Natl Acad Sci U S A*. 2009 Jul 21;106(29):11913-8. doi: 10.1073/pnas.0812138106.
- Miller WL. 1988. Molecular biology of steroid hormone synthesis. *Endocrine reviews* 9(3): 295-318.
- Nagahama Y, Yoshikumi M, Yamashita M, Sakai N, Tanaka M. 1993. Molecular endocrinology of oocyte growth and maturation in fish. *Fish Physiology and Biochemistry* 11: 3-14.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Payne AH, Hales DB. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine reviews* 25(6): 947-970.
- Quignot N, Bois FY. 2013. A computational model to predict rat ovarian steroid secretion from in vitro experiments with endocrine disruptors. *PloS one* 8(1): e53891.
- Senthilkumaran B, Yoshikuni M, Nagahama Y. A shift in steroidogenesis occurring in ovarian follicles prior to oocyte maturation. *Mol Cell Endocrinol*. 2004 Feb 27;215(1-2):11-8.

- Shoemaker JE, Gayen K, Garcia-Reyero N, Perkins EJ, Villeneuve DL, Liu L, et al. 2010. Fathead minnow steroidogenesis: in silico analyses reveals tradeoffs between nominal target efficacy and robustness to cross-talk. *BMC systems biology* 4: 89.
- Villeneuve DL, Ankley GT, Makynen EA, Blake LS, Greene KJ, Higley EB, et al. 2007. Comparison of fathead minnow ovary explant and H295R cell-based steroidogenesis assays for identifying endocrine-active chemicals. *Ecotoxicol Environ Saf* 68(1): 20-32.

Relationship: 5: Reduction, 17beta-estradiol synthesis by ovarian granulosa cells leads to Reduction, Plasma 17beta-estradiol concentrations

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Aromatase (Cyp19a1) reduction leading to impaired fertility in adult female	adjacent	High	
Aromatase inhibition leading to reproductive dysfunction	adjacent	High	Moderate
Androgen receptor agonism leading to reproductive dysfunction	adjacent	High	Low
Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	adjacent	High	Moderate
Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	adjacent		

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
human	Homo sapiens		NCBI
mouse	Mus musculus	Moderate	NCBI
rat	Rattus norvegicus	High	NCBI
fathead minnow	Pimephales promelas	Moderate	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

Key enzymes needed to synthesise 17 β -estradiol first appear in the common ancestor of amphioxus and vertebrates (Baker 2011). While some E2 synthesis can occur in other tissues, the ovary is recognized as the major source of 17 β -estradiol synthesis in female

vertebrates. Endocrine actions of ovarian E2 are facilitated through transport via the plasma. Consequently, this key event relationship is applicable to most female vertebrates.

Key Event Relationship Description

See plausibility, below.

Evidence Supporting this KER

Biological Plausibility

While brain, interrenal, adipose, and breast tissue (in mammals) are capable of synthesising estradiol, the gonads are generally considered the major source of circulating estrogens in vertebrates, including fish (Norris 2007). Consequently, if estradiol synthesis by ovarian granulosa cells is reduced, plasma E2 concentrations would be expected to decrease unless there are concurrent reductions in the rate of E2 catabolism. Synthesis in other tissues generally plays a paracrine role only, thus the contribution of other tissues to plasma E2 concentrations can generally be considered negligible.

Empirical Evidence

Fish

- In multiple studies with aromatase inhibitors (e.g., fadrozole, prochloraz), significant reductions in ex vivo E2 production have been linked to, and shown to precede, reductions in circulating E2 concentrations (Villeneuve et al. 2009; Skolness et al. 2011). It is also notable that compensatory responses at the level of ex vivo steroid production (i.e., rate of E2 synthesis per unit mass of tissue) tend to precede recovery of plasma E2 concentrations following an initial insult (Villeneuve et al. 2009; Ankley et al. 2009a; Villeneuve et al. 2013).
- Ex vivo E2 production by ovary tissue collected from female fish exposed to 30 or 300 µg ketoconazole/L showed significant decreases prior to significant effects on plasma estradiol being observed (Ankley et al. 2012).
- Ekman et al. (2011) reported significant reductions in ex vivo E2 production and plasma E2 concentrations in female fathead minnows exposed to 0.05 µg/L 17β-trenbolone. The effect on plasma E2 was observed at an earlier time point (24 h, versus 48 h for E2 production).
- Rutherford et al. (2015) reported significant reductions in both E2 production and circulating E2 concentrations in female *Fundulus heteroclitus* exposed to 5α-dihydrotestosterone or 17α-methyltestosterone for 14 d. The effects were equipotent in the case of 17α-methyltestosterone, but in the case of 5α-dihydrotestosterone, the effect on plasma E2 could be detected at a lower dose (10 µg/L) than that at which a significant effect on E2 production was detected (100 µg/L).
- In female *Fundulus heteroclitus* exposed to 17α-methyltestosterone for 7 or 14 d, both E2 production and plasma E2 were impacted at the same exposure concentrations (Sharpe et al. 2004).

Mammals

- MEHP /DEHP, mice, ex vivo DEHP (10 -100 µg/ml); MEHP (0.1 and 10 µg/ml) dose dependent reduction E2 production (Gupta et al., 2010)
- DEHP, rat, in vivo 300-600 mg/kg/day, dose dependent reduction of E2 plasma levels (Xu et al., 2010)

Evidence for rodent and human models is summarized in Table 1.

Table 1. Summary of the experimental data for decrease E2 production and decreased E2 levels. IC50- half maximal inhibitory concentration values reported if available, otherwise the concentration at which the effect was observed.

Compound class	Species	Study type	Dose	E2 production/levels	Reference
Phthalates (DEHP)	rat	ex vivo	1500 mg/kg/day	Reduced/increased E2 production in ovary culture	(Laskey & Berman, 1993)
Phthalates (MEHP)	rat	in vitro	From 50 µM	Reduced E2 production (concentration and time dependent in Granulosa cell)	(Davis, Weaver, Gaines, & Heindel, 1994)
Phthalates (MEHP)	rat	in vitro	100-200µM	reduction E2 production (dose dependent)	(Lovekamp & Davis, 2001)
Phthalates (DEHP)	rat	in vivo	300-600 mg/kg/day	reduction E2 levels dose dependent	(Xu et al., 2010),
Phthalates (MEHP)	human	in vitro	IC(50)= 49-138 µM (dependent on the stimulant)	reduction E2 production (dose dependent)	(Reinsberg, Wegener-Toper, van der Ven, van der Ven, & Klingmueller, 2009)
Phthalates (MEHP/DEHP)	mice	ex vivo	DEHP (10 - 100 g/ml); MEHP (0.1 and 10 g/ml)	reduction E2 production (dose dependent)	(Gupta et al., 2010)

Uncertainties and Inconsistencies

Based on the limited set of studies available to date, there are no known inconsistencies.

References

- Ankley GT, Bencic DC, Cavallin JE, Jensen KM, Kahl MD, Makynen EA, et al. 2009a. Dynamic nature of alterations in the endocrine system of fathead minnows exposed to the fungicide prochloraz. *Toxicological sciences : an official journal of the Society of Toxicology* 112(2): 344-353.

- Ankley GT, Cavallin JE, Durhan EJ, Jensen KM, Kahl MD, Makynen EA, et al. 2012. A time-course analysis of effects of the steroidogenesis inhibitor ketoconazole on components of the hypothalamic-pituitary-gonadal axis of fathead minnows. *Aquatic toxicology* 114-115: 88-95.
- Baker ME. 2011. Origin and diversification of steroids: co-evolution of enzymes and nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 14-20.
- Davis, B J, R Weaver, L J Gaines, and J J Heindel. 1994. "Mono-(2-Ethylhexyl) Phthalate Suppresses Estradiol Production Independent of FSH-cAMP Stimulation in Rat Granulosa Cells." *Toxicology and Applied Pharmacology* 128 (2) (October): 224–8. doi:10.1006/taap.1994.1201.
- Ekman DR, Villeneuve DL, Teng Q, Ralston-Hooper KJ, Martinović-Weigelt D, Kahl MD, Jensen KM, Durhan EJ, Makynen EA, Ankley GT, Collette TW. Use of gene expression, biochemical and metabolite profiles to enhance exposure and effects assessment of the model androgen 17 β -trenbolone in fish. *Environ Toxicol Chem.* 2011 Feb;30(2):319-29. doi: 10.1002/etc.406.
- Gupta, Rupesh K, Jeffery M Singh, Tracie C Leslie, Sharon Meachum, Jodi a Flaws, and Humphrey H-C Yao. 2010. "Di-(2-Ethylhexyl) Phthalate and Mono-(2-Ethylhexyl) Phthalate Inhibit Growth and Reduce Estradiol Levels of Antral Follicles in Vitro." *Toxicology and Applied Pharmacology* 242 (2) (January 15): 224–30. doi:10.1016/j.taap.2009.10.011.
- Laskey, J.W., and E. Berman. 1993. "Steroidogenic Assessment Using Ovary Culture in Cycling Rats: Effects of Bis (2-Diethylhexyl) Phthalate on Ovarian Steroid Production." *Reproductive Toxicology* 7 (1) (January): 25–33. doi:10.1016/0890-6238(93)90006-S.
- Li Z, Kroll KJ, Jensen KM, Villeneuve DL, Ankley GT, Brian JV, et al. 2011a. A computational model of the hypothalamic: pituitary: gonadal axis in female fathead minnows (*Pimephales promelas*) exposed to 17 α -ethynylestradiol and 17 β -trenbolone. *BMC systems biology* 5: 63.
- Lovekamp, T N, and B J Davis. 2001. "Mono-(2-Ethylhexyl) Phthalate Suppresses Aromatase Transcript Levels and Estradiol Production in Cultured Rat Granulosa Cells." *Toxicology and Applied Pharmacology* 172 (3) (May 1): 217–24. doi:10.1006/taap.2001.9156.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Reinsberg, Jochen, Petra Wegener-Toper, Katrin van der Ven, Hans van der Ven, and Dietrich Klingmueller. 2009. "Effect of Mono-(2-Ethylhexyl) Phthalate on Steroid Production of Human Granulosa Cells." *Toxicology and Applied Pharmacology* 239 (1) (August 15): 116–23. doi:10.1016/j.taap.2009.05.022.
- Rutherford R, Lister A, Hewitt LM, MacLatchy D. Effects of model aromatizable (17 α -methyltestosterone) and non-aromatizable (5 α -dihydrotestosterone) androgens on the adult mummichog (*Fundulus heteroclitus*) in a short-term reproductive endocrine bioassay. *Comp Biochem Physiol C Toxicol Pharmacol.* 2015 Apr;170:8-18. doi: 10.1016/j.cbpc.2015.01.004.
- Sharpe RL, MacLatchy DL, Courtenay SC, Van Der Kraak GJ. Effects of a model androgen (methyl testosterone) and a model anti-androgen (cyproterone acetate) on reproductive endocrine endpoints in a short-term adult mummichog (*Fundulus heteroclitus*) bioassay. *Aquat Toxicol.* 2004 Apr 28;67(3):203-15.
- Shoemaker JE, Gayen K, Garcia-Reyero N, Perkins EJ, Villeneuve DL, Liu L, et al. 2010. Fathead minnow steroidogenesis: in silico analyses reveals tradeoffs

between nominal target efficacy and robustness to cross-talk. *BMC systems biology* 4: 89.

- Skolness SY, Durhan EJ, Garcia-Reyero N, Jensen KM, Kahl MD, Makynen EA, et al. 2011. Effects of a short-term exposure to the fungicide prochloraz on endocrine function and gene expression in female fathead minnows (*Pimephales promelas*). *Aquat Toxicol* 103(3-4): 170-178.
- Villeneuve DL, Breen M, Bencic DC, Cavallin JE, Jensen KM, Makynen EA, et al. 2013. Developing Predictive Approaches to Characterize Adaptive Responses of the Reproductive Endocrine Axis to Aromatase Inhibition: I. Data Generation in a Small Fish Model. *Toxicological sciences : an official journal of the Society of Toxicology*.
- Villeneuve DL, Mueller ND, Martinovic D, Makynen EA, Kahl MD, Jensen KM, et al. 2009. Direct effects, compensation, and recovery in female fathead minnows exposed to a model aromatase inhibitor. *Environ Health Perspect* 117(4): 624-631.
- Xu, Chuan, Ji-An Chen, Zhiquan Qiu, Qing Zhao, Jiaohua Luo, Lan Yang, Hui Zeng, et al. 2010. "Ovotoxicity and PPAR-Mediated Aromatase Downregulation in Female Sprague-Dawley Rats Following Combined Oral Exposure to Benzo[a]pyrene and Di-(2-Ethylhexyl) Phthalate." *Toxicology Letters* 199 (3) (December 15): 323-32. doi:10.1016/j.toxlet.2010.09.015.

Relationship: 252: Reduction, Plasma 17 β -estradiol concentrations leads to Reduction, Vitellogenin synthesis in liver

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Aromatase inhibition leading to reproductive dysfunction	adjacent	High	Moderate
Androgen receptor agonism leading to reproductive dysfunction	adjacent	High	Moderate
Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	adjacent	High	Moderate
Unknown MIE leading to reproductive dysfunction via increased HIF-1 α transcription	adjacent		

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	Not Specified

Sex Applicability

Sex	Evidence
Female	High

Key enzymes needed to synthesise 17 β -estradiol first appear in the common ancestor of amphioxus and vertebrates (Baker 2011). However, non-oviparous vertebrates do not require vitellogenin. Consequently, this KER is applicable to oviparous vertebrates.

Key Event Relationship Description

See Plausibility below.

Evidence Supporting this KER

Biological Plausibility

Vitellogenin synthesis in fish is localised in the liver and is well documented to be regulated by estrogens via interaction with estrogen receptors (Tyler et al. 1996; Tyler and Sumpter 1996; Arukwe and Goksøyr 2003). The vitellogenin gene contains estrogen responsive elements in its promoter region and site directed mutagenesis has shown these to be essential for estrogen-dependent expression of vitellogenin (Chang et al. 1992; Teo et al. 1998). Liver is not regarded as a major site of E2 synthesis (Norris 2007), therefore the majority of E2 in liver comes from the circulation.

- Estrogen regulates expression of the vitellogenin gene in the amphibian *Xenopus laevis* (Skipper and Hamilton, 1977).

Empirical Evidence

- Empirical support for estrogen-dependent regulation of vitellogenin synthesis:
 - Many studies have demonstrated that exposure of hepatocytes to estrogens in vitro or in vivo induce vitellogenin mRNA synthesis (e.g., see reviews by (Navas and Segner 2006; Iguchi et al. 2006)).
 - In female fathead minnows exposed to 17 β -trenbolone, significant reductions in plasma E2 concentrations preceded significant reductions in plasma VTG (Ekman et al. 2011).
 - Intra-arterial injection of the estrogen 17 α ethynyl estradiol into male rainbow trout causes vitellogenin induction with about a 12 h lag time before increasing from basal levels (Schultz et al. 2001).
- Specific empirical support for reductions in plasma E2 leading to reductions in hepatic vitellogenin synthesis:
 - In a number of time-course experiments with aromatase inhibitors (e.g., fadrozole, prochloraz), decreases in plasma estradiol concentrations precede decreases in plasma vitellogenin concentrations (Villeneuve et al. 2009; Skolness et al. 2011; Ankley et al. 2009b). Recovery of plasma E2 concentrations also precedes recovery of plasma VTG concentrations after cessation of exposure (Villeneuve et al. 2009; Ankley et al. 2009a; Villeneuve et al. 2013).
 - It was demonstrated in *Danio rerio* that in vivo exposure to the aromatase inhibitor letrozole significantly reduced the expression of mRNA transcripts coding for vtg1, vtg2, and $er\alpha$, all of which are known to be regulated by estrogens (Sun et al. 2010). However, similar effects were not observed in primary cultured hepatocytes from *Danio rerio*, indicating that letrozole's effects on vtg transcription were not direct.

Uncertainties and Inconsistencies

Based on the limited set of studies available to date, there are no known inconsistencies.

References

- Ankley GT, Bencic D, Cavallin JE, Jensen KM, Kahl MD, Makynen EA, et al. 2009b. Dynamic nature of alterations in the endocrine system of fathead minnows exposed to the fungicide prochloraz. *Toxicol Sci* 112(2): 344-353.
- Ankley GT, Bencic DC, Cavallin JE, Jensen KM, Kahl MD, Makynen EA, et al. 2009a. Dynamic nature of alterations in the endocrine system of fathead minnows exposed to the fungicide prochloraz. *Toxicological sciences : an official journal of the Society of Toxicology* 112(2): 344-353.
- Ankley GT, Miller DH, Jensen KM, Villeneuve DL, Martinovic D. 2008. Relationship of plasma sex steroid concentrations in female fathead minnows to reproductive success and population status. *Aquatic toxicology* 88(1): 69-74.
- Arukwe A, Goksøyr A. 2003. Eggshell and egg yolk proteins in fish: hepatic proteins for the next generation: oogenetic, population, and evolutionary implications of endocrine disruption. *Comparative Hepatology* 2(4): 1-21.
- Chang TC, Nardulli AM, Lew D, and Shapiro, DJ. 1992. The role of estrogen response elements in expression of the *Xenopus laevis* vitellogenin B1 gene. *Molecular Endocrinology* 6:3, 346-354.
- Ekman DR, Villeneuve DL, Teng Q, Ralston-Hooper KJ, Martinovic-Weigelt D, Kahl MD, et al. 2011. Use of gene expression, biochemical and metabolite profiles to enhance exposure and effects assessment of the model androgen 17beta-trenbolone in fish. *Environmental toxicology and chemistry / SETAC* 30(2): 319-329.
- Iguchi T, Irie F, Urushitani H, Tooi O, Kawashima Y, Roberts M, et al. 2006. Availability of in vitro vitellogenin assay for screening of estrogenic and anti-estrogenic activities of environmental chemicals. *Environ Sci* 13(3): 161-183.
- Li Z, Kroll KJ, Jensen KM, Villeneuve DL, Ankley GT, Brian JV, et al. 2011a. A computational model of the hypothalamic: pituitary: gonadal axis in female fathead minnows (*Pimephales promelas*) exposed to 17alpha-ethynylestradiol and 17beta-trenbolone. *BMC systems biology* 5: 63.
- Murphy CA, Rose KA, Rahman MS, Thomas P. 2009. Testing and applying a fish vitellogenesis model to evaluate laboratory and field biomarkers of endocrine disruption in Atlantic croaker (*Micropogonias undulatus*) exposed to hypoxia. *Environmental toxicology and chemistry / SETAC* 28(6): 1288-1303.
- Murphy CA, Rose KA, Thomas P. 2005. Modeling vitellogenesis in female fish exposed to environmental stressors: predicting the effects of endocrine disturbance due to exposure to a PCB mixture and cadmium. *Reproductive toxicology* 19(3): 395-409.
- Navas JM, Segner H. 2006. Vitellogenin synthesis in primary cultures of fish liver cells as endpoint for in vitro screening of the (anti)estrogenic activity of chemical substances. *Aquat Toxicol* 80(1): 1-22.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Schultz IR, Orner G, Merdink JL, Skillman A. 2001. Dose-response relationships and pharmacokinetics of vitellogenin in rainbow trout after intravascular administration of 17alpha-ethynylestradiol. *Aquatic toxicology* 51(3): 305-318.
- Skipper JK, Hamilton TH. 1977. Regulation by estrogen of the vitellogenin gene. *Proc Natl Acad Sci USA* 74:2384-2388.
- Skolness SY, Durhan EJ, Garcia-Reyero N, Jensen KM, Kahl MD, Makynen EA, et al. 2011. Effects of a short-term exposure to the fungicide prochloraz on

endocrine function and gene expression in female fathead minnows (*Pimephales promelas*). *Aquat Toxicol* 103(3-4): 170-178.

- Sun L, Wen L, Shao X, Qian H, Jin Y, Liu W, et al. 2010. Screening of chemicals with anti-estrogenic activity using in vitro and in vivo vitellogenin induction responses in zebrafish (*Danio rerio*). *Chemosphere* 78(7): 793-799.
- Teo BY, Tan NS, Lim EH, Lam TJ, Ding JL. A novel piscine vitellogenin gene: structural and functional analyses of estrogen-inducible promoter. *Mol Cell Endocrinol*. 1998 Nov 25;146(1-2):103-20. PubMed PMID: 10022768.
- Tyler C, Sumpter J. 1996. Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287-318.
- Tyler C, van der Eerden B, Jobling S, Panter G, Sumpter J. 1996. Measurement of vitellogenin, a biomarker for exposure to oestrogenic chemicals, in a wide variety of cyprinid fish. *Journal of Comparative Physiology and Biology* 166: 418-426.
- Villeneuve DL, Breen M, Bencic DC, Cavallin JE, Jensen KM, Makynen EA, et al. 2013. Developing Predictive Approaches to Characterize Adaptive Responses of the Reproductive Endocrine Axis to Aromatase Inhibition: I. Data Generation in a Small Fish Model. *Toxicological sciences: an official journal of the Society of Toxicology*.
- Villeneuve DL, Mueller ND, Martinovic D, Makynen EA, Kahl MD, Jensen KM, et al. 2009. Direct effects, compensation, and recovery in female fathead minnows exposed to a model aromatase inhibitor. *Environ Health Perspect* 117(4): 624-631.

Relationship: 315: Reduction, Vitellogenin synthesis in liver leads to Reduction, Plasma vitellogenin concentrations

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Aromatase inhibition leading to reproductive dysfunction	adjacent	High	Moderate
Androgen receptor agonism leading to reproductive dysfunction	adjacent	High	Moderate
Estrogen receptor antagonism leading to reproductive dysfunction	adjacent	High	Moderate
Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	adjacent	High	High
Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	adjacent		

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	Not Specified

Sex Applicability

Sex	Evidence
Unspecific	Not Specified

This KER primarily applies to taxa that synthesise vitellogenin in the liver which is transported elsewhere in the body via plasma (i.e., oviparous vertebrates).

Key Event Relationship Description

See biological plausibility, below.

Evidence Supporting this KER

Biological Plausibility

Liver is the major source of VTG protein production in fish (Tyler and Sumpter 1996; Arukwe and Goksøyr 2003). Protein production involves transcription and subsequent translation. The time-lag between decreases in transcription/translation and decreases in plasma VTG concentrations can be expected to be dependent on vitellogenin elimination half-lives.

Empirical Evidence

- In a number of time-course experiments with aromatase inhibitors, decreases in plasma estradiol concentrations precede decreases in plasma vitellogenin concentrations (Villeneuve et al. 2009; Skolness et al. 2011; Ankley et al. 2009b). Recovery of plasma E2 concentrations also precedes recovery of plasma VTG concentrations after cessation of exposure (Villeneuve et al. 2009; Ankley et al. 2009a; Villeneuve et al. 2013).
- In experiments with strong estrogens, increases in vtg mRNA synthesis precede increases in plasma VTG concentration (Korte et al. 2000; Schmid et al. 2002).
- Elimination half-lives for VTG protein have been determined for induced male fish, but to our knowledge, similar kinetic studies have not been done for reproductively mature females (Korte et al. 2000; Schultz et al. 2001).
- In male sheepshead minnows injected with E2, induction of VTG mRNA precedes induction of plasma VTG (Bowman et al. 2000).
- In male *Cichlasoma dimerus* exposed to octylphenol for 28 days and then held in clean water, decline in induced VTG mRNA concentrations precedes declines in induced plasma VTG concentrations (Genovese et al. 2012).

Uncertainties and Inconsistencies

There are no known inconsistencies between these KERs which are not readily explained on the basis of the expected dose, temporal, and incidence relationships between these two KERs. This applies across a significant body of literature in which these two KEs have been measured.

References

- Ankley GT, Bencic D, Cavallin JE, Jensen KM, Kahl MD, Makynen EA, et al. 2009b. Dynamic nature of alterations in the endocrine system of fathead minnows exposed to the fungicide prochloraz. *Toxicol Sci* 112(2): 344-353.
- Ankley GT, Bencic DC, Cavallin JE, Jensen KM, Kahl MD, Makynen EA, et al. 2009a. Dynamic nature of alterations in the endocrine system of fathead minnows exposed to the fungicide prochloraz. *Toxicological sciences : an official journal of the Society of Toxicology* 112(2): 344-353.
- Ankley GT, Miller DH, Jensen KM, Villeneuve DL, Martinovic D. 2008. Relationship of plasma sex steroid concentrations in female fathead minnows to reproductive success and population status. *Aquatic toxicology* 88(1): 69-74.

- Arukwe A, Goksøyr A. 2003. Eggshell and egg yolk proteins in fish: hepatic proteins for the next generation: oogenetic, population, and evolutionary implications of endocrine disruption. *Comparative Hepatology* 2(4): 1-21.
- Bowman CJ, Kroll KJ, Hemmer MJ, Folmar LC, Denslow ND. 2000. Estrogen-induced vitellogenin mRNA and protein in sheepshead minnow (*Cyprinodon variegatus*). *General and comparative endocrinology* 120(3): 300-313.
- Genovese G, Regueira M, Piazza Y, Towle DW, Maggese MC, Lo Nostro F. 2012. Time-course recovery of estrogen-responsive genes of a cichlid fish exposed to waterborne octylphenol. *Aquatic toxicology* 114-115: 1-13.
- Korte JJ, Kahl MD, Jensen KM, Mumtaz SP, Parks LG, LeBlanc GA, et al. 2000. Fathead minnow vitellogenin: complementary DNA sequence and messenger RNA and protein expression after 17 β -estradiol treatment. *Environmental Toxicology and Chemistry* 19(4): 972-981.
- Li Z, Kroll KJ, Jensen KM, Villeneuve DL, Ankley GT, Brian JV, et al. 2011a. A computational model of the hypothalamic: pituitary: gonadal axis in female fathead minnows (*Pimephales promelas*) exposed to 17 α -ethynylestradiol and 17 β -trenbolone. *BMC systems biology* 5: 63.
- Murphy CA, Rose KA, Rahman MS, Thomas P. 2009. Testing and applying a fish vitellogenesis model to evaluate laboratory and field biomarkers of endocrine disruption in Atlantic croaker (*Micropogonias undulatus*) exposed to hypoxia. *Environmental toxicology and chemistry / SETAC* 28(6): 1288-1303.
- Murphy CA, Rose KA, Thomas P. 2005. Modeling vitellogenesis in female fish exposed to environmental stressors: predicting the effects of endocrine disturbance due to exposure to a PCB mixture and cadmium. *Reproductive toxicology* 19(3): 395-409.
- Schmid T, Gonzalez-Valero J, Rufli H, Dietrich DR. 2002. Determination of vitellogenin kinetics in male fathead minnows (*Pimephales promelas*). *Toxicol Lett* 131(1-2): 65-74.
- Schultz IR, Orner G, Merdink JL, Skillman A. 2001. Dose-response relationships and pharmacokinetics of vitellogenin in rainbow trout after intravascular administration of 17 α -ethynylestradiol. *Aquatic toxicology* 51(3): 305-318.
- Skolness SY, Durhan EJ, Garcia-Reyero N, Jensen KM, Kahl MD, Makynen EA, et al. 2011. Effects of a short-term exposure to the fungicide prochloraz on endocrine function and gene expression in female fathead minnows (*Pimephales promelas*). *Aquat Toxicol* 103(3-4): 170-178.
- Tyler C, Sumpter J. 1996. Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287-318.
- Villeneuve DL, Breen M, Bencic DC, Cavallin JE, Jensen KM, Makynen EA, et al. 2013. Developing Predictive Approaches to Characterize Adaptive Responses of the Reproductive Endocrine Axis to Aromatase Inhibition: I. Data Generation in a Small Fish Model. *Toxicological sciences : an official journal of the Society of Toxicology*.
- Villeneuve DL, Mueller ND, Martinovic D, Makynen EA, Kahl MD, Jensen KM, et al. 2009. Direct effects, compensation, and recovery in female fathead minnows exposed to a model aromatase inhibitor. *Environ Health Perspect* 117(4): 624-631.

Relationship: 255: Reduction, Plasma vitellogenin concentrations leads to Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor agonism leading to reproductive dysfunction	adjacent	Moderate	Low
Aromatase inhibition leading to reproductive dysfunction	adjacent	Moderate	Low
Estrogen receptor antagonism leading to reproductive dysfunction	adjacent	Moderate	Low
Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	adjacent	Moderate	
Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	adjacent		

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Moderate	NCBI
Oryzias latipes	Oryzias latipes	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

This KER is expected to be primarily applicable to oviparous vertebrates that synthesise vitellogenin in hepatic tissue which is ultimately incorporated into oocytes present in the ovary.

Key Event Relationship Description

SEE BIOLOGICAL PLAUSIBILITY BELOW

Evidence Supporting this KER

Biological Plausibility

Vitellogenin synthesised in the liver and transported to the ovary via the circulation is the primary source of egg yolk proteins in fish (Wallace and Selman 1981; Tyler and Sumpter 1996; Arukwe and Goksøyr 2003). In many teleosts vitellogenesis can account for up to 95% of total egg size (Tyler and Sumpter 1996).

Empirical Evidence

In some (Ankley et al. 2002; Ankley et al. 2003; Lalone et al. 2013), but not all (Ankley et al. 2005; Sun et al. 2007; Skolness et al. 2013) fish reproduction studies, reductions in plasma vitellogenin have been associated with visible decreases in yolk protein content in oocytes and overall reductions in ovarian stage.

Uncertainties and Inconsistencies

Not all fish reproduction studies showing reductions in plasma vitellogenin have caused visible decreases in yolk protein content in oocytes and overall reductions in ovarian stage (Ankley et al. 2005; Sun et al. 2007; Skolness et al. 2013).

While plasma vitellogenin is well established as the only major source of vitellogenins to the oocyte, the extent to which a decrease will impact an ovary that has already developed vitellogenic staged oocytes is less certain. It would be assumed that the more rapid the turnover of oocytes in the ovary, the tighter the linkage between these KEs. Thus, repeat spawning species with asynchronous oocyte development that spawn frequently would likely be more vulnerable than annual spawning species with synchronous oocyte development that had already reached late vitellogenic stages.

References

- Ankley GT, Jensen KM, Durhan EJ, Makynen EA, Butterworth BC, Kahl MD, et al. 2005. Effects of two fungicides with multiple modes of action on reproductive endocrine function in the fathead minnow (*Pimephales promelas*). *Toxicol Sci* 86(2): 300-308.
- Ankley GT, Jensen KM, Makynen EA, Kahl MD, Korte JJ, Hornung MW, et al. 2003. Effects of the androgenic growth promoter 17-b-trenbolone on fecundity and reproductive endocrinology of the fathead minnow. *Environmental Toxicology and Chemistry* 22(6): 1350-1360.
- Ankley GT, Kahl MD, Jensen KM, Hornung MW, Korte JJ, Makynen EA, et al. 2002. Evaluation of the aromatase inhibitor fadrozole in a short-term reproduction assay with the fathead minnow (*Pimephales promelas*). *Toxicological Sciences* 67: 121-130.
- Arukwe A, Goksøyr A. 2003. Eggshell and egg yolk proteins in fish: hepatic proteins for the next generation: oogenetic, population, and evolutionary implications of endocrine disruption. *Comparative Hepatology* 2(4): 1-21.

- Li Z, Villeneuve DL, Jensen KM, Ankley GT, Watanabe KH. 2011b. A computational model for asynchronous oocyte growth dynamics in a batch-spawning fish. *Can J Fish Aquat Sci* 68: 1528-1538.
- Skolness SY, Blanksma CA, Cavallin JE, Churchill JJ, Durhan EJ, Jensen KM, et al. 2013. Propiconazole Inhibits Steroidogenesis and Reproduction in the Fathead Minnow (*Pimephales promelas*). *Toxicological sciences : an official journal of the Society of Toxicology* 132(2): 284-297.
- Sun L, Zha J, Spear PA, Wang Z. 2007. Toxicity of the aromatase inhibitor letrozole to Japanese medaka (*Oryzias latipes*) eggs, larvae and breeding adults. *Comp Biochem Physiol C Toxicol Pharmacol* 145(4): 533-541.
- Tyler C, Sumpter J. 1996. Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287-318.
- Wallace RA, Selman K. 1981. Cellular and dynamic aspects of oocyte growth in teleosts. *American Zoologist* 21: 325-343.

Relationship: 337: Reduction, Vitellogenin accumulation into oocytes and oocyte growth/development leads to Reduction, Cumulative fecundity and spawning

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Aromatase inhibition leading to reproductive dysfunction	adjacent	Moderate	Moderate
Androgen receptor agonism leading to reproductive dysfunction	adjacent	Moderate	Moderate
Estrogen receptor antagonism leading to reproductive dysfunction	adjacent	Moderate	Moderate
Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	adjacent		
Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	adjacent		

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Moderate	NCBI
Oryzias latipes	Oryzias latipes	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

On the basis of the taxonomic relevance of the two KEs linked via this KER, this KER is likely applicable to aquatic, oviparous, vertebrates which both produce vitellogenin and deposit eggs/sperm into an aquatic environment.

Key Event Relationship Description

SEE BIOLOGICAL PLAUSIBILITY BELOW

*Evidence Supporting this KER***Biological Plausibility**

Vitellogenesis is a critical stage of oocyte development and accumulated lipids and yolk proteins make up the majority of oocyte biomass (Tyler and Sumpter 1996). At least in mammals, maintenance of meiotic arrest is supported by signals transmitted through gap junctions between the granulosa cells and oocytes (Jamnongjit and Hammes 2005). Disruption of oocyte-granulosa contacts as a result of cell growth has been shown to coincide with oocyte maturation (Eppig 1994). However, it remains unclear whether the relationship between vitellogenin accumulation and oocyte growth and eventual maturation is causal or simply correlative.

Empirical Evidence

- At present, to our best knowledge there are no studies that definitively demonstrate a direct cause-effect relationship between impaired VTG accumulation into oocytes and impaired spawning. There is, however, strong correlative evidence. Across a range of laboratory studies with small fish, there is a robust and statistically significant correlation between reductions in circulating VTG concentrations and reductions in cumulative fecundity (Miller et al. 2007). To date, we are unaware of any fish reproduction studies which show a large reduction in circulating VTG concentrations, but not reductions in cumulative fecundity.
- Ankley et al. (2003) reported significant reductions in VTG accumulation in oocytes along with significant reductions in cumulative fecundity, although fecundity was significantly impacted at a lower dose (0.05 ug/L 17beta-trenbolone versus 0.5 ug/L for VTG accumulation).
- Kang et al. (2008) reported significant reductions in both VTG accumulation in oocytes and cumulative fecundity in Japanese medaka, with cumulative fecundity being impacted at slightly lower concentrations (0.047 ug 17alpha-methyltestosterone/L versus 0.088 ug/L).

Uncertainties and Inconsistencies

Based on the limited number of studies available that have examined both of these KEs, there are no known, unexplained, results that are inconsistent with this relationship.

References

- Ankley GT, Jensen KM, Makynen EA, Kahl MD, Korte JJ, Hornung MW, Henry TR, Denny JS, Leino RL, Wilson VS, Cardon MC, Hartig PC, Gray LE. Effects of the androgenic growth promoter 17-beta-trenbolone on fecundity and reproductive endocrinology of the fathead minnow. *Environ Toxicol Chem.* 2003 Jun;22(6):1350-60.

-
- Eppig JJ. 1994. Further reflections on culture systems for the growth of oocytes in vitro. *Human reproduction* 9(6): 974-976.
 - Jamnongjit M, Hammes SR. 2005. Oocyte maturation: the coming of age of a germ cell. *Seminars in reproductive medicine* 23(3): 234-241.
 - Kang IJ, Yokota H, Oshima Y, Tsuruda Y, Shimasaki Y, Honjo T. The effects of methyltestosterone on the sexual development and reproduction of adult medaka (*Oryzias latipes*). *Aquat Toxicol.* 2008 Apr 8;87(1):37-46. doi: 10.1016/j.aquatox.2008.01.010.
 - Li Z, Villeneuve DL, Jensen KM, Ankley GT, Watanabe KH. 2011b. A computational model for asynchronous oocyte growth dynamics in a batch-spawning fish. *Can J Fish Aquat Sci* 68: 1528-1538.
 - Miller DH, Jensen KM, Villeneuve DL, Kahl MD, Makynen EA, Durhan EJ, et al. 2007. Linkage of biochemical responses to population-level effects: a case study with vitellogenin in the fathead minnow (*Pimephales promelas*). *Environ Toxicol Chem* 26(3): 521-527.
 - Tyler C, Sumpter J. 1996. Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287-318.

Relationship: 94: Reduction, Cumulative fecundity and spawning leads to Decrease, Populosity trajectory

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Aromatase inhibition leading to reproductive dysfunction	adjacent	Moderate	Moderate
Androgen receptor agonism leading to reproductive dysfunction	adjacent	Moderate	Moderate
Estrogen receptor antagonism leading to reproductive dysfunction	adjacent	Moderate	Moderate
Prolyl hydroxylase inhibition leading to reproductive dysfunction via increased HIF1 heterodimer formation	adjacent		
Unknown MIE leading to reproductive dysfunction via increased HIF-1alpha transcription	adjacent		

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
All life stages	Not Specified

Sex Applicability

Sex	Evidence
Unspecific	Not Specified

Spawning generally refers to the release of eggs and/or sperm into water, generally by aquatic or semi-aquatic organisms. Consequently, by definition, this KER is likely applicable only to organisms that spend a portion of their life-cycle in or near aquatic environments.

Key Event Relationship Description

SEE BIOLOGICAL PLAUSIBILITY BELOW

Evidence Supporting this KER

Biological Plausibility

Using a relatively simple density-dependent population model and assuming constant young of year survival with no immigration/emigration, reductions in cumulative fecundity have been predicted to yield declines in population size over time (Miller and Ankley 2004). Under real-world environmental conditions, outcomes may vary depending on how well conditions conform with model assumptions. Nonetheless, cumulative fecundity can be considered one vital rate that contributes to overall population trajectories (Kramer et al. 2011).

Empirical Evidence

- Using a relatively simple density-dependent population model and assuming constant young of year survival with no immigration/emigration, reductions in cumulative fecundity have been predicted to yield declines in population size over time (Miller and Ankley 2004). However, it should be noted that the model was constructed in such a way that predicted population size is dependent on cumulative fecundity, therefore this is a fairly weak form of empirical support.
- In a study in which an entire lake was treated with 17alpha-ethynyl estradiol, Kidd et al. (2007) declines in fathead minnow population size were associated with signs of reduced fecundity.

Uncertainties and Inconsistencies

- Wester et al. (2003) and references cited therein suggest that although egg production is an endpoint of demographic significance, incomplete reductions of egg production may not translate in a simple manner to population reductions. Compensatory effects of reduced predation and reduced competition for limited food and/or habitat resources may offset the effects of incomplete reductions in egg production.
- Fish and other egg laying animals employ a diverse range of reproductive strategies and life histories. The nature of the relationship between reduced spawning frequency and cumulative fecundity and overall population trajectories will depend heavily on the life history and reproductive strategy of the species in question. Relationships developed for one species will not necessarily hold for other species, particularly those with differing life histories.

References

- Kidd KA, Blanchfield KH, Palace VP, Evans RE, Lazorchak JM, Flick RW. 2007. Collapse of a fish population after exposure to a synthetic estrogen. PNAS 104:8897-8901.
- Kramer VJ, Etterson MA, Hecker M, Murphy CA, Roesijadi G, Spade DJ, Spromberg JA, Wang M, Ankley GT. Adverse outcome pathways and ecological

risk assessment: bridging to population-level effects. *Environ Toxicol Chem.* 2011 Jan;30(1):64-76. doi: 10.1002/etc.375. PubMed PMID: 20963853

- Miller DH, Ankley GT. 2004. Modeling impacts on populations: fathead minnow (*Pimephales promelas*) exposure to the endocrine disruptor 17 β -trenbolone as a case study. *Ecotoxicology and Environmental Safety* 59: 1-9.
- Miller DH, Tietge JE, McMaster ME, Munkittrick KR, Xia X, Ankley GT. 2013. Assessment of Status of White Sucker (*Catostomus Commersoni*) Populations Exposed to Bleached Kraft Pulp Mill Effluent. *Environmental toxicology and chemistry / SETAC* (in press).
- Wester P, van den Brandhof E, Vos J, van der Ven L. 2003. Identification of endocrine disruptive effects in the aquatic environment - a partial life cycle assay with zebrafish. (RIVM Report). Bilthoven, the Netherlands: Joint Dutch Environment Ministry.

List of Non Adjacent Key Event Relationships

Relationship: 32: Agonism, Androgen receptor leads to Reduction, Testosterone synthesis by ovarian theca cells

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor agonism leading to reproductive dysfunction	non-adjacent	Moderate	Low

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	High	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

This KER is potentially relevant to sexually mature female vertebrates and amphioxus. It is not relevant to invertebrates.

- Androgen receptor orthologs are primarily limited to vertebrates (Baker 1997; Thornton 2001; Eick and Thornton 2011; Markov and Laudet 2011).
- Cytochrome P45011a (Cyp11a), a rate limiting enzyme for the production of testosterone, is specific to vertebrates and amphioxus (Markov et al. 2009; Baker et al. 2011; Payne and Hales, 2004).
- Cyp11a does not occur in invertebrates, as a result, they do not synthesise testosterone, nor other steroid intermediates required for testosterone synthesis (Markov et al. 2009; Payne and Hales, 2004).

Key Event Relationship Description

At present, a direct structural/functional linkage between androgen receptor agonism and reduced testosterone production by ovarian theca cells is not known. This linkage is thought to operate indirectly through endocrine feedback along the hypothalamic-pituitary-gonadal axis. Consequently, the relationship is supported primarily via association/correlation.

Evidence Supporting this KER

Biological Plausibility

Synthesis of the steroidogenic enzymes that catalyze the formation of testosterone from cholesterol as a precursor is stimulated by gonadotropins, particularly luteinizing hormone, whose synthesis and secretion are in turn regulated by gonadotropin releasing hormone (GnRH) released from the hypothalamus (Payne and Hales 2004; Norris 2007; Miller 1988). Negative feedback of circulating androgens (e.g., testosterone) on GnRH release from the hypothalamus and/or gonadotropin release from the pituitary is a well established physiological phenomenon in vertebrate endocrinology (Norris 2007). While similar processes of negative feedback of sex steroids on gonadotropin expression and release have been established in fish (Levavi-Sivan et al. 2010), there are many remaining uncertainties about the exact mechanisms through which feedback takes place in fish as well as other vertebrates. For example, feedback is thought to involve a complex interplay of neurotransmitter signaling, kisspeptins, and the follistatin/inhibin/activin system (Trudeau et al. 2000; Trudeau 1997; Oakley et al. 2009; Cheng et al. 2007). In addition, the nature of the feedback produced by androgens is dependent on the concentration, form of the androgen (e.g., aromatizable versus non-aromatizable), life-stage and likely species (Habibi and Huggard 1998; Trudeau et al. 2000; Gopurappilly et al. 2013). At present, such negative feedback responses in vivo provide a biologically plausible connection between androgen receptor agonism and reduced testosterone production in ovarian theca cells, but uncertainty regarding the details of the underlying biology and the relevant applicability domain remain.

Empirical Evidence

Direct evidence based on measurements of T production following exposure to established AR agonists:

- Ekman et al. (2011) reported significant reductions in ex vivo T production by ovary tissue following 24 or 48 h of in vivo exposure to the synthetic, non-aromatizable, AR agonist 17 β -trenbolone.
- Glinka et al. (2015) reported significant reductions in T production by ovary tissue collected from *Fundulus heteroclitus* that had been exposed to the model (non-aromatizable) androgen 5 α -dihydrotestosterone.
- Sharpe et al. (2004) reported that testosterone biosynthetic capacity was inhibited in female *Fundulus heteroclitus* following 7 d of in vivo exposure to 0.25 μ g methyl testosterone/L or 14 d of exposure to 0.1 μ g methyl testosterone/L. Note - methyl testosterone is an aromatizable androgen.
- Indirect evidence based on measurements of circulating T following exposure to established AR agonists:
- Ankley et al. (2003) showed that 17 β -trenbolone binds the *Pimephales promelas* androgen receptor, induced turbercle formation (an androgen regulated male

secondary sex characteristic in females), and reduced circulating concentrations of T and 11-ketotestosterone following 21 d of continuous exposure.

- Jensen et al. (2006) reported tubercle formation in females and reduced circulating concentrations of T in female *Pimephales promelas* exposed to 17 α -trenbolone for 21 d.
- LaLone et al. (2013) reported tubercle formation in female *Pimephales promelas* and papillary complexes on the anal fin (a male secondary sex characteristic) of female *Oryzias latipes* following 21 d of exposure to spironolactone, along with significant reductions in plasma T measured in *Pimephales promelas* (that endpoint was not measured in medaka due to limited plasma volumes).

Uncertainties and Inconsistencies

See biological plausibility section above regarding current uncertainties in the mechanisms through which AR agonists may reduce gonadotropin secretion.

- Rutherford et al. (2015) reported an increase in plasma T concentrations in female and no change in gonadal T production in *Fundulus heteroclitus* following 14 d of exposure to 100 μ g/L 5 α -dihydrotestosterone.

References

- Baker ME. 1997. Steroid receptor phylogeny and vertebrate origins. *Molecular and cellular endocrinology* 135(2): 101-107.
- Baker ME. 2011. Origin and diversification of steroids: Co-evolution of enzymes and nuclear receptors. *Mol Cell Endocrinol* 334: 14-20.
- Cheng GF, Yuen CW, Ge W. 2007. Evidence for the existence of a local activin follistatin negative feedback loop in the goldfish pituitary and its regulation by activin and gonadal steroids. *The Journal of endocrinology* 195(3): 373-384.
- Eick GN, Thornton JW. 2011. Evolution of steroid receptors from an estrogen-sensitive ancestral receptor. *Molecular and cellular endocrinology* 334(1-2): 31-38.
- Ekman DR, Villeneuve DL, Teng Q, Ralston-Hooper KJ, Martinović-Weigelt D, Kahl MD, Jensen KM, Durhan EJ, Makynen EA, Ankley GT, Collette TW. Use of gene expression, biochemical and metabolite profiles to enhance exposure and effects assessment of the model androgen 17 β -trenbolone in fish. *Environ Toxicol Chem.* 2011 Feb;30(2):319-29. doi: 10.1002/etc.406.
- Ankley GT, Jensen KM, Makynen EA, Kahl MD, Korte JJ, Hornung MW, Henry TR, Denny JS, Leino RL, Wilson VS, Cardon MC, Hartig PC, Gray LE. Effects of the androgenic growth promoter 17-beta-trenbolone on fecundity and reproductive endocrinology of the fathead minnow. *Environ Toxicol Chem.* 2003 Jun;22(6):1350-60.
- Glinka CO, Frasca S Jr, Provatas AA, Lama T, DeGuise S, Bosker T. The effects of model androgen 5 α -dihydrotestosterone on mummichog (*Fundulus heteroclitus*) reproduction under different salinities. *Aquat Toxicol.* 2015 Aug;165:266-76. doi: 10.1016/j.aquatox.2015.05.019.
- Gopurappilly R, Ogawa S, Parhar IS. 2013. Functional significance of GnRH and kisspeptin, and their cognate receptors in teleost reproduction. *Frontiers in endocrinology* 4: 24.

- Habibi HR, Huggard DL. 1998. Testosterone regulation of gonadotropin production in goldfish. *Comparative biochemistry and physiology Part C, Pharmacology, toxicology & endocrinology* 119(3): 339-344.
- Jensen KM, Makynen EA, Kahl MD, Ankley GT. Effects of the feedlot contaminant 17 α -trenbolone on reproductive endocrinology of the fathead minnow. *Environ Sci Technol*. 2006 May 1;40(9):3112-7.
- LaLone CA, Villeneuve DL, Cavallin JE, Kahl MD, Durhan EJ, Makynen EA, Jensen KM, Stevens KE, Severson MN, Blanksma CA, Flynn KM, Hartig PC, Woodard JS, Berninger JP, Norberg-King TJ, Johnson RD, Ankley GT. Cross-species sensitivity to a novel androgen receptor agonist of potential environmental concern, spironolactone. *Environ Toxicol Chem*. 2013 Nov;32(11):2528-41. doi: 10.1002/etc.2330.
- Levavi-Sivan B, Bogerd J, Mananos EL, Gomez A, Lareyre JJ. 2010. Perspectives on fish gonadotropins and their receptors. *General and comparative endocrinology* 165(3): 412-437.
- Li Z, Kroll KJ, Jensen KM, Villeneuve DL, Ankley GT, Brian JV, Sepúlveda MS, Orlando EF, Lazorchak JM, Kostich M, Armstrong B, Denslow ND, Watanabe KH. A computational model of the hypothalamic: pituitary: gonadal axis in female fathead minnows (*Pimephales promelas*) exposed to 17 α -ethynylestradiol and 17 β -trenbolone. *BMC Syst Biol*. 2011 May 5;5:63. doi: 10.1186/1752-0509-5-63.
- Markov GV, Laudet V. 2011. Origin and evolution of the ligand-binding ability of nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 21-30.
- Markov GV, Tavares R, Dauphin-Villemant C, Demeneix BA, Baker ME, Laudet V. Independent elaboration of steroid hormone signaling pathways in metazoans. *Proc Natl Acad Sci U S A*. 2009 Jul 21;106(29):11913-8. doi: 10.1073/pnas.0812138106.
- Miller WL. 1988. Molecular biology of steroid hormone synthesis. *Endocrine reviews* 9(3): 295-318.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Oakley AE, Clifton DK, Steiner RA. 2009. Kisspeptin signaling in the brain. *Endocrine reviews* 30(6): 713-743.
- Payne AH, Hales DB. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine reviews* 25(6): 947-970.
- Rutherford R, Lister A, Hewitt LM, MacLatchy D. Effects of model aromatizable (17 α -methyltestosterone) and non-aromatizable (5 α -dihydrotestosterone) androgens on the adult mummichog (*Fundulus heteroclitus*) in a short-term reproductive endocrine bioassay. *Comp Biochem Physiol C Toxicol Pharmacol*. 2015 Apr;170:8-18. doi: 10.1016/j.cbpc.2015.01.004.
- Sharpe RL, MacLatchy DL, Courtenay SC, Van Der Kraak GJ. Effects of a model androgen (methyl testosterone) and a model anti-androgen (cyproterone acetate) on reproductive endocrine endpoints in a short-term adult mummichog (*Fundulus heteroclitus*) bioassay. *Aquat Toxicol*. 2004 Apr 28;67(3):203-15.
- Thornton JW. 2001. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by ligand exploitation and serial genome expansions. *Proceedings of the National Academy of Sciences of the United States of America* 98(10): 5671-5676.

- Trudeau VL, Spanswick D, Fraser EJ, Larivière K, Crump D, Chiu S, et al. 2000. The role of amino acid neurotransmitters in the regulation of pituitary gonadotropin release in fish. *Biochemistry and Cell Biology* 78: 241-259.
- Trudeau VL. 1997. Neuroendocrine regulation of gonadotropin II release and gonadal growth in the goldfish, *Carassius auratus*. *Reviews of Reproduction* 2: 55-68.

Relationship: 1384: Agonism, Androgen receptor leads to Reduction, 17beta-estradiol synthesis by ovarian granulosa cells

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor agonism leading to reproductive dysfunction	non-adjacent	Moderate	Low

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	Moderate	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	Moderate

Sex Applicability

Sex	Evidence
Female	High

This KER is potentially applicable to sexually mature, female, vertebrates.

- Androgen receptor orthologs are primarily limited to vertebrates (Baker 1997; Thornton 2001; Eick and Thornton 2011; Markov and Laudet 2011).
- Key enzymes needed to synthesise 17 β -estradiol first appear in the common ancestor of amphioxus and vertebrates (Markov et al. 2009; Baker 2011).
- Key Event Relationship Description
- At present, a direct structural/functional link between androgen receptor agonism and reduced estradiol synthesis by ovarian granulosa cells is not known. The linkage is thought to operate indirectly via endocrine feedback along the hypothalamic-pituitary-gonadal axis and subsequent effects on the regulation of enzymes involved in ovarian steroidogenesis. This relationship is primarily supported by association/correlation.

Evidence Supporting this KER

Biological Plausibility

Synthesis of the steroidogenic enzymes that catalyze the formation of testosterone from cholesterol as a precursor as well as 17 β -estradiol (E2) from testosterone is stimulated by gonadotropins whose synthesis and secretion are in turn regulated by gonadotropin releasing hormone (GnRH) released from the hypothalamus (Payne and Hales 2004; Norris 2007; Miller 1988). Strong AR agonists are thought to exert negative feedback along the hypothalamic-pituitary-gonadal axis, leading to decreased stimulation of the steroidogenic pathway and subsequent declines in E2 production.

Empirical Evidence

Direct support for the effect of AR agonists on estrogen production by ovary tissue:

- Ekman et al. (2011) reported reductions in ex vivo E2 production by ovary tissue collected from fathead minnows exposed to 17 β -trenbolone in vivo. However, among four exposure durations assessed (1, 2, 4, 8 d), E2 production was only reduced following 2 d of exposure.
- Glinka et al. (2015) reported reductions in E2 production in female *Fundulus heteroclitus* following 21 d of exposure to 0.5 or 5.0 ug 5 α -dihydrotestosterone.
- Rutherford et al. (2015) demonstrate reduced E2 production in *Fundulus heteroclitus* following exposure to 100 ug 5 α -dihydrotestosterone (a non-aromatizable androgen) or 0.1 and 1.0 ug methyltestosterone (an aromatizable androgen) for 14 d,
- Sharpe et al. (2004) reported reductions in E2 production in female *Fundulus heteroclitus* following exposure to 0.25 or 1.0 ug methyltestosterone/L for 7d, and in females exposed to 0.001, 0.01, or 0.1 ug methyltestosterone/L for 14 d.

Uncertainties and Inconsistencies

The work of Ekman et al. (2011) demonstrates the effects can be transient due to complex compensatory behaviors.

References

- Baker ME. 1997. Steroid receptor phylogeny and vertebrate origins. *Molecular and cellular endocrinology* 135(2): 101-107.
- Baker ME. 2011. Origin and diversification of steroids: Co-evolution of enzymes and nuclear receptors. *Mol Cell Endocrinol* 334: 14-20.
- Eick GN, Thornton JW. 2011. Evolution of steroid receptors from an estrogen-sensitive ancestral receptor. *Molecular and cellular endocrinology* 334(1-2): 31-38.
- Ekman DR, Villeneuve DL, Teng Q, Ralston-Hooper KJ, Martinović-Weigelt D, Kahl MD, Jensen KM, Durhan EJ, Makynen EA, Ankley GT, Collette TW. Use of gene expression, biochemical and metabolite profiles to enhance exposure and effects assessment of the model androgen 17 β -trenbolone in fish. *Environ Toxicol Chem.* 2011 Feb;30(2):319-29. doi: 10.1002/etc.406.
- Markov GV, Laudet V. 2011. Origin and evolution of the ligand-binding ability of nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 21-30.
- Markov GV, Tavares R, Dauphin-Villemant C, Demeneix BA, Baker ME, Laudet V. Independent elaboration of steroid hormone signaling pathways in metazoans.

Proc Natl Acad Sci U S A. 2009 Jul 21;106(29):11913-8. doi: 10.1073/pnas.0812138106.

- Miller WL. 1988. Molecular biology of steroid hormone synthesis. *Endocrine reviews* 9(3): 295-318.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Payne AH, Hales DB. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine reviews* 25(6): 947-970.
- Thornton JW. 2001. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by ligand exploitation and serial genome expansions. *Proceedings of the National Academy of Sciences of the United States of America* 98(10): 5671-5676.

Relationship: 1385: Agonism, Androgen receptor leads to Reduction, Vitellogenin synthesis in liver

AOPs Referencing Relationship

AOP Name	Adjacency	Weight Evidence	of Quantitative Understanding
Androgen receptor agonism leading to reproductive dysfunction	non-adjacent	High	Low

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	High	NCBI
Danio rerio	Danio rerio	Moderate	NCBI
medaka	Oryzias latipes	Moderate	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

This KER is potentially applicable to reproductively mature, adult, oviparous vertebrates.

- Androgen receptor orthologs are primarily limited to vertebrates (Baker 1997; Thornton 2001; Eick and Thornton 2011; Markov and Laudet 2011).
- Oviparous vertebrates synthesise yolk precursor proteins that are transported in the circulation for uptake by developing oocytes. Many invertebrates also synthesise vitellogenins that are taken up into developing oocytes via active transport mechanisms. However, invertebrate vitellogenins are transported in hemolymph or via other transport mechanisms rather than plasma.

Key Event Relationship Description

At present, a direct structural/functional linkage between androgen receptor agonism and reduced plasma vitellogenin concentrations is not known. Consequently, the relationship is supported primarily via association/correlation.

Evidence Supporting this KER

Biological Plausibility

Synthesis of the steroidogenic enzymes that catalyze the formation of testosterone from cholesterol as a precursor as well as 17 β -estradiol (E2) from testosterone is stimulated by gonadotropins whose synthesis and secretion are in turn regulated by gonadotropin releasing hormone (GnRH) released from the hypothalamus (Payne and Hales 2004; Norris 2007; Miller 1988). Strong AR agonists are thought to exert negative feedback along the hypothalamic-pituitary-gonadal axis, leading to decreased stimulation of the steroidogenic pathway and subsequent declines in E2 production. E2 is known to be a major regulator of hepatic vitellogenin production (Tyler et al. 1996; Tyler and Sumpter 1996; Arukwe and Goksøyr 2003).

Empirical Evidence

Direct support for the effect of AR agonists on plasma vitellogenin (VTG) concentrations:

- Ekman et al. (2011) reported significant reductions in plasma VTG in female fathead minnows exposed to 0.5 ng 17beta-trenbolone/L for 4 or 8 d, or to 0.05 ng/L for 8 d.
- Ankley et al. (2010) showed significant reductions in plasma VTG in four independent experiments in which female fathead minnows were exposed to 17beta-trenbolone for 14 d and a fifth experiment where they were exposed for 21 d.
- Seki et al. (2006) showed significant reductions in plasma VTG in three different species of female fish (*Oryzias latipes*, *Danio rerio*, *Pimephales promelas*) following 21 d of exposure to 17beta-trenbolone.
- Villeneuve et al. (2016) reported significant reductions in plasma VTG following 22 d of exposure to 0.5 ng 17beta-trenbolone/L.
- Jensen et al. (2006) observed significant reductions in plasma VTG in female fathead minnows following exposure to 17alpha-trenbolone for 21 d.
- LaLone et al. (2013) reported significant reductions in plasma VTG in female fathead minnows exposed to 5 ug/L or higher concentrations of spironolactone.
- Rutherford et al. (2015) reported significant reductions in plasma VTG in female *Fundulus heteroclitus* after 14 d of exposure to 100 ug 5alpha-dihydrotestosterone/L as well as after exposure to 1 ug methyltestosterone/L.
- Sharpe et al. (2004), detected significant reductions in plasma VTG in female *Fundulus heteroclitus* exposed to 0.25 ug/L methyltestosterone for 7 d or 0.01 ug/L for 14 d.

Uncertainties and Inconsistencies

None noted.

References

- Ankley GT, Jensen KM, Kahl MD, Durhan EJ, Makynen EA, Cavallin JE, Martinović D, Wehmas LC, Mueller ND, Villeneuve DL. Use of chemical mixtures

- to differentiate mechanisms of endocrine action in a small fish model. *Aquat Toxicol*. 2010 Sep 1;99(3):389-96. doi: 10.1016/j.aquatox.2010.05.020.
- Arukwe A, Goksøyr A. 2003. Eggshell and egg yolk proteins in fish: hepatic proteins for the next generation: oogenetic, population, and evolutionary implications of endocrine disruption. *Comparative Hepatology* 2(4): 1-21.
 - Baker ME. 1997. Steroid receptor phylogeny and vertebrate origins. *Molecular and cellular endocrinology* 135(2): 101-107.
 - Eick GN, Thornton JW. 2011. Evolution of steroid receptors from an estrogen-sensitive ancestral receptor. *Molecular and cellular endocrinology* 334(1-2): 31-38.
 - Ekman DR, Villeneuve DL, Teng Q, Ralston-Hooper KJ, Martinović-Weigelt D, Kahl MD, Jensen KM, Durhan EJ, Makynen EA, Ankley GT, Collette TW. Use of gene expression, biochemical and metabolite profiles to enhance exposure and effects assessment of the model androgen 17 β -trenbolone in fish. *Environ Toxicol Chem*. 2011 Feb;30(2):319-29. doi: 10.1002/etc.406.
 - Jensen KM, Makynen EA, Kahl MD, Ankley GT. Effects of the feedlot contaminant 17 α -trenbolone on reproductive endocrinology of the fathead minnow. *Environ Sci Technol*. 2006 May 1;40(9):3112-7.
 - Jolly C, Katsiadaki I, Le Belle N, Mayer I, Dufour S. 2006. Development of a stickleback kidney cell culture assay for the screening of androgenic and anti-androgenic endocrine disrupters. *Aquatic toxicology* 79(2): 158-166.
 - LaLone CA, Villeneuve DL, Cavallin JE, Kahl MD, Durhan EJ, Makynen EA, Jensen KM, Stevens KE, Severson MN, Blanksma CA, Flynn KM, Hartig PC, Woodard JS, Berninger JP, Norberg-King TJ, Johnson RD, Ankley GT. Cross-species sensitivity to a novel androgen receptor agonist of potential environmental concern, spironolactone. *Environ Toxicol Chem*. 2013 Nov;32(11):2528-41. doi: 10.1002/etc.2330.
 - Li Z, Kroll KJ, Jensen KM, Villeneuve DL, Ankley GT, Brian JV, Sepúlveda MS, Orlando EF, Lazorchak JM, Kostich M, Armstrong B, Denslow ND, Watanabe KH. A computational model of the hypothalamic: pituitary: gonadal axis in female fathead minnows (*Pimephales promelas*) exposed to 17 α -ethynylestradiol and 17 β -trenbolone. *BMC Syst Biol*. 2011 May 5;5:63. doi: 10.1186/1752-0509-5-63.
 - Markov GV, Laudet V. 2011. Origin and evolution of the ligand-binding ability of nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 21-30.
 - Miller WL. 1988. Molecular biology of steroid hormone synthesis. *Endocrine reviews* 9(3): 295-318.
 - Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
 - Payne AH, Hales DB. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine reviews* 25(6): 947-970.
 - Rutherford R, Lister A, Hewitt LM, MacLatchy D. Effects of model aromatizable (17 α -methyltestosterone) and non-aromatizable (5 α -dihydrotestosterone) androgens on the adult mummichog (*Fundulus heteroclitus*) in a short-term reproductive endocrine bioassay. *Comp Biochem Physiol C Toxicol Pharmacol*. 2015 Apr;170:8-18. doi: 10.1016/j.cbpc.2015.01.004.
 - Seki M, Fujishima S, Nozaka T, Maeda M, Kobayashi K. Comparison of response to 17 beta-estradiol and 17 beta-trenbolone among three small fish species. *Environ Toxicol Chem*. 2006 Oct;25(10):2742-52.
 - Sharpe RL, MacLatchy DL, Courtenay SC, Van Der Kraak GJ. Effects of a model androgen (methyl testosterone) and a model anti-androgen (cyproterone acetate) on

reproductive endocrine endpoints in a short-term adult mummichog (*Fundulus heteroclitus*) bioassay. *Aquat Toxicol.* 2004 Apr 28;67(3):203-15.

- Thornton JW. 2001. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by ligand exploitation and serial genome expansions. *Proceedings of the National Academy of Sciences of the United States of America* 98(10): 5671-5676.
- Tyler C, Sumpter J. 1996. Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287-318.
- Tyler C, van der Eerden B, Jobling S, Panter G, Sumpter J. 1996. Measurement of vitellogenin, a biomarker for exposure to oestrogenic chemicals, in a wide variety of cyprinid fish. *Journal of Comparative Physiology and Biology* 166: 418-426.
- Villeneuve DL, Jensen KM, Cavallin JE, Durhan EJ, Garcia-Reyero N, Kahl MD, Leino RL, Makynen EA, Wehmas LC, Perkins EJ, Ankley GT. Effects of the antimicrobial contaminant triclocarban, and co-exposure with the androgen 17 β -trenbolone, on reproductive function and ovarian transcriptome of the fathead minnow (*Pimephales promelas*). *Environ Toxicol Chem.* 2017 Jan;36(1):231-242. doi: 10.1002/etc.3531.

Relationship: 1386: Reduction, Plasma 17 β -estradiol concentrations leads to Reduction, Plasma vitellogenin concentrations

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
Androgen receptor agonism leading to reproductive dysfunction	non-adjacent	High	Moderate
Aromatase inhibition leading to reproductive dysfunction	non-adjacent	High	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
fathead minnow	Pimephales promelas	High	NCBI
Fundulus heteroclitus	Fundulus heteroclitus	High	NCBI

Life Stage Applicability

Life Stage	Evidence
Adult, reproductively mature	High

Sex Applicability

Sex	Evidence
Female	High

This key event relationship likely applies to oviparous vertebrates only.

- Key enzymes needed to synthesise 17 β -estradiol first appear in the common ancestor of amphioxus and vertebrates (Baker 2011).
- Vitellogenesis is common to a range of egg-laying vertebrates and invertebrates. However, in the case of invertebrates, vitellogenins are transported via hemolymph rather than plasma and vitellogenesis is regulated by invertebrate hormones, not estradiol.
- Key Event Relationship Description
- There is not a direct structural/functional relationship between reduced concentrations of 17 β -estradiol in plasma and reduced plasma VTG concentrations. The relationship is thought to be mediated through additional events of hepatic estrogen receptor activation, vitellogenin protein synthesis in the liver, and subsequent secretion of vitellogenin into the plasma.

Evidence Supporting this KER

Biological Plausibility

The mechanisms through which 17 β -estradiol stimulates the transcription and translation of hepatic vitellogenin are well understood.

- In fish, see: Tyler et al. 1996; Tyler and Sumpter 1996; Arukwe and Goksøyr 2003; Teo et al. 1998
- In frogs: Chang et al. 1992; Wangh and Knowland 1975
- In reptiles: Ho et al. 1980
- Ho (1987)
- In birds: Deeley et al. 1975;
- 17 β -estradiol is not synthesised in significant amounts in the liver. Its synthesis originates in other tissues, principally the gonads. It is then transported to the liver and other tissues via circulation (Norris 2007; Payne and Hales 2004; Miller 1988; Nagahama et al. 1993).

Empirical Evidence

- Under conditions of continuous flow through exposure to 17 β -trenbolone (a non-aromatizable androgen receptor agonist), plasma E2 concentrations were reduced in female fathead minnows after 2, 4, or 8 d of exposure to concentrations of 0.05 ug/L or greater. Plasma VTG concentrations were significantly reduced only after 4 or 8 d of exposure, and at 4 d, only at a concentration of 0.5 ug/L, not 0.05 ug/L (Ekman et al. 2011).
- In the same study by Ekman et al. (2011), once exposure ceased, plasma E2 concentrations returned to control levels within 48 h, while plasma VTG concentrations remained significantly depressed until d. 4, post-exposure.
- Ankley et al. (2003) detected reductions in both plasma E2 and plasma VTG in female fathead minnows following 21 d of continuous exposure to 17 β -trenbolone. At 21 d, plasma E2 concentrations were impacted at concentrations of 0.5 ug/L or greater, while plasma VTG was significantly reduced at 0.05 ug/L or greater.
- Villeneuve et al. (2016) observed significant reductions in both plasma E2 and plasma VTG in female fathead minnows exposed to 0.5 ug/L 17 β -trenbolone for 14 d.
- Jensen et al. (2006) observed significant reductions in both plasma E2 and plasma VTG following exposure to 0.03 ug/L 17 α -trenbolone for 21 d.
- Following 21 d of continuous exposure to spironolactone, plasma E2 and plasma VTG were both significantly reduced in female fathead minnows. The lowest effect concentration for plasma E2 was 0.5 ug/L, while that for plasma VTG was 5 ug/L (LaLone et al. 2013).
- In female *Fundulus heteroclitus* exposed to 5 α -dihydrotestosterone for 14 d, plasma E2 was significantly reduced following exposure to 10 ug/L, while plasma VTG was reduced at 100 ug/L (Rutherford et al. 2015).
- In two experiments in which female *Fundulus heteroclitus* were exposed to 17 α -methyltestosterone, both plasma E2 and plasma VTG were significantly reduced. In both cases, plasma E2 was impacted at lower concentrations (0.25 ug/L in a 7 d study; 0.01 ug/L in a 14 d study) than plasma VTG (1 ug/L in the 7 d study; 0.1 ug/L in the 14 d study; Sharpe et al. 2004).
- In two experiments where plasma E2 and plasma VTG were measured in female fathead minnows (*Pimephales promelas*) in a time-course following continuous

exposure the aromatase inhibitor fadrozole, both plasma VTG and plasma E2 were depressed (Villeneuve et al. 2009; 2013). In both cases, following cessation of exposure, plasma E2 concentrations recovered to control levels before plasma VTG concentrations recovered (Villeneuve et al. 2009; 2013).

- Schroeder et al. (in preparation) reported effects on plasma E2 concentrations within 4 h of initiating exposure to 5 or 50 ug/L fadrozole. Plasma VTG concentrations did not decline until 24 h or later (Schroeder et al. 2009; Villeneuve et al. 2009; 2013).
- In female fathead minnows exposed to 300 ug/L prochloraz, plasma E2 concentrations were significantly reduced after 12 h of exposure, while plasma VTG concentrations were not significantly reduced until 24 h of exposure (Skolness et al. 2011).
- Ankley et al. (2009) reported significant reductions in plasma E2 in female fathead minnows following 24 h of exposure to 30 ug/L prochloraz. In the same study, plasma VTG concentrations did not significantly decline until 48 h of exposure, and then only at 300 ug/L prochloraz.
- In a 21 d exposure to prochloraz, plasma E2 was significantly reduced in females exposed to 300 ug prochloraz/L, while plasma VTG was significantly reduced in females exposed to 100 ug/L (Ankley et al. 2005).

Uncertainties and Inconsistencies

- In several studies, significant decreases in plasma vitellogenin are detected at lower concentrations than those that result in significant decreases in plasma E2. However, detection of differences in plasma VTG is often enhanced by the greater dynamic range in the concentrations of the protein that occur in plasma, compared to the dynamic range of steroid hormone concentrations.

References

- Ankley GT, Jensen KM, Makynen EA, Kahl MD, Korte JJ, Hornung MW, Henry TR, Denny JS, Leino RL, Wilson VS, Cardon MC, Hartig PC, Gray LE. Effects of the androgenic growth promoter 17-beta-trenbolone on fecundity and reproductive endocrinology of the fathead minnow. *Environ Toxicol Chem.* 2003 Jun;22(6):1350-60.
- Arukwe A, Goksøyr A. 2003. Eggshell and egg yolk proteins in fish: hepatic proteins for the next generation: oogenetic, population, and evolutionary implications of endocrine disruption. *Comparative Hepatology* 2(4): 1-21.
- Baker ME. 2011. Origin and diversification of steroids: co-evolution of enzymes and nuclear receptors. *Molecular and cellular endocrinology* 334(1-2): 14-20.
- Chang TC, Nardulli AM, Lew D, and Shapiro, DJ. 1992. The role of estrogen response elements in expression of the *Xenopus laevis* vitellogenin B1 gene. *Molecular Endocrinology* 6:3, 346-354\
- Chang TC, Nardulli AM, Lew D, Shapiro DJ. The role of estrogen response elements in expression of the *Xenopus laevis* vitellogenin B1 gene. *Mol Endocrinol.* 1992 Mar;6(3):346-54.
- Cheng WY, Zhang Q, Schroeder A, Villeneuve DL, Ankley GT, Conolly R. Computational Modeling of Plasma Vitellogenin Alterations in Response to Aromatase Inhibition in Fathead Minnows. *Toxicol Sci.* 2016 Nov;154(1):78-89.

- Deeley RG, Mullinix DP, Wetekam W, Kronenberg HM, Meyers M, Eldridge JD, Goldberger RF. Vitellogenin synthesis in the avian liver. Vitellogenin is the precursor of the egg yolk phosphoproteins. *J Biol Chem.* 1975 Dec 10;250(23):9060-6.
- Ekman DR, Villeneuve DL, Teng Q, Ralston-Hooper KJ, Martinović-Weigelt D, Kahl MD, Jensen KM, Durhan EJ, Makynen EA, Ankley GT, Collette TW. Use of gene expression, biochemical and metabolite profiles to enhance exposure and effects assessment of the model androgen 17 β -trenbolone in fish. *Environ Toxicol Chem.* 2011 Feb;30(2):319-29. doi: 10.1002/etc.406.
- Ho DM, L'Italien J, Callard IP. 1980. Studies on reptilian yolk:Chrysemys. *Comp. Biochem. Physiol.* 65B: 139-144.
- Ho SM. Endocrinology of vitellogenesis. In Norris DO, Jones RE Eds, *Hormones and reproduction in fishes, amphibians, and reptiles*, Plenum, New York, (1987), pp. 146-169.
- Jensen KM, Makynen EA, Kahl MD, Ankley GT. Effects of the feedlot contaminant 17 α -trenbolone on reproductive endocrinology of the fathead minnow. *Environ Sci Technol.* 2006 May 1;40(9):3112-7.
- LaLone CA, Villeneuve DL, Cavallin JE, Kahl MD, Durhan EJ, Makynen EA, Jensen KM, Stevens KE, Severson MN, Blanksma CA, Flynn KM, Hartig PC, Woodard JS, Berninger JP, Norberg-King TJ, Johnson RD, Ankley GT. Cross-species sensitivity to a novel androgen receptor agonist of potential environmental concern, spironolactone. *Environ Toxicol Chem.* 2013 Nov;32(11):2528-41. doi: 10.1002/etc.2330.
- Li Z, Kroll KJ, Jensen KM, Villeneuve DL, Ankley GT, Brian JV, Sepúlveda MS, Orlando EF, Lazorchak JM, Kostich M, Armstrong B, Denslow ND, Watanabe KH. A computational model of the hypothalamic: pituitary: gonadal axis in female fathead minnows (*Pimephales promelas*) exposed to 17 α -ethynylestradiol and 17 β -trenbolone. *BMC Syst Biol.* 2011 May 5;5:63. doi: 10.1186/1752-0509-5-63.
- Miller WL. 1988. Molecular biology of steroid hormone synthesis. *Endocrine reviews* 9(3): 295-318.
- Nagahama Y, Yoshikumi M, Yamashita M, Sakai N, Tanaka M. 1993. Molecular endocrinology of oocyte growth and maturation in fish. *Fish Physiology and Biochemistry* 11: 3-14.
- Norris DO. 2007. *Vertebrate Endocrinology*. Fourth ed. New York: Academic Press.
- Payne AH, Hales DB. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine reviews* 25(6): 947-970.
- Rutherford R, Lister A, Hewitt LM, MacLatchy D. Effects of model aromatizable (17 α -methyltestosterone) and non-aromatizable (5 α -dihydrotestosterone) androgens on the adult mummichog (*Fundulus heteroclitus*) in a short-term reproductive endocrine bioassay. *Comp Biochem Physiol C Toxicol Pharmacol.* 2015 Apr;170:8-18. doi: 10.1016/j.cbpc.2015.01.004.
- Sharpe RL, MacLatchy DL, Courtenay SC, Van Der Kraak GJ. Effects of a model androgen (methyl testosterone) and a model anti-androgen (cyproterone acetate) on reproductive endocrine endpoints in a short-term adult mummichog (*Fundulus heteroclitus*) bioassay. *Aquat Toxicol.* 2004 Apr 28;67(3):203-15.
- Teo BY, Tan NS, Lim EH, Lam TJ, Ding JL. A novel piscine vitellogenin gene: structural and functional analyses of estrogen-inducible promoter. *Mol Cell*

-
- Tyler C, Sumpter J. 1996. Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287-318.
 - Tyler C, van der Eerden B, Jobling S, Panter G, Sumpter J. 1996. Measurement of vitellogenin, a biomarker for exposure to oestrogenic chemicals, in a wide variety of cyprinid fish. *Journal of Comparative Physiology and Biology* 166: 418-426.
 - Villeneuve DL, Jensen KM, Cavallin JE, Durhan EJ, Garcia-Reyero N, Kahl MD, Leino RL, Makynen EA, Wehmas LC, Perkins EJ, Ankley GT. Effects of the antimicrobial contaminant triclocarban, and co-exposure with the androgen 17 β -trenbolone, on reproductive function and ovarian transcriptome of the fathead minnow (*Pimephales promelas*). *Environ Toxicol Chem.* 2017 Jan;36(1):231-242. doi: 10.1002/etc.3531.
 - Wangh LJ, Knowland J. 1975. Synthesis of vitellogenin in cultures of male and female frog liver regulated by estradiol treatment in vitro. *Proc. Nat. Acad. Sci.* 72: 3172-3175.

Appendix 3: Concordance Table

AOP:23 - Androgen receptor agonism leading to reproductive dysfunction
Dose-response and temporal concordance evaluation table.
Bold indicates lowest reported concentration (nominal) at which the effect was observed
Yellow highlight indicates the earliest time point at which the effect was observed.

KE1 (MIE)															KE2	KE3	KE4	KE5	KE6	KE7	KE8	KE9 (AO)	KE10 (AO)
															gonadotropins, circulating concentrations, reduction	testosterone synthesis by ovarian theca cells, reduction	ovarian granulosa cells, reduction	plasma 17β-estradiol concentrations, reduction	transcription and translation of vitellogenin in liver, reduction	plasma vitellogenin concentrations, reduction	vitellogenin uptake into oocytes and oocyte growth/development, reduction	cumulative fecundity and spawning, reduction	Population trajectory, decrease
Time point (d)	Reference	Chemical Initiator	Species	Concentrations tested	androgen receptor, agonism																		
n/a	Ankley et al. 2003, ET&C 22:1350-1360	17β-trenbolone	fathead minnow	0.00270-81.1 ug/L	2.70 ug/L IC50	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a									
0.333	Zhang et al. 2008, ET&C 27:2504-2511	17β-trenbolone	fathead minnow	2 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a									
1	Ekman et al. 2011 ET&C 30:319-329	17β-trenbolone	fathead minnow	0.05, 0.5 ug/L	n/a	n/a	0.05	n/a	n/a	n/a	n/a	n/a	n/a	n/a									
1.333	Zhang et al. 2008, ET&C 27:2504-2511	17β-trenbolone	medaka	2 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	2	n/a	n/a	n/a									
2	Ekman et al. 2011 ET&C 30:319-329	17β-trenbolone	fathead minnow	0.05, 0.5 ug/L	n/a	n/a	0.05	n/a	n/a	0.05	n/a	n/a	n/a	n/a									
4	Ekman et al. 2011 ET&C 30:319-329	17β-trenbolone	fathead minnow	0.05, 0.5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a									
7	Zhang et al. 2008 Aquat. Toxicol. 88:173-182	17β-trenbolone	medaka	0.05, 0.5, 5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	5	n/a	n/a	n/a									
7	Zhang et al. 2008, ET&C 27:2504-2511	17β-trenbolone	medaka	2 ug/L	n/a	n/a	n/a	n/a	n/a	2	n/a	n/a	n/a	n/a									
8	Ekman et al. 2011 ET&C 30:319-329	17β-trenbolone	fathead minnow	0.05, 0.5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	0.05	n/a	n/a	n/a									
14	Ankley et al. 2010, Aquat Toxicol. 99:389-396	17β-trenbolone	fathead minnow	0.5 ug/L	0.5 (b, c)	n/a	n/a	n/a	n/a	n/a	n/a	0.5 (c)	n/a	n/a									
14	Seki et al. 2006 ET&C 25:2742-2752	17β-trenbolone	fathead minnow	0.05, 0.5, 5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	4.06	n/a	n/a	n/a									
14	Seki et al. 2006 ET&C 25:2742-2752	17β-trenbolone	medaka	0.05, 0.5, 5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	0.365	n/a	n/a	n/a									
14	Seki et al. 2006 ET&C 25:2742-2752	17β-trenbolone	zebrafish	0.05, 0.5, 5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	0.351	n/a	n/a	n/a									
21	Ankley et al. 2003, ET&C 22:1350-1360	17β-trenbolone	fathead minnow	0.005, 0.05, 0.5, 5, 50 ug/L	0.05 (b)	n/a	0.5 (a)	n/a	0.5	n/a	0.05	0.5	0.05	n/a									
21	Ankley et al. 2010, Aquat Toxicol. 99:389-396	17β-trenbolone	fathead minnow	0.5 ug/L	0.5 (b)	n/a	n/a	n/a	n/a	n/a	0.5	n/a	n/a	n/a									
21	Seki et al. 2006 ET&C 25:2742-2752	17β-trenbolone	fathead minnow	0.05, 0.5, 5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	4.06	n/a	n/a	n/a									
21	Seki et al. 2006 ET&C 25:2742-2752	17β-trenbolone	medaka	0.05, 0.5, 5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	39.7	n/a	n/a	n/a									
21	Seki et al. 2006 ET&C 25:2742-2752	17β-trenbolone	zebrafish	0.05, 0.5, 5 ug/L	n/a	n/a	n/a	n/a	n/a	n/a	0.351	n/a	n/a	n/a									
22	Villeneuve et al. 2016 ET&C 36: 231-242	17β-trenbolone	fathead minnow	0.5 µg/L	0.5 (b)	n/a	n/a	n/a	0.5	n/a	0.5	0.5	0.5	n/a									
730	Miller and Ankley 2004 Ecotoxicol. Environ. Safe.	17β-trenbolone	fathead minnow	ug/L	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	0.027 ug/L (d)									
21	Jensen et al. 2006, ES&T 40:3112-3117	17α-trenbolone	fathead minnow	0.003, 0.01, 0.03, 0.10 ug/L	0.10 (b)	n/a	0.03 (a)	n/a	0.03	0.03	0.03	nd	0.01										
21	Laloue et al. 2013, ET&C 32:2528-2541	spironolactone	fathead minnow	0.5, 5, 50, 500 ug/L	50 (b)	n/a	0.5 (a)	n/a	0.5	n/a	5	5	5										
14	Rutherford et al. 2015 Compar. Biochem. Physiol. Part C. 170: 8-18	5α-dihydrotestosterone	Fundulus heteroclitus	10, 100 µg/L	n/a	n/a	n/a	100	10	n/a	100	n/a	n/a	n/a									
21	Glinka et al. 2015. Aquat. Toxicol. 165: 266-276	5α-dihydrotestosterone	heteroclitus	0.05, 0.5, 5.0 µg/L	n/a	n/a	0.5	0.5	n/a	n/a	n/a	n/a	0.05 (f)	n/a									
Aromatizable androgens [effects of aromatizable androgens may not be consistent with the AOP, which is considered applicable to non-aromatizable androgens]																							
21	Pawlowski et al. 2004 Aquat. Toxicol. 68: 277-291	17α-methyltestosterone	fathead minnow	0.1, 1, 5, 50 µg/L	1 (b)	n/a	n/a	n/a	n/a	n/a	n/a	5	n/a	n/a									
21	Kang et al. 2008 Aquat. Toxicol. 87: 37-46	17α-methyltestosterone	medaka	0.0225, 0.0468, 0.0881, 0.188, 0.380 µg/L	0.0468 (e)	n/a	n/a	n/a	n/a	0.188	n/a	0.0881	0.0468	n/a									
10	Korsgaard 2006 Mar Environ Res 62: 5205-5210	17α-methyltestosterone	Zoarcetes viviparus	0.01, 0.025, 0.050, 0.100, 0.500 µg/L	n/a	n/a	n/a	n/a	n/a	n/a	100 (but not 500)	n/a	n/a	n/a									
14	Rutherford et al. 2015 Compar. Biochem. Physiol. Part C. 170: 8-18	17α-methyltestosterone	Fundulus heteroclitus	0.1, 1µg/L	n/a	n/a	n/a	1	1	n/a	n/a	n/a	n/a	n/a									
7	Sharpe et al. 2004. Aquat. Toxicol. 67: 203-215	17α-methyltestosterone	Fundulus heteroclitus	0.25, 1 µg/L	n/a	n/a	0.25	0.25	0.25	n/a	1	n/a	n/a	n/a									
14	Sharpe et al. 2004. Aquat. Toxicol. 67: 203-215	17α-methyltestosterone	Fundulus heteroclitus	0.001, 0.01, 0.1 µg/L	n/a	n/a	0.1	0.001	0.01	n/a	0.1	n/a	n/a	n/a									

(a)indirect evidence based on plasma T concentrations

(b)indirect evidence based on tubercle formation in females

(c)consistent across 4 independent experiments

(d)modeled, 50% population reduction over 2 years.

(e)indirect evidence based on fin morphology

(f) at salinity of 16 ppt; reproduction was reduced overall at salinity 2 ppt