



3.7. FINLAND

Figure 3.7.1. **National agri-environmental and economic profile, 2002-04: Finland**



StatLink  <http://dx.doi.org/10.1787/300081400114>

1. Data refer to the period 2001-03.

2. Data refer to the year 2004.

Source: OECD Secretariat. For full details of these indicators, see Chapter 1 of the *Main Report*.

3.7.1. Agricultural sector trends and policy context

Primary agriculture's contribution to the economy is small and declining, accounting for 1.2% of GDP and 3.9% of employment in 2004 [1] (Figure 3.7.1). Agricultural productivity improved at around 1% annually between 1992 and 2003, with production remaining about stable (reflecting rising crop production largely offset by declining livestock output) and reduced input use [1, 2, 3, 4]. The intensity of farming has diminished with the area farmed over the period 1990-92 to 2002-04 declining by 12%, one of the largest decreases across the OECD, with even larger reductions in purchased farm input use: nitrogen (-20%) and phosphorus (-60%) inorganic fertilisers; pesticides (-9%); and on-farm direct energy consumption fell by 12% (Figure 3.7.2).

Finland's accession to the EU in 1995 brought major price and structural changes to farming [1, 2, 3, 4]. In 1995 while producer prices declined by 40-50%, although for milk the reduction was 15%, the decrease in input prices was less dramatic [1, 3, 4]. Also the average farm size has increased as their number declined, and a third of farmers are full time. The climate limits farm production, and the share of agricultural land is only 7% of the total land area, among the lowest share across the OECD, with crop production largely in the south, whereas livestock farming is concentrated in the central, eastern and northern regions [1, 5]. As agriculture is largely rain-fed, use of total water resources is extremely limited with irrigation, mainly for vegetables, accounting for only 4% of total farmland in 2000 [6, 7].

Farming is mainly supported under the Common Agricultural Policy (CAP) with support also provided through national expenditure within the CAP framework. Support to EU farmers has on average declined from 41% of farm receipts in the mid-1980s to 34% in 2002-04 (as measured by the OECD Producer Support Estimate – PSE) compared to the 31% OECD average. Nearly 70% of EU support to farmers is still output and input linked (compared to over 90% in the mid-1980s), which are the forms of support that most encourage production intensity. Finnish accession to the EU brought a considerable reduction in farm support with the EU PSE 50% compared to 67% for Finland in 1994 [8]. Finland's national expenditure to support farming was around EUR 1.0 (USD 1.25) billion and in conjunction with EU co-financing amounted to EUR 1.8 (USD 2.25) billion in 2004 of which about a third is allocated to agri-environmental schemes [1]. Agri-environmental support accounted for a third of total government environmental expenditure in 2004 [9].

Agri-environmental policies seek to reduce environmental damage and promote biodiversity and landscape conservation. The main agri-environmental measure is the Horizontal Rural Development Programme (HRDP, 2000-06) based on the EU Rural Development Programme [10, 11], with a new *Rural Development Strategy and Programme for 2007-13* approved by the EU Commission in June 2007. The key emphasis of the HRDP is on water protection, but measures also aim to limit air pollution, reduce pesticide risks, and promote conservation of biodiversity and cultural landscapes [12]. The HRDP consists of mandatory basic and additional measures (*general schemes*) and special measures (*specific schemes*). *General schemes* provide payments (EUR 259 [USD 324] million in 2004) for the adoption of nationwide agri-environmental practices, such as nutrient and pesticide management plans, creation of filter strips, and biodiversity and landscape conservation, with payments varying per hectare per year from EUR 93 (USD 116) for arable crops, EUR 117 (USD 146) for livestock farms, and EUR 333-484 (USD 416-605) for horticultural crops. Over 90% of working farms and cultivated area were covered by general schemes in 2004 [1], with 5% of farms receiving this support monitored to verify that the required measures are being adopted by farmers [6]. *Specific schemes* are more focused and are only provided if commitments under the general schemes have been applied by a given farmer. They provide payments (EUR 39 [USD 49] million in 2004) for covering investment and maintenance costs, such as the establishment of riparian buffer zones and wetlands, and promoting organic farming. The government has set a target to increase the area under organic farming to 15% of farmland by 2010, with 7% of agricultural land under organic management by 2004 [5].

Agriculture is influenced by national environmental and taxation policies. There are a number of measures that have economy-wide objectives to reduce eutrophication of water courses [13]. *Water Protection Targets* for 2005 set a 50% reduction target from 1991-95 levels for both nitrogen and phosphorus loads from agriculture. As these targets were not met, a new target for 2015 was agreed in 2007, seeking to reduce nitrogen and phosphorus loads from agriculture by 30% from 2002-05 levels. Under the *Environmental Protection Act* (2000) large-scale agricultural activities may undergo an Environmental Impact Assessment (EIA). The *Water Services Act* (2004) implements the EU *Water Framework Directive*, which for farming involves control of nutrient emissions under the EU *Nitrates Directive*, with action plans established at the water catchment level [1, 14]. A pesticide tax is levied on the pesticide industry, rather than farmers, averaging EUR 2 (USD 2.5) million annually to cover the administrative costs of registering new pesticides and improving pesticide productivity [9, 15, 16]. A tax on phosphorus fertiliser was introduced in 1990, but abolished in 1994 in preparation for entry

into the EU [10]. Policies promoting wood fuel production and agri-environmental management were implemented jointly in the early 1990s to assist rural areas and increase bioenergy production, especially as forestry is an integral part of farming with 95% of working farms having forests [17]. Payments are provided for production of energy crops (e.g. reed canary grass – *Phalaris arundinacea*) at EUR 45 (USD 56) per hectare [1]. Producers receiving farm support are entitled to reimbursement of energy (including fuel and electricity) and carbon dioxide taxes [16, 18], which was around EUR 245 (USD 304) per farm, equal to nearly EUR 16 (USD 20) million of budget revenue forgone in 2005.

International environmental agreements important to farming include [9]: those seeking to curb nutrient emissions and pesticides into the Baltic Sea (*HELCOM Convention*) [13]; the *Gothenburg Protocol* concerning ammonia emissions; greenhouse gases (*Kyoto Protocol*); and commitments under the *Convention of Biological Diversity*.

3.7.2. Environmental performance of agriculture

The main agri-environmental issues are water pollution and biodiversity conservation. Water pollution from the run-off and leaching of excess farm nutrients is a major source of degradation of aquatic ecosystems in both inland surface water and marine waters, and to a much lesser extent pesticides. Other agri-environmental issues of importance, include soil quality, emissions of ammonia and greenhouse gases, and conservation of cultural features in agricultural landscapes.

Soil erosion is primarily a concern for its off-farm impacts as a carrier of nutrients to water bodies. Soil water erosion is within the tolerable range and is usually under 1 tonne/hectare/year, reaching a maximum of 3 tonnes/hectare/year in some south-western areas [6, 19]. Although erosion rates are low it is a key factor affecting the quality of water as soil particles transport nutrients, especially phosphorus, to water bodies causing eutrophication and algal blooms [6, 19, 20]. With about 30% of arable land under plant cover or reduced tillage this has led to a decrease in the areas with the highest rates of erosion [6]. But research has shown that with mild winters over the past 20 years in Finland, nutrient loadings into water bodies were substantial and tended to override load reductions from decreased autumn tillage and increased use of green cover [20]. Greater adoption of low till has led to a rise in pesticide use, and potentially an increase in water pollution from their use, because of the need to use pesticides (Glyphosate), as perennial weeds are more abundant under low till than conventional tillage. However, besides reducing soil erosion and sediment bound nutrients, there are other environmental benefits from low till, such as greater soil carbon sequestration and habitat conservation for wild species [21].

The main focus on controlling water pollution has been on agriculture recently, since the control of urban and industrial pollution is well developed. For example, over 95% of phosphorus is removed from wastewater treatment plants [19]. Overall environmental pressure on inland and marine water quality has eased as there has been a substantial reduction of farm pollutants, largely nutrients (nitrogen and phosphorus) and pesticides. But while emissions have been lowered eutrophication of water continues and the state of water bodies has not improved over the past decade [1, 10].

There has been a substantial reduction over the past 15 years in agricultural nutrient surpluses (input minus output of nutrients, nitrogen – N – and phosphorus – P), among the highest reduction across OECD countries. The decrease in the quantity of nutrient surpluses was more significant for P (–65%) compared to N (–42%), with nutrient surpluses per hectare

of agricultural land now below both EU15 and OECD average levels (Figure 3.7.2). As a result of these changes there has been a considerable improvement in nutrient use efficiency (i.e. ratio of N/P output to N/P input), and while N use efficiency is now similar to the EU15 and OECD averages, for P use efficiency it is lower. The reduction in nutrient surpluses has been mainly due to the large reduction in nutrient inputs – inorganic fertiliser use and livestock numbers (i.e. less manure) – compared to a much smaller rise in nutrient uptake from crops and pasture. Over 90% of agricultural land was under a nutrient management plan in 2001-04, which includes conducting a soil nutrient test on farms once every 4 to 5 years.

Eutrophication of water has become the most serious environmental problem caused by agriculture [10]. Farming remains the single most important source of nutrient loading into water bodies, accounting for around 50% of N and 60% of P [1, 9]. A study for the period 1993 to 1998 estimated that agriculture's N contribution to river catchments varied from 35-85% in the intensively farmed south-west to 0-25% in the north [22]. The proportion of slightly eutrophic waters has increased and signs of early eutrophication have been pronounced in many small rivers, lakes and the Baltic Sea [9, 10, 23]. A study concluded that, annual variation notwithstanding, there has been little or no decrease in nutrient loadings into lakes from agriculture in southern Finland during the period 1976 to 2002 [23]. It is estimated that around 2% of shallow wells (and 1.5% of aquifers) in agricultural regions contained nitrates in excess of drinking water standards in 2002 [6]. The Gulf of Finland is one of the most eutrophied sub-basins of the Baltic Sea with a marked increase in algal blooms and dead zones, and nutrient loadings 2-3 times above that of the Baltic Sea average, although Finland is not among the major polluters of the Baltic Sea [10, 24, 25]. The share of Finnish agriculture in the total Finnish N (P) load into the Gulf of Finland rose from 31% (35%) in 1986-90 to 35% (48%) by 1997-2001, and while the quantity of agricultural N has increased over this period, for P it has declined [6].

Despite large reductions in nutrient surpluses this has not yet led to improvements in water quality (Figure 3.7.3). One reason for this is that caution is required in linking changes in nutrient balances to loading of nutrients into water, because of the importance of other factors, for example, nutrient management, crop rotations, and soil drainage systems [26]. Another reason is the long time lags of reductions in external nutrient loads showing up in changes in water quality because of the accumulation of nutrients in soils, especially for P [27]. Moreover, an increasing share of green set aside land has been converted to cereal production which has resulted in higher P losses [28], and there has been a decrease in the area of perennial vegetation from 34% to 28% of farmland between 1995 and 2002, which is important to slow soil erosion and the transport of nutrients into water [6, 28]. Also restrictions on the timing of manure applications and the pressure of time for farmers in the spring has led to them to spread manure on fields close to the farm, which already have high nutrient levels, rather than more distant fields with lower nutrient levels [17]. This problem is being accentuated with the trend toward the regional concentration of livestock production [17].

Pesticide use rose from the mid-1990s to 2003, although over the period 1990-92 to 2001-03 use declined by 9% (Figure 3.7.2) [1, 9]. The main reasons for the increase in pesticide use is due to the: wider adoption of reduced tillage and direct sowing; an increase in the arable area since the mid-1990s; establishment of buffer strips (and enlarged field boundaries); and a shift to pesticides that are used in larger doses [1, 9, 10, 21]. This trend has to some extent been offset by the major increase in the area under organic farming rising from a share in agricultural land area of less than 2% in the mid-1990s to over 7% in 2004,

among the highest share across OECD countries. The intensity of pesticide use is low compared to many OECD countries, mainly because of climatic conditions, especially colder winters, which limit pest populations. As a result the detection of pesticides in watercourses is infrequent and low, with an estimated 0.1-1% of active substances causing water pollution [10], although there is not yet regular monitoring of pesticides in water courses.

Agricultural ammonia emissions decreased by 13% between 1990-92 and 2001-03, a rate of reduction higher than achieved for the EU15 on average (-7%) (Figure 3.7.2). Farming accounts for almost all ammonia emissions (97% in 2001-03), largely from manure management and inorganic fertiliser use. The reduction of nitrogen fertilisers and livestock numbers over the past 15 years has been the main reasons for the decline in emissions. Finland has agreed to cut total ammonia emissions to 31 000 tonnes by 2010 under the *Gothenburg Protocol*, and by 2001-03 emissions totalled 33 000 tonnes, so a further 7% cut will be required to meet the target. While it is likely that the reduction in farm ammonia emissions has contributed to an overall decline in acidifying pollutants, easing pressure on ecosystems (terrestrial and aquatic) sensitive to excess acidity, there is little research or data on this issue.

There was a 14% decline in agricultural greenhouse gas (GHG) emissions between 1990-92 and 2002-04 (Figure 3.7.2). This compares to the 12% growth in GHG emissions from other sources across the country over this period, and a reduction in EU15 agricultural GHG emissions of 7%. The Kyoto commitment requires Finland under the *EU Burden Sharing Agreement* to stabilise total GHG emissions at 0% by 2008-12. In 2000-02 farming accounted for 10% of total GHG emissions, mainly methane and nitrous oxide [1, 29]. Agricultural emissions reductions are largely a co-benefit from decreasing nutrient loadings into the environment, including lower livestock numbers, reduced use of fertilisers, and improved manure management [29]. Even so, the increase in livestock numbers reared in slurry-based manure management systems, compared to solid storage or pasture, has led to a slight rise in methane but a reduction in nitrous oxide emissions. With the projected continued contraction of farming, the downward trend of agricultural GHGs is expected to persist up to 2010 [29]. **Carbon sequestration in agricultural soils** has the potential to reduce GHG emissions. With the increase in low tillage on cropland there was a small rise in GHG removals between 1990 and 2003 [29].

The reduction in on-farm energy consumption of -12% compared to a rise of 18% across the economy over the period 1990-92 to 2002-04 has also helped to lower GHG emissions, with agriculture accounting for 3% of total energy consumption (Figure 3.7.2). In 2005 the first large-scale agricultural biogas power plant was opened in Vehmaa, processing liquid manure from 20 pig farms [1]. In 2006 there were 17 000 hectares of energy crops, which is less than 0.5% of total arable land, but the area is expanding rapidly. Research suggests that production of reed canary grass for bioenergy, for example, is only profitable if located between 50-100 kilometres from the energy plant, but could provide environmental benefits not only in terms of lower GHG emissions, but also by reducing nutrient run-off and by replacing the use of peat for energy [3, 30].

The state of farmland biodiversity deteriorated over the period 1990 to 2004 [1, 31, 32, 33, 34]. There are, however, positive signs that recently the pressure from farming on biodiversity are easing, for example, for butterfly species (see below). Protection of Finnish **agricultural genetic resource diversity**, domestic plants and livestock breeds, combines both in situ and ex situ conservation [34, 35]. The diversity of most crop varieties and livestock breeds used in

production increased over the period 1990 to 2002. All nine “endangered” and “critical” livestock breeds (cattle, poultry and sheep breeds) were by 2004 maintained under *in situ* conservation programmes, compared with only two in 1985, while for crops there are limited areas of *in situ* conservation for certain fruit, berries, cereals and grass varieties [34]. Finland contributes plant material to the Nordic Gene Bank, while the *National Animal Genetic Resources Programme* finalised in 2004, covers *ex situ* conservation for livestock [34, 35].

Overall there has been a reduction in the abundance and richness of wild species associated with agriculture [1]. Around 25% of Finnish wild flora and fauna species use agricultural land as habitat, with nearly 30% of endangered species found in farmed habitats [36]. A comprehensive evaluation of threatened species associated with farmland habitats in 1985, 1990 and 2000, showed an increasing number of species under threat across five taxonomic groups – *Lepidoptera* (e.g. butterflies and moths), *Coleoptera* (e.g. beetles), *Hymenoptera* (e.g. bees, ants), vascular plants (e.g. ferns) and macro-fungi – although the increase was partly due to improved monitoring [31]. Moreover, the latest survey in 2000 showed a rise in the number of threatened species that was higher than in earlier years [37]. There is, however, great variation in the numbers of threatened species across different types of farmland habitats, with almost 50% of them species of dry meadows, and a further 25% found on marginal agricultural habitats, such as field boundaries and field and forest margins [31]. About 20% of all endangered plant species are found in agricultural habitats, but around 60% of these species are threatened by the disappearance of pasture or forest pastures following the cessation of grazing and mowing [33, 38].

Overall populations of many farmland birds also declined from the late 1970s to 2005, although this masks trends of individual species, as some bird species numbers have risen [33, 37]. The decrease in numbers of certain bird species is of particular importance as Finland is host to some of the largest European populations of the threatened ortolan bunting (*Emberiza hortulana*), northern wheatear (*Oenanthe oenanthe*) and whinchat (*Saxicola rubetra*) [39]. The deteriorating trend in bird populations on farmland habitats is also apparent for other species, including insect pollinators, and dung beetles [33, 40]. For butterflies, however, monitoring data for the period 1999 to 2006 indicate an increase in grassland and field margin butterfly species (Figure 3.7.4).

Changes in the quality of semi-natural farmed habitats is a key reason for the decline in wild species linked to agriculture [1, 31, 33]. But other factors are also important in leading to adverse impacts on wild species including: changes in cropping patterns (e.g. increase in the area of spring cereals and reduction of winter cereals); greater use of sub-surface drainage which has led to the removal of ditches [1, 39]; effects of pesticides; and the lack of economic incentives for farmers to provide ecosystem services [41]. During the 1990s up to 2004 the area of **semi-natural extensive pasture** increased by about 15 000 hectares less than 1% of the total agricultural land area in 2004, largely because agri-environmental payments have encouraged their conservation [6, 36]. But while the total area of pasture has expanded, there are concerns that its quality has diminished, including the fragmentation of pasture into patches, and diminution in the diversity of different types of pasture (e.g. loss of fen meadows, forested pasture) [36]. There has been a decrease in the area (number/length) of **small scale habitats** on farmland which serve as important habitats for wild species and can provide benefits in terms of the visual landscape. In particular, the loss of open ditches (due to the expansion of sub-surface drainage), small woodland patches within fields, and field boundaries (due to larger field sizes), have had adverse impacts on biodiversity and landscape [42, 43]. However, farmers have been given incentives since 1995 to maintain and

develop buffer strips in agricultural landscapes [43, 44, 45]. It is estimated that around 60-90 000 hectares of **peatland** are currently in agricultural use, or about 4% of the farmland area [46, 47], with Finland one of the world's richest countries in terms of peatlands [48].

Cultural agricultural landscapes and agri-biodiversity have in general been adversely affected by a lack of grazing of extensive pastures [1, 33, 49]. Research has shown that there are positive effects from low intensity grazing for many plant species in semi-natural pastures in Finland compared to either discontinuing grazing or subjecting these habitats to high intensity stocking levels [50, 51]. For some plant species, including certain rare species, other factors were also important for their survival, such as mowing rather than grazing pasture, the timing of grazing, and the type and breed of grazing livestock [50, 52, 53]. But the increase in tree cover where grazing is discontinued can also be detrimental to plant species rather than the lack of grazing *per se* [51]. However, changes in the grazing system or its abandonment seem less important for bird than plant species [49]. For butterflies and moths where grazing was restored on abandoned pasture, there was little evidence of the colonisation of old pasture species in restored sites. Moreover, species richness and abundance of butterflies and moths was found to be greatest in abandoned pastures [54], and higher with low intensity compared to high intensity grazed pasture [55].

3.7.3. Overall agri-environmental performance

Despite the significant reduction in pollution from agriculture over the period since 1990, this has not been yet reflected in an overall improvement in the state of environment [1]. From 1990 to 2004 there were substantial reductions in nutrient surpluses and ammonia and GHG emissions, in most cases well below the changes for the EU15 and OECD average. While this has helped ease pressure on the environment, the quality of water in rivers, lakes and the Baltic Sea has not improved, but it must be noted, however, that Finland is only one of a number of countries polluting the Baltic. There has been an increase in pesticide use since the mid-1990s, but because of a lack of monitoring the potential impact of greater pesticide loadings on the environment is unclear, although the intensity of pesticide use remains low compared to many OECD countries. There has been a deterioration in the quality and quantity of wild species, and the habitats important to them, in agricultural landscapes, notably semi-natural grassland.

Environmental monitoring has a long history in Finland but tracking agri-environmental performance is more recent [56]. Indicators are used to evaluate the effectiveness of the *Strategy for the Sustainable Use of Renewable Natural Resources*, which encompasses agriculture [57, 58, 59].

Since joining the EU agri-environmental policies have been strengthened. A new agri-environmental support scheme for the period 2007-13 was approved by the EU Commission in June 2007 as part of the 2007-13 *Rural Development Strategy and Programme*. **Changes in farming practices** have brought various environmental benefits, such as low tillage helping to reduce nutrient pollution through lowering rates of soil erosion, and improvements in manure management leading to reduced nutrient loadings, ammonia and greenhouse gas emissions. Improved manure management practices has been a factor raising the efficiency of nutrient use, nevertheless, phosphorus use efficiency remains lower than that of many OECD countries. Deterioration of **biodiversity** in farming environments has continued, although there does appear to be some recent success in


increasing the area of extensive semi-natural grasslands, towards the goal of 60 000 ha by 2010 [1]. But the trend toward greater sub-surface drainage has led to the removal of ditches, which has been harmful for biodiversity dependent on small scale habitats.

While further improvements in agri-environmental performance are likely some environmental concerns remain. Projections of the agricultural sector to 2013 suggest a further contraction of sector, which is expected to lead to a further concentration of production on fewer and larger farms, while productivity will increase [60]. Some researchers consider there could be trade-offs between fewer but more concentrated and intensive agricultural production and environmental quality, especially stemming from the loss of semi-natural grasslands to non-agricultural uses [61, 62, 63]. Hence, a major challenge facing Finnish policy makers is the conservation of semi-natural grazed grasslands which are recognised as providing biodiversity and cultural landscape benefits.

The mid-term assessment of the Water Protection Targets concluded that even with the implementation of agri-environmental measures the nutrient reduction targets for 2005 could not be reached, despite progress in lowering agricultural nutrient surpluses [7, 25]. As a result of the 2005 target not being reached a new reduction target to 2015 was approved in 2007, seeking to reduce nitrogen and phosphorus loads from agriculture by 30% from 2002-05 levels. Moreover, the nutrients now stored in water bodies will continue to deteriorate water quality for many years, suggesting that action may be required for their recovery [1, 23], especially if Finland is to meet its international commitment to reduce nutrient loadings into the Baltic Sea. Exemptions for farmers from **energy and climate change taxes** act as disincentives to further limit on-farm energy consumption, improve energy efficiency and reduce GHG emissions [18]. Given the continuing deterioration of **biodiversity** for both flora and fauna [1], some researchers have noted that the uptake of biodiversity conservation by farmers under agri-environmental schemes has been low and the share of government agri-environmental expenditure (12%, between 2000 and 2003) is too low to adequately improve agri-biodiversity [64]. While on the one hand peat production and use of peatlands can have harmful impacts on biodiversity, GHG emissions and water systems, they can on the other hand provide economic and social benefits. The agricultural sector makes a positive contribution to meeting international environment agreement commitments to reduce *ammonia and greenhouse gas emissions*. The contribution of farming in cutting GHG emissions might be further enhanced with the cessation of the agricultural use of peatlands, which potentially could decrease agricultural sector GHG emissions by up to 10% [46].

Figure 3.7.2. **National agri-environmental performance compared to the OECD average**

Percentage change 1990-92 to 2002-04¹



Absolute and economy-wide change/level


Variable	Unit	Finland	OECD
Agricultural production volume	Index (1999-01 = 100) 1990-92 to 2002-04	100	105
Agricultural land area	000 hectares 1990-92 to 2002-04	-298	-48 901
Agricultural nitrogen (N) balance	Kg N/hectare 2002-04	55	74
Agricultural phosphorus (P) balance	Kg P/hectare 2002-04	8	10
Agricultural pesticide use	Tonnes 1990-92 to 2001-03	-157	-46 762
Direct on-farm energy consumption	000 tonnes of oil equivalent 1990-92 to 2002-04	-104	+1 997
Agricultural water use	Million m ³ 1990-92 to 2001-03	n.a.	+8 102
Irrigation water application rates	Megalitres/ha of irrigated land 2001-03	n.a.	8.4
Agricultural ammonia emissions	000 tonnes 1990-92 to 2001-03	-5	+115
Agricultural greenhouse gas emissions	000 tonnes CO ₂ equivalent 1990-92 to 2002-04	-922	-30 462

n.a.: Data not available. Zero equals value between -0.5% to < +0.5%.

1. For agricultural water use, pesticide use, irrigation water application rates, and agricultural ammonia emissions the % change is over the period 1990-92 to 2001-03.
2. Percentage change in nitrogen and phosphorus balances in tonnes.

Source: OECD Secretariat. For full details of these indicators, see Chapter 1 of the *Main Report*.


Figure 3.7.3. **Nitrogen fluxes in the Paimionjoki river¹ and agricultural nitrogen balances**



1. The Paimionjoki river is situated in the main agricultural area of Finland.

Source: Ministry of Agriculture and Forestry, Finland.

Figure 3.7.4. **Population trends of Finnish farmland butterflies in three ecological species groups**



Source: Heliola, J., M. Kuussaari and I. Niininen (2007), "Results of the butterfly monitoring scheme in Finnish agricultural landscapes for the year 2005", *Baptria*, Vol. 32 (in press).

StatLink <http://dx.doi.org/10.1787/300102623372>

Bibliography

- [1] Agrifood Research Finland (2006), *Finnish Agriculture and Rural Industries 2006*, J. Niemi and J. Ahstedt (eds.), Economics Research Publications 106a, Helsinki, Finland, www.mtt.fi/english/publications/mtt_dp.html.
- [2] Kuosmanen, T. and T. Sipiläinen (2004), *On the anatomy of productivity growth: A decomposition of the Fisher Ideal TFP index*, Discussion Paper No. 2004:17, MTT Economic Research, Agrifood Research Finland, Helsinki, Finland, www.mtt.fi/english/publications/mtt_dp.html.
- [3] Agrifood Research Finland (2005), *Finnish Agriculture and Rural Industries 2005 – Ten Years in the European Union*, Niemi, J and J. Ahstedt (ed.), Economics Research Publications 105a, Helsinki, Finland, www.mtt.fi/english/publications/mtt_dp.html.
- [4] Niemi, J. (2005), “The static welfare effects of the accession to the European Union on the Finnish agricultural markets”, *Agricultural and Food Science*, Vol. 14, pp. 224-235.
- [5] Information Centre of the Ministry of Agriculture and Forestry, *2005 Yearbook of Farm Statistics*, Helsinki, www.mmmmtike.fi/en/index/statistics_information_services/publications.html.
- [6] The Finnish response to the OECD Agri-environmental Indicator Questionnaire, unpublished.
- [7] Nikkola, E. and K. Tolonen (2006), “Tool for Monitoring and Evaluating the On-farm Environment Management and Nutrient Use on Finnish Cattle Farms”, in OECD, *Agriculture and Water: Sustainability, Markets and Policies*, Paris, France, www.oecd.org/tad/env.
- [8] OECD (1995), *Agricultural Policies, Markets and Trade in OECD Countries: Monitoring and Outlook 1995*, Paris, France, www.oecd.org/agr/policy.
- [9] Statistics Finland (2006), *Finland's Natural Resources and the Environment 2005*, Helsinki, Finland, www.stat.fi/tk/tt/ymparisto_en.html.
- [10] Kröger, L., J. Lankoski and A. Huhtala (2005), *Review of the literature on Agri-environmental schemes in Finland*, report to the EU Commission Sixth Framework Programme on Integrated tools to design and implement agro environmental schemes, web-based report, <http://merlin.lusignan.inra.fr/ITAES/website/Publicdeliverables>.
- [11] Kaljonen, M. (2006), “Co-construction of agency and environmental management. The case of agri-environmental policy implementation at Finnish farms”, *Journal of Rural Studies*, Vol. 22, pp. 205-216.
- [12] Yliskylä-Peuralahti, J. (2003), “Biodiversity – a new spatial challenge for Finnish agri-environmental policies?”, *Journal of Rural Studies*, Vol. 19, pp. 215-231.
- [13] Kohonen, J.T. (2003), “Finnish strategies for reduction and control of effluents to the marine environment – Examples from agriculture, municipalities and industry”, *Marine Pollution Bulletin*, Vol. 47, pp. 162-168.
- [14] Kröger, L. (2005), “Development of the Finnish agri-environmental policy as a learning process”, *European Environment*, Vol. 15, pp. 13-26.
- [15] Hiltunen, M. (2004), *Economic environmental policy instruments in Finland*, Finnish Environment Institute, Helsinki, Finland, www.environment.fi/default.asp?node=6284&lan=EN.
- [16] OECD (2005), *Taxation and social security in agriculture*, Paris, France, www.oecd.org/agr.
- [17] Åkerman, M., M. Kaljonen and T. Peltola (2005), “Integrating environmental policies into local practices: The politics of agri-environmental and energy policies in rural Finland”, *Local Environment*, Vol. 10, No. 6, pp. 595-611.
- [18] Vehmas, J. (2005), “Energy-related taxation as an environmental policy tool – the Finnish experience 1990-2003”, *Energy Policy*, Vol. 33, pp. 2175-2182.
- [19] Tattari, S. and S. Rekolainen (2006), “Soil Erosion in Finland”, in J. Boardman and J. Poesen (eds.), *Soil Erosion in Europe*, John Wiley, London, United Kingdom.
- [20] Puustinen, M., S. Tattari, J. Koskiaho and J. Linjama (2007), “Influence of seasonal and annual hydrological variations on erosion and phosphorus transport from arable areas in Finland”, *Soil and Tillage Research*, Vol. 93, Issue 1, pp. 44-55.
- [21] Lankoski, J., M. Ollikainen and P. Uusitalo (2006), “No-till technology: benefits to farmers and the environment? Theoretical analysis and application to Finnish agriculture”, *European Review of Agricultural Economics*, Vol. 33, No. 2, pp. 193-221.
- [22] Lepistö, A., K. Granlund, P. Kortelainen and A. Räike (2006), “Nitrogen in river basins: Sources, retention in the surface waters and peatlands, and fluxes to estuaries in Finland”, *Science of the Total Environment*, Vol. 365, pp. 238-259.

- [23] Ekholm, P. and S. Mitikka (2006), "Agricultural lakes in Finland: Current water quality and trends", *Environmental Monitoring and Assessment*, Vol. 116, pp. 111-135.
- [24] Laukkanen, M. and A. Huhtala (2006), *Optimal management of a eutrophied coastal ecosystem: Balancing agricultural and municipal abatement measures*, Agrifood Research Finland Discussion Paper No. 4, Helsinki, Finland, www.mtt.fi/english/publications/mtt_dp.html.
- [25] Kohonen, J.T. (2003), "Finnish strategies for reduction and control of effluents to the marine environment – examples from agriculture, municipalities and industry", *Marine Pollution Bulletin*, Vol. 47, pp. 162-168.
- [26] Salo, T. and E. Turtola (2006), "Nitrogen balance as an indicator of nitrogen leaching in Finland", *Agriculture, Ecosystems and Environment*, Vol. 113, pp. 98-107.
- [27] Virtanen, H. (2005), "Nitrogen and phosphorus balances on Finnish dairy farms", *Agricultural and Food Science*, Vol. 14, pp. 166-180.
- [28] Granlund, K., A. Räike, P. Ekholm, K. Rankinen and S. Rekolainen (2005), "Assessment of water protection targets for agricultural nutrient loading in Finland", *Journal of Hydrology*, Vol. 304, pp. 251-260.
- [29] Statistics Finland (2005), *Finland's Fourth National Communication under the United Nations Framework Convention on Climate Change*, see the UNFCCC website at http://unfccc.int/national_reports/annex_i_natcom/submitted_natcom/items/3625.php.
- [30] Lankoski, J. and M. Ollikainen (2006), *Bioenergy crop production and climate policies: A von Thunen model and case of reed canary grass in Finland*, unpublished manuscript, Department of Economics and Management, Discussion Paper, No. 17, University of Helsinki, Finland, www.tiedekirjasto.helsinki.fi:8080/dspace/browse-date?top=1975%2F718.
- [31] Kuussaari, M., J. Heliölä and M. Luoto (2004), "Farmland biodiversity indicators and monitoring in Finland", in G. Gross (ed.), *Developments in Strategic Landscape Monitoring for the Nordic Countries*, Nordic Council of Ministers, Diverse Series Report, Nordic Council, Copenhagen, Denmark, www.ymparisto.fi/default.asp?contentid=198676&lan=en.
- [32] Kuussaari, M. and J. Heliölä (2003), "National and regional level farmland biodiversity indicators in Finland", in OECD, *Agriculture and Biodiversity: Developing Indicators for Policy Analysis*, Paris, France, www.oecd.org/tad/env/indicators.
- [33] Pitkänen, M. and J. Tiainen (eds.) (2001), *Biodiversity of agricultural landscapes in Finland*, Birdlife Finland Conservation Series No. 3, Birdlife, Suomi, Finland, www.birdlife.fi/julkaisut/julkaisusarja/index.shtml.
- [34] Ministry of the Environment (2005), *Third National Report of Finland to the Convention on Biological Diversity*, Secretariat to the Convention on Biological Diversity, Montreal, Canada, www.biodiv.org/reports/list.aspx?menu=chm.
- [35] Agrifood Research Finland (2006), *Genetic Resources – Securing Biodiversity*, Helsinki, Finland, www.mtt.fi/english/research/plants/genetics.pdf.
- [36] Kivinen, S. (2005), "Regional distribution and biodiversity perspectives of Finnish grasslands", *Fennia*, Vol. 183, Part 1, pp. 37-56.
- [37] Rassi, P., A. Alanen, T. Kanerva and I. Mannerkoski (eds.) (2001), *The 2000 Red List of Finnish Species*, in Finnish with English summary, Finnish Environment Institute, Helsinki, Finland, www.ymparisto.fi/default.asp?contentid=179629&lan=en.
- [38] Luoto, M., S. Rekolainen, J. Aakkula and J. Pykälä (2003), "Loss of plant species richness and habitat connectivity in grasslands associated with agricultural change in Finland", *Ambio*, Vol. 32, No. 7, pp. 447-452.
- [39] Virkkala, R., M. Luoto and K. Rainio (2004), "Effects of landscape composition on farmland and red-listed birds in boreal agricultural-forest mosaics", *Ecography*, Vol. 27, pp. 273-284.
- [40] Finnish Environment Institute, *Monitoring of Butterflies in Finnish Agricultural Landscapes*, website database on trends in butterfly and moth species, www.environment.fi/default.asp?node=17966&lan=en.
- [41] Miettinen, A. and A. Huhtala (2005), *Modelling environmental effects of agriculture: The case of organic rye and grey partridge*, paper presented to the European Association of Agricultural Economists, 24-27 August, Copenhagen, Denmark, <http://agecon.lib.umn.edu/cgi-bin/view.pl>.
- [42] Hietala-Koivu, R., J. Lankoski and S. Tarmi (2004), "Loss of biodiversity and its social cost in an agricultural landscape", *Agriculture, Ecosystems and Environment*, Vol. 103, pp. 75-83.
- [43] Hietala-Koivu, R. (2002), "Landscape and modernizing agriculture: a case study of three areas in Finland in 1954-1998", *Agriculture, Ecosystems and Environment*, Vol. 91, pp. 273-281.

- [44] Lankoski, J. (2005), "Alternative Approaches for Evaluating the Performance of Buffer Strip Policy in Finland", in OECD, *Evaluating Agri-environmental Policies: Design, Practice and Results*, Paris, France, www.oecd.org/tad/env.
- [45] Ma, M., S. Tarmi and J. Helenius (2002), "Revisiting the species-area relationship in a semi-natural habitat: floral richness in agricultural buffer zones in Finland", *Agriculture, Ecosystems and Environment*, Vol. 89, pp. 137-148.
- [46] Lehtonen, H., J. Peltola and M. Sinkkonen (2006), "Co-effects of climate policy and agricultural policy on regional agricultural viability in Finland", *Agricultural Systems*, Vol. 88, pp. 472-493.
- [47] Vasander, H. (2006), "The use of mires for agriculture and forestry", in Lindholm, T. and R. Heikkilä, *Finland land of mires*, pp. 173-178, Finnish Environment Institute, Helsinki, Finland, www.ymparisto.fi/default.asp?contentid=194173&lan=en.
- [48] Takko, A. and H. Vasander (2004), "Socioeconomic aspects of commercial uses of peatlands in Finland", *Proceedings of the 12th International Peat Conference*, Vol. 2, pp. 1313-1322.
- [49] Luoto, M. and J. Pykälä (2003), "Decline of landscape-scale habitat and species diversity after the end of cattle grazing", *Journal of Nature Conservation*, Vol. 11, pp. 171-178.
- [50] Pykälä, J. (2005), "Plant species responses to cattle grazing in mesic semi-natural grassland", *Agriculture, Ecosystems and Environment*, Vol. 108, pp. 109-117.
- [51] Pykälä, J., M. Luoto, R.K. Heikkinen and T. Kontula (2005), "Plant species richness and persistence of rare plants in abandoned semi-natural grasslands in northern Europe", *Basic and Applied Ecology*, Vol. 6, pp. 25-33.
- [52] Pykälä, J. (2004), "Cattle grazing increases plant species richness of most species trait groups in mesic semi-natural grasslands", *Plant Ecology*, Vol. 175, pp. 217-226.
- [53] Pykälä, J. (2003), "Effects of restoration with cattle grazing on plant species composition and richness of semi-natural grasslands", *Biodiversity and Conservation*, Vol. 12, pp. 2211-2226.
- [54] Pöyry, J., S. Lindgren, J. Salminen and M. Kuussaari (2005), "Responses of butterfly and moth species to restored cattle grazing in semi-natural grasslands", *Biological Conservation*, Vol. 122, pp. 465-478.
- [55] Pöyry, J., S. Lindgren, J. Salminen and M. Kuussaari (2004), "Restoration of butterfly and moth communities in semi-natural grasslands by cattle grazing", *Ecological Applications*, Vol. 14, No. 6, pp. 1656-1670.
- [56] Niemi, J. (ed.) (2006), *Environmental Monitoring in Finland 2006-2008*, Finnish Environment Institute, Helsinki, Finland, www.environment.fi/default.asp?node=19251&lan=en.
- [57] Ministry of Agriculture and Forestry (2004), *Indicators for renewable natural resources*, Helsinki, Finland, www.mmm.fi/en/index/frontpage.html.
- [58] Yli-Viikari, A. and R. Lemola (2005), "Usability of management indicators – Considerations from a Finnish perspective", in OECD, *Farm Management and the Environment: Developing Indicators for Policy Analysis*, Paris, France, www.oecd.org/tad/env/indicators.
- [59] Yli-Viikari, A., H. Risku-Norja, V. Nuutinen, E. Heinonen, R. Hietala-koivu, E. Huusela-veistola, T. Hyvönen, J. Kantanen, S. Raussi, P. Rikkinen, A. Seppälä and E. Vehmasto (2002), *Agri-environmental and rural development indicators – A proposal*, Agrifood Research Reports 5, Agrifood Research Finland, Helsinki, Finland, www.mtt.fi/english/publications/mtt_dp.html.
- [60] Lehtonen, H. and P. Pyykkönen (2005), *Structural development of Finnish agriculture until 2013*, English summary only, Agrifood Research Working Papers No. 100, Agrifood Research Finland, Helsinki, Finland, www.mtt.fi/english/press/pressrelease.html.
- [61] Lehtonen, H., J. Aakkula and P. Rikkinen (2005), "Alternative agricultural policy scenarios, sector modelling and indicators: A sustainability assessment", *Journal of Sustainable Agriculture*, Vol. 26, No. 4, pp. 63-93.
- [62] Rikkinen, P. (2005), "Scenarios for future agriculture in Finland: a Delphi study among agri-food sector stakeholders", *Agricultural and Food Science*, Vol. 14, pp. 205-223.
- [63] Lehtonen, H., M. Kuussaari, T. Hyvönen and J. Lankoski (2006), *Performance of alternative policy measures to increase biodiversity value of farmlands in Finland*, unpublished paper, Agrifood Research Finland, Helsinki, Finland.
- [64] Herzon, I. and M. Mikk (2007), "Farmers' perceptions of biodiversity and their willingness to enhance it through agri-environment schemes: A comparative study from Estonia and Finland", *Journal of Nature Conservation*, Vol. 15, pp. 10-25.

Table of Contents

I. Highlights	15
Overall agri-environmental performance.	15
Agri-environmental performance in specific areas	16
Caveats and limitations	19
Matching indicator criteria	20
II. Background and Scope of the Report.	23
1. Objectives and scope.	23
2. Data and information sources.	24
3. Progress made since the OECD 2001 Agri-environmental Indicator Report	25
4. Structure of the Report	26
Bibliography	28
Annex II.A1. List of indicators in Chapter 1	29
Annex II.A2. Indicators in Chapter 1 assessed according to the OECD indicator criteria	31
Chapter 1. OECD Trends of Environmental Conditions related to Agriculture since 1990	37
1.1. Agricultural production and land	38
1.1.1. Introduction	39
1.1.2. Agricultural production	39
1.1.3. Agricultural land use	40
1.1.4. Linkages between agricultural production and land use	46
Bibliography	47
1.2. Nutrients	48
1.2.1. Nitrogen balance	52
1.2.2. Phosphorus balance	56
1.2.3. Regional (sub-national) nutrient balances	60
Bibliography	62
1.3. Pesticides	63
1.3.1. Pesticide use	63
1.3.2. Pesticide risk indicators	67
Bibliography	74
1.4. Energy	76
Bibliography	83
1.5. Soil	84
Bibliography	90

1.6. Water.....	92
1.6.1. Water use	93
1.6.2. Water quality	100
Bibliography	108
1.7. Air	109
Background	110
1.7.1. Ammonia emissions, acidification and eutrophication.....	110
1.7.2. Methyl bromide use and ozone depletion	117
1.7.3. Greenhouse gas emissions and climate change	122
Bibliography	130
1.8. Biodiversity	133
Background	134
1.8.1. Genetic diversity	136
1.8.2. Wild species diversity	146
1.8.3. Ecosystem diversity.....	148
Bibliography	159
1.9. Farm management	160
1.9.1. Overview of environmental farm management	163
1.9.2. Nutrient management	163
1.9.3. Pest management	168
1.9.4. Soil management.....	169
1.9.5. Water management.....	172
1.9.6. Biodiversity management	173
1.9.7. Organic management	174
Bibliography	176
Chapter 2. OECD Progress in Developing Agri-environmental Indicators	179
2.1. Introduction.....	180
2.2. Progress in developing OECD Agri-environmental Indicators	180
2.2.1. Soil: Erosion, biodiversity and soil organic carbon	180
2.2.2. Water: Use and water quality	184
2.2.3. Biodiversity: Genetic, wild species and ecosystem diversity	188
2.2.4. Land: Landscapes and ecosystem functions	192
2.2.5. Farm management	195
2.3. Overall assessment.....	196
Annex 2.A1. Agri-environmental Indicators of Regional Importance and/or under Development.....	200
Annex 2.A2. A Qualitative Assessment of the Agri-environmental Indicators included in Annex 2.A1 according to the OECD Indicator Criteria	202
Bibliography	207
Chapter 3. OECD Country Trends of Environmental Conditions related to Agriculture since 1990	209
Background to the country sections	210
3.1. Australia	212
3.2. Austria	224
3.3. Belgium.....	234
3.4. Canada	243

3.5. Czech Republic	256
3.6. Denmark.....	269
3.7. Finland	284
3.8. France	296
3.9. Germany	305
3.10. Greece.....	313
3.11. Hungary	324
3.12. Iceland	336
3.13. Ireland.....	344
3.14. Italy	357
3.15. Japan.....	366
3.16. Korea.....	377
3.17. Luxembourg.....	386
3.18. Mexico.....	393
3.19. Netherlands	402
3.20. New Zealand	413
3.21. Norway	423
3.22. Poland.....	433
3.23. Portugal.....	448
3.24. Slovak Republic	459
3.25. Spain.....	472
3.26. Sweden.....	486
3.27. Switzerland	498
3.28. Turkey.....	507
3.29. United Kingdom	522
3.30. United States	532
3.31. European Union.....	545
Chapter 4. Using Agri-environmental Indicators for Policy Analysis	551
4.1. Policy context to OECD agri-environmental performance	552
4.2. Tracking agri-environmental performance.....	554
4.2.1. Evolution of Agri-environmental Indicators to track sustainable development.....	554
4.2.2. Tracking national agri-environmental performance	556
4.2.3. International reporting on environmental conditions in agriculture	559
4.2.4. Non-governmental organisations (NGOs)	561
4.3. Using Agri-environmental Indicators for policy analysis	562
4.3.1. OECD member countries	563
4.3.2. International governmental organisations	565
4.3.3. Research community	567
4.4. Knowledge gaps in using Agri-environmental Indicators.....	568
Bibliography	571
 List of boxes	
II.1. OECD Expert Meetings on Agri-environmental Indicators: 2001-04	25
1.7.1. Towards a net agricultural greenhouse gas balance indicator?.....	123

1.8.1. Defining agricultural biodiversity	134
2.1. Soil biodiversity in agricultural land	182
2.2. Agricultural livestock pathogens and water pollution	187
2.3. The impact of agriculture on aquatic ecosystems.	188
4.1. Main agri-environmental measures in OECD countries.	553
4.2. Selected international and regional environmental agreements relevant to agriculture.	555

List of tables

1.1.1. OECD and world agricultural production	39
1.1.2. OECD and world agricultural exports	40
1.3.1. Germany: Percentage risk indices	70
1.7.1. Total OECD emissions of acidifying pollutants	114
1.7.2. Ammonia emission targets to 2010 under the Convention on Long-range Transboundary Air Pollution.	116
1.7.3. Methyl bromide use and progress in meeting the phase-out schedule under the <i>Montreal Protocol</i>	120
1.7.4. Critical Use Exemptions (CUEs) for methyl bromide agreed under the <i>Montreal Protocol</i> for 2005.	121
1.7.5. Total OECD gross greenhouse gas emissions	124
1.7.6. Main sources and types of gross greenhouse gas emissions	127
1.8.1. Area of transgenic crops for major producing countries	139
1.8.2. Plant genetic resource conservation activities for OECD countries	139
1.8.3. Livestock genetic resource conservation activities for OECD countries.	144
1.8.4. Share of farm woodland in agricultural land area.	157
1.8.5. Share of farm fallow in agricultural land area	157
1.9.1. Countries recording adoption of environmental farm management practices	164
1.9.2. Overview of farmer incentives to adopt environmental farm management practices	166
2.1. Net water balance in a Japanese rice field irrigation system: 2003.	185

List of figures

II.1. The Driving Force-State-Response framework: Coverage of indicators	24
1.1.1. Production, yields and area harvested and future projections for selected commodities and OECD countries	41
1.1.2. Volume of total agricultural production	43
1.1.3. Share of agricultural land use in the national land area	44
1.1.4. Agricultural land area	45
1.1.5. Agricultural production volume index and agricultural land area	46
1.2.1. Main elements in the OECD gross nutrient (nitrogen and phosphorus) balance calculation	50
1.2.2. Gross nitrogen balance estimates	51
1.2.3. Gross nitrogen balances for selected OECD countries	53
1.2.4. Inorganic nitrogen fertilisers and livestock manure nitrogen input in nitrogen balances.	54

1.2.5. Agricultural use of inorganic nitrogen and phosphate fertilisers	54
1.2.6. Contribution of the main sources of nitrogen inputs and outputs in nitrogen balances	56
1.2.7. Nitrogen efficiency based on gross nitrogen balances	57
1.2.8. Gross phosphorus balance estimates	58
1.2.9. Gross phosphorus balance for selected OECD countries	59
1.2.10. Contribution of the main sources of phosphorus inputs and outputs in phosphorus balances	60
1.2.11. Phosphorus efficiency based on phosphorus balances	61
1.2.12. Spatial distribution of nitrogen balances in Canada and Poland	62
1.3.1. Pesticide use in agriculture	65
1.3.2. Pesticide use for selected OECD countries	66
1.3.3. Belgium: Risk for aquatic species due to use of pesticides in arable land, horticulture and outside of agriculture	69
1.3.4. Denmark: The annual trend in frequency of pesticide application	70
1.3.5. The Netherlands: Potential chronic effects scores for aquatic and terrestrial organisms and leaching into groundwater	71
1.3.6. Norway: Trends of health risk, environmental risk and sales of pesticides	72
1.3.7. Sweden: National level pesticide risk indicators and the number of hectare doses	73
1.3.8. United Kingdom (England and Wales): Total area of pesticide applications	74
1.4.1. Simplified energy “model” of an agricultural system	78
1.4.2. Direct on-farm energy consumption	79
1.4.3. Direct on-farm energy consumption for selected OECD countries	80
1.4.4. Agricultural employment and farm machinery use	81
1.4.5. Composition of on-farm energy consumption in the EU15 and the United States	82
1.5.1. Agricultural land area classified as having moderate to severe water erosion risk	87
1.5.2. Trends in agricultural land area classified as having moderate to severe water erosion risk	88
1.5.3. Agricultural land area classified as having moderate to severe wind erosion risk	89
1.6.1. Agricultural water use	95
1.6.2. Share of national water use in annual freshwater resources and share of agricultural water use in national use	96
1.6.3. Irrigated area, irrigation water use and irrigation water application rates	97
1.6.4. Share of agricultural groundwater use in total groundwater use, and total groundwater use in total water use	99
1.6.5. Share of agriculture in total emissions of nitrates and phosphorus in surface water	102
1.6.6. Share of agriculture in total emissions of nitrates and phosphorus in coastal water	103
1.6.7. Share of monitoring sites in agricultural areas exceeding national drinking water limits for nitrates and phosphorus in surface water	104
1.6.8. Share of monitoring sites in agricultural areas exceeding national drinking water limits for nitrates in groundwater	105

1.6.9. Share of monitoring sites in agricultural areas where one or more pesticides are present in surface and groundwater	106
1.6.10. Share of monitoring sites in agricultural areas exceeding national drinking water limits for pesticides in surface water and groundwater	107
1.7.1. Impacts of agriculture on air quality: Multi-pollutants, multi-effects	110
1.7.2. Ammonia emissions from agriculture	112
1.7.3. Emissions of acidifying airborne pollutants for the EU15, US and OECD.	113
1.7.4. Agricultural ammonia emission trends for selected OECD countries	114
1.7.5. Share of the main sources of agricultural ammonia emissions in OECD countries	117
1.7.6. Methyl bromide use	119
1.7.7. Global methyl bromide use by major sectors.	121
1.7.8. Agricultural gross greenhouse gas emissions	125
1.7.9. Gross agricultural greenhouse gas emissions in carbon dioxide equivalent for selected OECD countries	126
1.7.10. Agricultural production and agricultural greenhouse gas emissions.	128
1.7.11. Main sources of methane and nitrous oxide emissions in OECD agriculture	129
1.7.12. Contribution of main sources in agricultural greenhouse gas emissions	130
1.8.1. OECD agri-biodiversity indicators framework	135
1.8.2. Change in the number of plant varieties registered and certified for marketing	137
1.8.3. Change in the share of the one-to-five dominant crop varieties in total marketed crop production	138
1.8.4. Change in the number of livestock breeds registered and certified for marketing	141
1.8.5. Change in the share of the three major livestock breeds in total livestock numbers.	142
1.8.6. Total number of cattle, pigs, poultry and sheep in endangered and critical risk status and under conservation programmes	143
1.8.7. Share of selected wild species that use agricultural land as primary habitat.	148
1.8.8. Population trends of farmland birds	149
1.8.9. Change in agricultural land use and other uses of land.	152
1.8.10. Permanent pasture and arable and permanent cropland	155
1.8.11. Share of arable and permanent cropland, permanent pasture and other agricultural land in total agricultural land area.	156
1.8.12. Share of national Important Bird Areas where intensive agricultural practices pose a serious threat or a high impact on the areas' ecological functions	158
1.9.1. OECD farm management indicator framework	162
1.9.2. Share of agricultural land area under nutrient management plans.	168
1.9.3. Share of total number of farms under nutrient management plans	169
1.9.4. Share of total number of farms using soil nutrient testing	170
1.9.5. Share of total arable and permanent crop area under integrated pest management.	171
1.9.6. Share of arable crop area under soil conservation practices	172
1.9.7. Share of total arable and permanent crop area under all-year vegetative cover	173
1.9.8. Share of irrigated land area using different irrigation technology systems	174

1.9.9. Share of agricultural land area under biodiversity management plans	175
1.9.10. Share of agricultural land area under certified organic farm management	176
2.1. Canadian soil organic carbon stocks in agricultural soils by different classes	183
2.2. United States soil organic carbon stocks in agricultural soils by different classes	184
2.3. Agricultural, industrial, and household water charges	186
2.4. National crop varieties that are endangered	189
2.5. National crop varieties that are not at risk.	190
2.6. Edge density of agricultural fields in Finland.	190
2.7. Share of Canadian farmland in various classes of the habitat capacity index.	191
2.8. Cultural landscape features on agricultural land	193
2.9. Water retaining capacity of agriculture	194
2.10. Water retaining capacity for agricultural facilities	195
2.11. Share of farmers participating in agri-environmental education programmes	197
3.1.1. National agri-environmental and economic profile, 2002-04: Australia	212
3.1.2. National agri-environmental performance compared to the OECD average.	220
3.1.3. National Landcare membership.	220
3.1.4. Annual quantities of insecticide and acaricide applied to the cotton crop	220
3.2.1. National agri-environmental and economic profile, 2002-04: Austria	224
3.2.2. National agri-environmental performance compared to the OECD average.	231
3.2.3. Area under non-use of inputs, organic farming and erosion control measures of the ÖPUL agri-environmental programme.	231
3.2.4. Greenhouse gas emissions from agriculture	231
3.3.1. National agri-environmental and economic profile, 2002-04: Belgium	234
3.3.2. National agri-environmental performance compared to the OECD average.	240
3.3.3. Total pesticide use	240
3.3.4. Greenhouse gas emissions and sinks	240
3.4.1. National agri-environmental and economic profile, 2002-04: Canada	243
3.4.2. National agri-environmental performance compared to the OECD average.	252
3.4.3. Share of cropland in different soil organic carbon change classes.	252
3.4.4. Share of farmland in different wildlife habitat capacity change classes.	252
3.5.1. National agri-environmental and economic profile, 2002-04: Czech Republic	256
3.5.2. National agri-environmental performance compared to the OECD average.	265
3.5.3. Share of samples above Czech drinking water standards for nitrates in surface water	265
3.5.4. Monitored numbers of partridge population	265
3.6.1. National agri-environmental and economic profile, 2002-04: Denmark	269
3.6.2. National agri-environmental performance compared to the OECD average.	280
3.6.3. Share of monitoring sites with occurrences of pesticides in groundwater used for drinking	280
3.6.4. Share of meadows and dry grasslands, heath, and bogs and marshes in the total land area	280
3.7.1. National agri-environmental and economic profile, 2002-04: Finland	284
3.7.2. National agri-environmental performance compared to the OECD average.	292
3.7.3. Nitrogen fluxes in the Paimionjoki river and agricultural nitrogen balances	292

3.7.4. Population trends of Finnish farmland butterflies in three ecological species groups.	292
3.8.1. National agri-environmental and economic profile, 2002-04: France.	296
3.8.2. National agri-environmental performance compared to the OECD average.	302
3.8.3. Trends in key agri-environmental indicators.	302
3.8.4. Trends in key agri-environmental indicators.	302
3.9.1. National agri-environmental and economic profile, 2002-04: Germany	305
3.9.2. National agri-environmental performance compared to the OECD average.	310
3.9.3. Share of the number of farms and Utilised Agricultural Area (UAA) under organic farming.	310
3.9.4. Share of renewable biomass and energy crop area in the total agricultural land area	310
3.10.1. National agri-environmental and economic profile, 2002-04: Greece	313
3.10.2. National agri-environmental performance compared to the OECD average.	321
3.10.3. Irrigated area and irrigation water application rates	321
3.10.4. <i>Ex situ</i> accessions of plant landraces, wild and weedy relatives.	321
3.11.1. National agri-environmental and economic profile, 2002-04: Hungary.	324
3.11.2. National agri-environmental performance compared to the OECD average.	333
3.11.3. Agricultural land affected by various classes of water erosion	333
3.11.4. Support payments for agri-environmental schemes and the number of paid applications.	333
3.12.1. National agri-environmental and economic profile, 2002-04: Iceland	336
3.12.2. National agri-environmental performance compared to the OECD average.	342
3.12.3. Annual afforestation	342
3.12.4. Annual area of wetland restoration.	342
3.13.1. National agri-environmental and economic profile, 2002-04: Ireland	344
3.13.2. National agri-environmental performance compared to the OECD average.	353
3.13.3. River water quality	353
3.13.4. Population changes for key farmland bird populations	353
3.14.1. National agri-environmental and economic profile, 2002-04: Italy.	357
3.14.2. National agri-environmental performance compared to the OECD average.	363
3.14.3. Actual soil water erosion risk.	363
3.14.4. Regional change in agricultural land area: 1990 to 2000.	363
3.15.1. National agri-environmental and economic profile, 2002-04: Japan	366
3.15.2. National agri-environmental performance compared to the OECD average.	373
3.15.3. National water retaining capacity of agriculture.	373
3.15.4. Share of eco-farmers in the total number of farmers.	373
3.16.1. National agri-environmental and economic profile, 2002-04: Korea	377
3.16.2. National agri-environmental performance compared to the OECD average.	383
3.16.3. Composition of soils	383
3.16.4. National water retaining capacity of agriculture.	383
3.17.1. National agri-environmental and economic profile, 2002-04: Luxembourg	386
3.17.2. National agri-environmental performance compared to the OECD average.	391
3.17.3. Nitrate and phosphorus concentration in river sampling stations.	391
3.17.4. Agricultural land under agri-environmental schemes	391
3.18.1. National agri-environmental and economic profile, 2002-04: Mexico	393
3.18.2. National agri-environmental performance compared to the OECD average.	399

3.18.3. Trends in key agri-environmental indicators	399
3.18.4. Trends in key agri-environmental indicators	399
3.19.1. National agri-environmental and economic profile, 2002-04: Netherlands	402
3.19.2. National agri-environmental performance compared to the OECD average	409
3.19.3. Annual mean concentrations of nitrogen and phosphorus in surface water of rural and agricultural water catchments	409
3.19.4. Farmland bird populations	409
3.20.1. National agri-environmental and economic profile, 2002-04: New Zealand	413
3.20.2. National agri-environmental performance compared to the OECD average	420
3.20.3. Sectoral use of pesticides: 2004	420
3.20.4. Dairy cattle enteric methane emissions per litre of milk	420
3.21.1. National agri-environmental and economic profile, 2002-04: Norway	423
3.21.2. National agri-environmental performance compared to the OECD average	430
3.21.3. National sales of pesticides	430
3.21.4. Net change in agricultural land for five counties	430
3.22.1. National agri-environmental and economic profile, 2002-04: Poland	433
3.22.2. National agri-environmental performance compared to the OECD average	444
3.22.3. Agriculture and forest land at risk to erosion	444
3.22.4. Index of population trends of farmland birds	444
3.23.1. National agri-environmental and economic profile, 2002-04: Portugal	448
3.23.2. National agri-environmental performance compared to the OECD average	456
3.23.3. Numbers of local breeds under <i>in situ</i> conservation programmes: 2006	456
3.23.4. Relation between land use and Designated Nature Conservation Areas (DNCA): 2004	456
3.24.1. National agri-environmental and economic profile, 2002-04: Slovak Republic . .	459
3.24.2. National agri-environmental performance compared to the OECD average	468
3.24.3. Agricultural methane (CH ₄) and nitrous oxide (N ₂ O) emissions	468
3.24.4. Share of agricultural land under different types of protected areas: 2003	468
3.25.1. National agri-environmental and economic profile, 2002-04: Spain	472
3.25.2. National agri-environmental performance compared to the OECD average	482
3.25.3. Area of organic farming	482
3.25.4. Share of Dehesa area in total land area for five regions	482
3.26.1. National agri-environmental and economic profile, 2002-04: Sweden	486
3.26.2. National agri-environmental performance compared to the OECD average	494
3.26.3. Losses of nutrients from arable areas and the root zone	494
3.26.4. Cultural features on arable land	494
3.27.1. National agri-environmental and economic profile, 2002-04: Switzerland	498
3.27.2. National agri-environmental performance compared to the OECD average	504
3.27.3. Support for agricultural semi-natural habitats	504
3.27.4. Input/output efficiency of nitrogen, phosphorous and energy in agriculture . . .	504
3.28.1. National agri-environmental and economic profile, 2002-04: Turkey	507
3.28.2. National agri-environmental performance compared to the OECD average	518
3.28.3. Trends in key agri-environmental indicators	518
3.28.4. Trends in key agri-environmental indicators	518
3.29.1. National agri-environmental and economic profile, 2002-04: United Kingdom	522
3.29.2. National agri-environmental performance compared to the OECD average	528

3.29.3. Agri-environmental trends	528
3.29.4. Greenhouse gas emission trends and projections.	528
3.30.1. National agri-environmental and economic profile, 2002-04: United States.	532
3.30.2. National agri-environmental performance compared to the OECD average.	540
3.30.3. Soil erosion on cropland	540
3.30.4. Change in palustrine and estuarine wetlands on non-federal land and water area	540
3.31.1. National agri-environmental and economic profile, 2002-04: European Union (15)	545
3.31.2. EU15 agri-environmental performance compared to the OECD average.	548
3.31.3. Agri-environmental trends, EU15	548
3.31.4. Agri-environmental trends, EU15	548

This book has...



StatLinks 

**A service that delivers Excel® files
from the printed page!**

Look for the *StatLinks* at the bottom right-hand corner of the tables or graphs in this book. To download the matching Excel® spreadsheet, just type the link into your Internet browser, starting with the <http://dx.doi.org> prefix.

If you're reading the PDF e-book edition, and your PC is connected to the Internet, simply click on the link. You'll find *StatLinks* appearing in more OECD books.



From:
**Environmental Performance of Agriculture in
OECD Countries Since 1990**

Access the complete publication at:
<https://doi.org/10.1787/9789264040854-en>

Please cite this chapter as:

OECD (2008), "OECD Country Trends of Environmental Conditions related to Agriculture since 1990: Finland", in *Environmental Performance of Agriculture in OECD Countries Since 1990*, OECD Publishing, Paris.

DOI: <https://doi.org/10.1787/9789264040854-12-en>

This work is published under the responsibility of the Secretary-General of the OECD. The opinions expressed and arguments employed herein do not necessarily reflect the official views of OECD member countries.

This document and any map included herein are without prejudice to the status of or sovereignty over any territory, to the delimitation of international frontiers and boundaries and to the name of any territory, city or area.

You can copy, download or print OECD content for your own use, and you can include excerpts from OECD publications, databases and multimedia products in your own documents, presentations, blogs, websites and teaching materials, provided that suitable acknowledgment of OECD as source and copyright owner is given. All requests for public or commercial use and translation rights should be submitted to rights@oecd.org. Requests for permission to photocopy portions of this material for public or commercial use shall be addressed directly to the Copyright Clearance Center (CCC) at info@copyright.com or the Centre français d'exploitation du droit de copie (CFC) at contact@cfcopies.com.